

Northwest Forest Plan – The First 15 Years [1994-2008]

Watershed Condition Status and Trend

Steven H. Lanigan, Sean N. Gordon, Peter Eldred, Mark Isley,
Chris Moyer, Steve Wilcox, and Heidi Andersen

Abstract

Lanigan, Steven H.; Gordon, Sean N.; Eldred, Peter; Isley, Mark; Wilcox, Steve; Moyer, Chris;

Andersen, Heidi. 2011. Northwest Forest Plan – the first 15 years (1994-2008): status and trend of watershed condition. Gen. Tech. Rep. PNW-GTR-XXX. Portland, OR:U.S. Department of Agriculture, Forest Service, Pacific Northwest Research Station. Xxx p.

We used two different data sets to evaluate stream and watershed condition for sixth-field watersheds in each aquatic province within the Northwest Forest Plan (NWFP) area. A stream evaluation was based on inchannel data (e.g., substrate, pieces of large wood, water temperature, pool frequency, and macroinvertebrates) we sampled from 2002 to the 2009 (193 watersheds) as part of a repeating sample design. We just completed our first round of sampling, so only current condition was calculated for this data set. Grouping condition scores for the inchannel data into categories, relatively few fell into the low (10 percent) and very low (1 percent) categories. The majority of inchannel attribute scores fell into the moderate (35 percent) and high (41 percent) condition ranges, with relatively few (12 percent) in the very high category. For low-scoring watersheds, water temperature was often the most influential factor. Aquatic invertebrate scores also appeared influential in producing the low scores. An evaluation of upslope and riparian (watershed-wide) conditions for all 1,379 sixth-field watersheds in the NWFP area with significant federal ownership was based on mapped data, e.g., road density, based on U.S. Forest Service and Bureau of Land Management geographic information system road layers, and vegetation data, e.g., tree canopy cover, derived from satellite imagery. Watershed-wide condition scores were calculated for 1994 and 2008, and the difference in these scores was used to represent trend. Regarding status, the overall condition scores of the 1,379 watersheds mostly fell into the low (21 percent), moderate (27 percent) high (26 percent), and very high (22 percent) categories; relatively few watersheds scored in the very low (4 percent) category. The majority of watersheds (69 percent) had a positive change in

THE FINDINGS AND CONCLUSIONS IN THIS REPORT ARE IN PRESS AND SUBJECT TO CHANGE PRIOR TO FORMAL DISSEMINATION BY THE AGENCIES AND SHOULD NOT BE CONSTRUED TO REPRESENT AGENCY DETERMINATION OR POLICY.

condition scores (trend). Of those with larger positive changes, most were driven by both improvements in road (decommissioning) and vegetation (natural growth) scores. The greatest negative score changes were caused by the Biscuit Fire and other fires along the eastern side of the Cascades. Half of the fire-affected watersheds were in congressional reserves, 35 percent in late-successional reserves, and 15 percent in matrix (lands identified for timber production).

Keywords: Effectiveness monitoring, status and trend monitoring, aquatic ecosystems, riparian ecosystems, watersheds, decision-support models, Northwest Forest Plan, aquatic conservation strategy, Pacific Northwest.

Authors

Steve Lanigan is the module leader, **Peter Eldred** is the geographic information system (GIS) coordinator, **Mark Isley** is the database manager, **Steve Wilcox** is the GIS cartographer, and **Heidi Andersen** is a fish biologist for the Aquatic and Riparian Effectiveness Monitoring Program, U.S. Department of Agriculture, Forest Service, Pacific Northwest Region, 333 NW First, Portland, OR 97204; **Chris Moyer** is a fish biologist for the Aquatic and Riparian Effectiveness Monitoring Program, U.S. Department of the Interior, Bureau of Land Management, Oregon State Office, 4077 Research Way, Corvallis, OR 97333; **Sean Gordon** is a research associate with the Dept. Forest Engineering, Resources and Management, Oregon State University, 333 NW First, Portland, OR 97204.

Executive Summary

The watershed monitoring module (also known as the Aquatic and Riparian Effectiveness Monitoring Program or AREMP) determines if the Northwest Forest Plan's (NWFP) aquatic conservation strategy is achieving the goals of maintaining and restoring the condition of watersheds. The NWFP area being evaluated includes USDA Forest Service (FS), USDI Bureau of Land Management (BLM), and USDI National Park Service (NPS) lands. Only the federal portion of sixth-field watersheds was included when determining watershed condition status and trend because federal agency land managers have no jurisdiction over nonfederal lands.

We used two different data sets to evaluate stream and watershed condition for each aquatic province within the NWFP. A stream evaluation was based on inchannel data (e.g., substrate, pieces of large wood, water temperature, pool frequency, and macroinvertebrates) we sampled from 2002 to 2009 (193 watersheds) as part of a repeating sample design. We just completed our first round of sampling, so only current condition was calculated for this data set. An evaluation of upslope and riparian (watershed-wide) conditions for all 1,379 sixth-field watersheds in the NWFP area with significant federal ownership was based on mapped data, e.g., road density, based on FS and BLM geographic information system road layers, and vegetation data, e.g., tree canopy cover, derived from satellite imagery. Watershed-wide condition scores were calculated for 1994 and 2008, and the difference in these scores was used to represent trend. Experts from six aquatic provinces decided which indicators to use, and how to evaluate and combine them into an overall condition index for each level. We used decision-support modeling software to calculate the index scores for each watershed to a standardized range between -1 ("poor") and +1 ("good").

Grouping condition scores for the inchannel data into categories, relatively few fell into the low (10 percent) and very low (1 percent) categories. The majority of inchannel attribute scores fell into the

THE FINDINGS AND CONCLUSIONS IN THIS REPORT ARE IN PRESS AND SUBJECT TO CHANGE PRIOR TO FORMAL DISSEMINATION BY THE AGENCIES AND SHOULD NOT BE CONSTRUED TO REPRESENT AGENCY DETERMINATION OR POLICY.

moderate (35 percent) and high (41 percent) condition ranges, with relatively few (12 percent) in the very high category. For low-scoring watersheds, water temperature was often the most influential factor. In many of the provincial evaluation models, a poor water temperature score carried more weight than other attributes because it was only measured once for each watershed (at the lowest point), in contrast to the other attributes, which were averaged over 6 to 8 sites. Aquatic invertebrate scores also appeared influential in producing the low scores.

Regarding status, the overall condition scores of the 1,379 watersheds were clustered in the center of the distribution and skewed slightly positive. Most fell into the low (21 percent), moderate (27 percent) high (26 percent), and very high (22 percent) categories; relatively few watersheds scored in the very low (4 percent) category. The spatial distribution of watershed scores showed some noticeable patterns. High scores were found in the central Olympic Peninsula (Olympic National Park), the north central Cascades, the Oregon Coast Range, and scattered pockets along the Klamath-Siskiyou mountain range (mostly corresponding to designated wilderness areas). Low condition scores were present in the southern Olympic region, eastern Klamath-Siskiyou, and along the eastern and western flanks of the Cascade Range in Oregon and Washington. These low-scoring areas are generally closer to existing development and lower in elevation and slope, making them historically more accessible to roading and timber harvest. Watershed condition was most positive for congressionally reserved lands, followed by late-successional reserves (LSR), and then matrix lands. Key watersheds, which provide high-quality habitat or refugia for aquatic- and riparian-dependent species, or would be able to after restoration, were in better condition than nonkey watersheds in both 1994 and 2008.

The majority of watersheds (69 percent) had a positive change in condition scores (trend). By far the largest trend category was score increases between 0 and +0.1 (55 percent). This trend was largely due to small increases in vegetation scores from natural tree growth moving acres out of early seral classes or into late seral classes. The second largest trend category was a minor decrease between 0 and -0.1 (18

THE FINDINGS AND CONCLUSIONS IN THIS REPORT ARE IN PRESS AND SUBJECT TO CHANGE PRIOR TO FORMAL DISSEMINATION BY THE AGENCIES AND SHOULD NOT BE CONSTRUED TO REPRESENT AGENCY DETERMINATION OR POLICY.

percent). Because there has been little road building on federal lands, this trend was again due to vegetation, but in this case, losses in average tree diameter or canopy cover. Positive trends in the +0.1 to +0.3 range were mostly due to road decommissioning (42%) but with a fair contribution from vegetation (30%) and landslide risk (28%). Reduced landslide risk was the dominant driver of improvement about the +0.3 level (54%), apparently due to road decommissioning in landslide prone areas, since road improvements contributed considerably more (38%) than vegetation (8%) at this level. The greatest negative score changes were caused by the Biscuit Fire and other fires along the eastern side of the Cascades. Half of the fire-affected watersheds were in congressional reserves, 35 percent in late-successional reserves, and 15 percent in matrix (lands identified for timber production). Looking at changes by land use allocations, the reserved class actually declined slightly (-0.01), while the LSR (+0.05) and matrix (+0.04) both showed slight increases on average. Considering that the reserved class is already generally in good condition with respect to roads and vegetation, it is not surprising that it did not increase, and as harvest and road building are not permitted in these areas, the main driver was vegetation losses from fires.

Table of Contents

CHAPTER 1: INTRODUCTION.....	8
CHAPTER 2: METHODS.....	16
CHAPTER 3: RESULTS.....	38
CHAPTER 4: DISCUSSION.....	56
CHAPTER 5: EMERGING ISSUES	64
REFERENCES.....	68
APPENDIX 1: OBJECTIVES OF THE AQUATIC CONSERVATION STRATEGY ¹	83
APPENDIX 2: PROJECT-SCALE RESTORATION EFFORTS.	85
APPENDIX 3: WATERSHEDS SELECTED FOR INCHANNEL MONITORING	86
APPENDIX 4: MODEL STRUCTURES, EVALUATION CRITERIA, AND STATUS AND TREND MAPS FOR EACH OF THE PROVINCIAL DECISION-SUPPORT MODELS	87
APPENDIX 5: ADDITIONAL INFORMATION ON INCHANNEL, UPSLOPE AND RIPARIAN DATA INPUTS.....	92
APPENDIX 6: QUALITY ASSESSMENT OF UPSLOPE/RIPARIAN AND INCHANNEL DATA INPUTS.	101
APPENDIX 7: CONTACT INFORMATION	105
TABLES	106

THE FINDINGS AND CONCLUSIONS IN THIS REPORT ARE IN PRESS AND SUBJECT TO CHANGE PRIOR TO FORMAL DISSEMINATION BY THE AGENCIES AND SHOULD NOT BE CONSTRUED TO REPRESENT AGENCY DETERMINATION OR POLICY.

Chapter 1: Introduction

In 1994, the Northwest Forest Plan (NWFP) Record of Decision amended 19 National Forest and 7 Bureau of Land Management (BLM) and resource plans within the range of the northern spotted owl (*Strix occidentalis caurina*) (USDA and USDI 1994). An interagency effectiveness monitoring framework was implemented to meet requirements for tracking status and trend for watershed condition, late and old forests, social and economic conditions, tribal relationships, and population and habitat for marbled murrelets (*Brachyramphus marmoratus*), and northern spotted owls. Monitoring results are evaluated and reported in 1- and 5-year intervals. Monitoring results for the first 10 years are documented in a series of reports posted at <http://www.reo.gov/monitoring/>.

This “15-year report” evaluates the status of watershed condition and changes in condition under the NWFP aquatic conservation strategy during years 1994–2008. Although this report was originally intended to evaluate 15 years of data, delays in producing this and other NWFP monitoring reports allowed us to include stream condition data from 2009 in our evaluation of stream condition status. Gallo et al. (2005) described the status of aquatic and riparian resources and changes in their condition for the first 10 years under the NWFP aquatic conservation strategy. That assessment was based on evaluating 250 randomly chosen watersheds throughout the NWFP area. Because of the limited number of watersheds, inference was appropriate only for the whole NWFP area. The Regional Executive Interdisciplinary Team (REIT) responded to the Gallo et al. (2005) assessment by asking Aquatic and Riparian Effectiveness Monitoring Program (AREMP) staff to develop a way to evaluate watershed condition at smaller scales than the entire NWFP area. The AREMP staff proposed a “GIS and field monitoring” option that the REIT supported.¹ This option continued the sampling of inchannel attributes in 250 watersheds along with using a geographic information system (GIS) framework to evaluate all

¹ Regional Interagency Executive Committee Meeting Notes for March 17, 2006

watersheds with at least 25 percent of the 1:100,000 stream layer within federal land ownership. This report describes the results of evaluating all watersheds that met this criterion.

Background

In the early 1990s, controversy over harvest in old-growth forests led to sweeping changes in management of federal forests in western Washington, Oregon, and northwest California. These changes were prompted by a series of lawsuits in the late 1980s and early 1990s that effectively shut down federal timber harvest in the Pacific Northwest. In response, President Clinton convened a summit in Portland, Oregon, in 1993, where he issued a mandate for federal land management and regulatory agencies to work together to develop a plan to resolve the conflict. The President's guiding principles followed shortly after the summit in his *Forest Plan for a Sustainable Economy and a Sustainable Environment* (Tuchmann et al. 1996)—otherwise known as the Northwest Forest Plan.

Immediately after the summit, a team of scientists and technical experts were convened to conduct an assessment of options (FEMAT 1993). This assessment provided the scientific basis for the environmental impact statement and record of decision (USDA and USDI 1994) to amend Forest Service and Bureau of Land Management planning documents within the range of the northern spotted owl.

The record of decision (ROD), covering 24 million acres of federal lands, put in place a new approach to federal land management. Key components of the ROD included a new set of land use allocations—late-successional reserves, matrix lands, riparian reserves, adaptive management areas, and key watersheds. The NWFP standards and guidelines provided direction regarding how these land use allocations were to be managed. In addition, the NWFP put in place a variety of strategies and processes to be implemented. These included adaptive management, an aquatic conservation strategy, late-successional reserve and watershed assessments, a survey and manage program, an interagency executive organization, social and economic mitigation initiatives, and monitoring.

Monitoring provides a means to address the uncertainty of our predictions and compliance with forest management laws and policy. The ROD stated that monitoring is essential and required:

Monitoring is an essential component of the selected alternative. It ensures that management actions meet the prescribed standards and guidelines and that they comply with applicable laws and policies. Monitoring will provide information to determine if the standards and guidelines are being followed, verify if they are achieving the desired results, and determine if underlying assumptions are sound. [USDA and USDI 1994]

Judge Dwyer reinforced the importance of monitoring in his 1994 decision declaring the NWFP legally acceptable: “Monitoring is central to the [NWFP’s] validity. If it is not funded, or not done for any reason, the plan will have to be reconsidered” (Dwyer 1994).

The ROD monitoring plan provided a general framework to begin development of an interagency monitoring program. It identified key areas to monitor, initial sets of questions, types and scope of monitoring, the need for common protocols and quality assurance, and the need to develop a common design framework. In 1995, the effectiveness monitoring program plan (Mulder et al. 1995) and initial protocols for implementation monitoring (Alegria et al. 1995) were approved by the Regional Interagency Executive Committee. Approval of the effectiveness monitoring plan led to the formation of technical teams to develop the overall program strategy and design (Mulder et al. 1999) and monitoring protocols for late-successional and old-growth forests (termed older forests) (Hemstrom et al. 1998), northern spotted owls (Lint et al. 1999), marbled murrelets (Madsen et al. 1999), tribal relations (Bown and others 2002), and watershed condition (Reeves et al. 2004). Periodic analysis and interpretation of monitoring data is essential to completing the monitoring task critical to completing the adaptive management cycle. This important step was described in the overall monitoring strategy (Mulder et al. 1999) and approved by the Regional Interagency Executive Committee.

THE FINDINGS AND CONCLUSIONS IN THIS REPORT ARE IN PRESS AND SUBJECT TO CHANGE PRIOR TO FORMAL DISSEMINATION BY THE AGENCIES AND SHOULD NOT BE CONSTRUED TO REPRESENT AGENCY DETERMINATION OR POLICY.

Gallo et al. (2005) completed one of a series of assessments describing current status and trends for the first 10 years of the NWFP, which also included northern spotted owls (Lint 2005), older forests (Moeur et al. 2005), marbled murrelets (Huff et al. 2006), socioeconomic conditions (Charnley 2006), tribal relations (Stewart and Martine 2006), and implementation or compliance monitoring (Baker et al., 2005). This series of reports was accompanied by a synthesis report by a panel of scientists and managers that integrates and interprets the findings from the status and trend reports and offers alternatives to policymakers (Haynes et al. 2006). These “10-year reports” were the first comprehensive analysis and interpretation of monitoring data since the ROD.

Overview of the Aquatic Conservation Strategy

The aquatic conservation strategy (ACS) is a comprehensive, regionwide strategy designed to maintain, restore, and protect those processes and landforms that create good ecological conditions in watersheds, such as high-quality habitat for aquatic and riparian organisms and good water quality (FEMAT 1993). The strategy contains nine objectives that describe general characteristics of functional aquatic and riparian ecosystems that are intended to maintain and restore good habitat in the context of ecological disturbance (see appendix 1). This approach was intended to prevent further degradation of aquatic ecosystems and restore habitat over broad landscapes, as opposed to focusing on individual projects or species (USDA and USDI 1994). Aquatic and riparian organisms evolved in a dynamic environment influenced by natural disturbance. The authors of the strategy believed that stewardship of aquatic resources is most likely to protect biological diversity and productivity when land use activities do not substantially alter the natural disturbance regime to which organisms are adapted (FEMAT 1993). Therefore, the strategy used several tactics to try to maintain the natural disturbance regime in watersheds. The strategy also includes standards and guidelines that apply to management activities in riparian reserves and key watersheds. The four components of the strategy were intended to work in concert to maintain and restore the health of aquatic and riparian ecosystems:

THE FINDINGS AND CONCLUSIONS IN THIS REPORT ARE IN PRESS AND SUBJECT TO CHANGE PRIOR TO FORMAL DISSEMINATION BY THE AGENCIES AND SHOULD NOT BE CONSTRUED TO REPRESENT AGENCY DETERMINATION OR POLICY.

- Watershed analysis—used to characterize watersheds and provide a basis (context) for making management decisions.
- Riparian reserves—used to enhance habitat for riparian-dependent organisms, to provide good water quality, to provide dispersal corridors for terrestrial species, and to provide connectivity within watersheds.
- Key watersheds—provide high-quality habitat or refugia for aquatic- and riparian-dependent species, or would be able to after restoration.
- Watershed restoration—designed to recover degraded habitat and maintain existing good conditions.

Although late-successional reserves are not listed among the components of the strategy, they provide increased protection for aquatic and riparian ecosystems. Late-successional reserves contain areas of high-quality stream habitat that serve as refuge for aquatic and riparian organisms and as source areas from which organisms may move to recolonize formerly degraded areas (USDA and USDI 1994).

The aquatic conservation strategy for the Forest Service in Region 6 (Oregon and Washington) has evolved to become known as the Aquatic and Riparian Conservation Strategy (ARCS).² The ARCS integrates and refines three earlier aquatic strategies in the region: the aquatic conservation strategy of the NWFP, PACFISH,³ and INFISH.⁴ It maintains and builds upon the essential components of these earlier strategies including: riparian management areas, key watersheds, watershed analysis, watershed restoration, and monitoring. It is based on the best available science, is ecologically sound, and complies

² Staab, B. 2010. Personal communication, Hydrologist, USDA Forest Service, Pacific Northwest Region, 333 SW First Ave., Portland, OR 97204.

³ Interim Strategies for Managing Anadromous Fish-Producing Watersheds in Eastern Oregon and Washington, Idaho and portions of California

⁴ Inland Native Fish Strategy

with the requirements of federal law including the National Forest Management Act, Endangered Species Act, and Clean Water Act.

Specific refinements include better recognition that disturbance is integral to the resiliency of ecosystems, acknowledgment that climate change may have large effects on these resources, consideration of scale effects on ecosystem processes, better linkages between management intent and direction, and a more robust and transparent process for selecting key watersheds. In addition, the ARCS now includes a more explicit and strategic approach to executing forest watershed restoration programs.

Monitoring was included in the strategy to achieve three goals: ensure that management actions follow the standards and guidelines and comply with applicable laws and policies (implementation monitoring), determine the effectiveness of management practices at multiple spatial scales ranging from individual watersheds to the entire NWFP area (effectiveness monitoring), and determine whether the assumptions underlying the strategy are sound (validation monitoring). The first goal was accomplished through the implementation monitoring program (Baker et al. 2005). The aquatic and riparian effectiveness monitoring program was developed to reach the effectiveness monitoring goal.

Overview of the Aquatic and Riparian Effectiveness Monitoring Program

The AREMP is responsible for the effectiveness monitoring component of the ACS. Its purpose is to assess the effectiveness of the NWFP by periodically determining the status of watershed condition and using this information to track trends in the condition of watersheds through time. Watershed condition refers to a combination of aquatic, riparian, and upslope characteristics. While the original intent of the AREMP was to combine all these characteristics into a single watershed evaluation (Reeves et al. 2004), the evaluation process has evolved to consider stream condition separately from upslope and riparian (“watershed”) condition. Stream condition is based on inchannel data, e.g., substrate, pieces of large wood, water temperature, pool frequency, and macroinvertebrates. Watershed condition is based on mapped data, e.g., road density and vegetation data.

THE FINDINGS AND CONCLUSIONS IN THIS REPORT ARE IN PRESS AND SUBJECT TO CHANGE PRIOR TO FORMAL DISSEMINATION BY THE AGENCIES AND SHOULD NOT BE CONSTRUED TO REPRESENT AGENCY DETERMINATION OR POLICY.

Stream and watershed condition are determined by integrating biological and physical indicator information (Reeves et al. 2004). The results are presented in the form of a distribution of the condition of streams and watersheds across the NWFP area and not on the condition of individual streams and watersheds. If the NWFP is effective, the distribution of conditions should either stay the same or improve over time (Reeves et al. 2004). Note that the Forest Ecosystem Management Assessment Team (FEMAT) scientists did not intend for each of the ACS's objectives to be monitored individually, nor did they expect that the objectives would be met in each watershed at all times.

The AREMP was pilot tested in 2000 and 2001 to evaluate sampling protocols and determine the funding and crew structure needed to implement the monitoring program (Figure 1). Monitoring officially began in 2002, although funding was about half the amount identified as being needed to fully implement the program. As of fall 2009, 193 of an expected 250 watersheds were sampled for inchannel attributes.

Figure 1—Timeline of the monitoring program development and implementation.

Evaluating the effectiveness of the ACS is based on measuring changes in the distribution of stream and watershed condition scores through time. Few details on the changes in individual watersheds are provided. The ACS does not describe the baseline condition of streams and watersheds, nor does it define a desired distribution. We infer that if the strategy has been effective in maintaining or improving the condition of watersheds, then the distribution of stream and watershed condition scores should shift in a direction that indicates improvement (Reeves et al. 2004). To spotlight some of the success local units have achieved with project-scale restoration, we describe several case studies in appendix 2. Some of the projects may have had immediate effects, such as opening up habitat to fish by replacing poorly designed culverts that previously blocked fish passage. But most restoration projects should be viewed as a critical first step in restoring natural watershed processes.

Monitoring Questions

The AREMP is charged with answering questions related to evaluating the effectiveness of the aquatic conservation strategy in achieving its goal of maintaining and improving the condition of watersheds in the NWFP area (Reeves et al. 2004). This report focuses on responding to three questions, whose answers provide insight for evaluating the success of the aquatic conservation strategy:

1. What is the status of inchannel conditions?
2. What is the status of upslope and riparian conditions?
3. What is the trend in watershed conditions?

Chapter 2: Methods

Each of the three principal monitoring questions is answered using somewhat different data sources and methods. First we describe the common elements of the study area and the conceptual model before moving onto sections providing more details on study designs, data sources, and analytical procedures for each of the four principal monitoring questions.

Study Area

The Northwest Forest Plan (NWFP) encompasses more than 24 million acres of federal lands in western Washington, western Oregon, and northwestern California and includes the entire geographic range of the northern spotted owl (Figure 2). Stream and riparian habitat conditions differ greatly across the NWFP area because of natural and management-related factors. Geologic and climatic history influence topographic relief, landforms, channel patterns, and the dominant erosion processes. Precipitation ranges from more than 200 inches per year in some areas near the coast to less than 20 inches on the east side of the Cascade Range. Riparian vegetation communities are structured by climate and the disturbance regime of the area, including hydrologic processes and disturbance such as forest fires region (Benda et al. 1998, Naiman et al. 1992). Many of these critical components of landscape form and function are in distinctive combinations characteristic of each physiographic province in the region. Physiographic provinces incorporate physical, biological, and environmental factors that shape broad-scale landscapes and therefore reflect differences in responses such as soil development and plant community structure.

The evaluation of upslope and riparian conditions in watersheds was tailored to specific physiographic provinces. Although physiographic provinces are useful in describing both terrestrial and aquatic ecosystems, different processes dominate the functioning of these ecosystems. Consequently, the Forest Ecosystem Management Assessment Team (FEMAT 1993) used different physiographic province

boundaries for aquatic and terrestrial ecosystems. The physiographic boundaries used in this analysis were developed from those used in the aquatic ecosystem assessment, and are based on broadly drawn precipitation and geologic zones, as well as political boundaries (state lines). These province boundaries differ from those used by the other effectiveness monitoring components (e.g., the late-successional old-growth and the northern spotted owl), which were delineated primarily by vegetation type and political boundaries. The aquatic province boundaries used by the FEMAT (1993) were not available in a digital format, so their province boundary lines were refined by using level-four lines described by Omernik in Oregon and Washington (Bryce et al. 1999), Bailey ecological subsections lines in California (Bailey et al. 1994), and the Cascade crest derived from the Forest Service Pacific Northwest Region sixth-field watershed layer.

The NWFP area contains eight aquatic physiographic provinces including the Olympic Peninsula, North Cascades, Willamette/Puget Trough, West Cascades, Washington/Oregon Coast, High Cascades, Klamath/Siskiyou, and Franciscan. Land ownership in the Willamette/Puget Trough is predominantly private, and none of the watersheds in this province met the monitoring program minimum criterion of at least 25 percent of the 1:100,000 stream layer within federal land ownership. Consequently, this province is not included in the analysis. The Klamath/Siskiyou and Franciscan Provinces were combined into one provincial area for this report, along with the California portion of what was formally part of the High Cascades. Descriptions of the provinces based largely on those presented by FEMAT (1993) are provided in Gallo et al. (2005).

The subwatershed (sixth-field hydrological unit, hereafter called watershed) was chosen as the basic geographic unit for monitoring, as recommended by Reeves et al. (2004), because it was the smallest consistently delineated unit available at the time. These watersheds are 10,000 to 40,000 acres, and include both complete (contains all headwaters for a main stream) and composite (contains only part of the source waters) watersheds. Because the NWFP applies only to federally managed lands, watersheds

THE FINDINGS AND CONCLUSIONS IN THIS REPORT ARE IN PRESS AND SUBJECT TO CHANGE PRIOR TO FORMAL DISSEMINATION BY THE AGENCIES AND SHOULD NOT BE CONSTRUED TO REPRESENT AGENCY DETERMINATION OR POLICY.

must contain a minimum of 25 percent of the total length of the stream (1:100,000 National Hydrography Dataset stream layer) within federal ownership (USDA Forest Service [FS], USDI Bureau of Land Management [BLM], or USDI National Park Service [NPS]) to be considered for sampling and analysis in the monitoring program. The ownership criterion was recommended by Reeves et al. (2004) to gauge the influence of the strategy by sampling watersheds in which the strategy was implemented to varying degrees while avoiding sampling watersheds in which the contribution of federal lands to the condition of the watershed was insignificant. The NWFP area contains 2,657 watersheds, of which 2,151 contain some land that is federally owned, and 1,379 have at least 25 percent of stream channels in federal ownership. The ownership criterion excludes about 7 percent of the federal lands in the NWFP area from this analysis.

Only the federal portion of watersheds was included when determining watershed condition status and trend because federal agency land managers have no jurisdiction over management of nonfederal lands.

Figure 2—Aquatic provinces used to assess watershed condition in the Northwest Forest Plan (NWFP) area. The NWFP area extends from the U.S-Canada border to Point Reyes, California, and includes the eastern flank of the Cascade Mountain range and encompasses the range of the northern spotted owl. The NWFP area being evaluated includes USDA Forest Service, USDI Bureau of Land Management, and USDI National Park Service lands.

Land Use Categories

Land use categories provide a key spatial component of the NWFP by assigning different management guidelines and priorities to zones within the NWFP area. We review our three monitoring questions in the context of two types of land classification: the general NWFP land use allocations (reserved, late-successional reserve, matrix) and the NWFP aquatic conservation strategy (ACS) designations of key versus nonkey watersheds. We collapsed some of the land use allocations that have

THE FINDINGS AND CONCLUSIONS IN THIS REPORT ARE IN PRESS AND SUBJECT TO CHANGE PRIOR TO FORMAL DISSEMINATION BY THE AGENCIES AND SHOULD NOT BE CONSTRUED TO REPRESENT AGENCY DETERMINATION OR POLICY.

similar guidelines for management (Table 1). The land use allocation categories presented here are the same as those described by Tuchmann et al. (1996). Boundaries for land use categories did not follow watershed boundaries; consequently multiple land use categories may have been present in individual watersheds. Within each classification, each watershed was assigned to the class covering the largest amount of its area. Table 1 shows the number of watersheds falling into each class. The following paragraphs briefly describe each allocation, and their spatial distribution is displayed in Figure 3.

Congressional Reserves (CR)—Congressional reserves include national parks and monuments, wilderness areas, wildlife refuges, and other areas reserved by the administrative unit or act of Congress. These lands are generally not managed for timber production.

Late-successional and old-growth reserves (LSR)—Late-successional reserves contain largely old-growth forest and were designated to provide habitat for old-growth-dependent species such as the northern spotted owl. Adaptive management areas managed under LSR guidelines were grouped into LSR (see below).

Matrix—The matrix land use allocation includes all lands not included in one of the other allocations. Scheduled timber harvest activities may take place in matrix lands. For analysis and reporting purposes, we grouped some adaptive management with matrix (see below). Riparian reserves were not included as a separate land allocation because they have not been mapped; they are included as part of the matrix land allocation. Results for upslope and riparian attributes are based on the riparian buffers described in chapter 2.

Adaptive management areas—Ten areas were identified to develop and test innovative management approaches to integrate and achieve ecological, economic, and other social and community objectives (USDA and USDI 1994). They are a mix of lands where timber production can occur and where timber production must follow LSR guidelines. For analysis and reporting purposes, we grouped

watersheds in adaptive management areas into either matrix lands or LSR, depending on which allocation covered the largest amount of its area.

Key and nonkey watersheds—Key watersheds are one of the primary components of the aquatic conservation strategy. They are intended to “serve as refugia for aquatic organisms, particularly in the short term for at-risk fish populations, to have the greatest potential for restoration, or to provide sources of high-quality water” (Haynes et al. 2006). Key watersheds were identified as part of the ACS and independent of the land use allocations in the NWFP, thus key and nonkey watershed designations overlay the other land use allocations. Key watershed delineation was begun prior to the development of the interagency standard 5th and 6th field watershed boundaries, so their boundaries are not always coincident. For this analysis, 469 of our 1,379 watersheds are considered key because they have >50 percent of the area designated as key watershed. The remaining 910 watersheds are considered as nonkey in this assessment.

Table 1—Collapsed land use and watershed categories used in this analysis.

Figure 3—Land use allocations in the Northwest Forest Plan (NWFP) area.

Study Design

Definition of Watershed Condition

The definition of watershed condition developed by the monitoring program is based on the goals of the NWFP aquatic conservation strategy (see app. 1) and on guidance provided by the aquatic monitoring plan (Reeves et al. 2004). The NWFP was designed to account for the complex and dynamic nature of aquatic ecosystems resulting from the wide range of physical characteristics, natural disturbance events, and climatic features of the region (Benda et al. 1998, Naiman et al. 1992). Monitoring these dynamic watershed processes is accomplished by linking them to measurable physical attributes (e.g., vegetation structure, road density, water temperature). Reeves et al. (2004) initially identified 90 potential attributes

that represent key functions and processes in watersheds. This number of attributes was reduced based on criteria established by Noon et al. (1999). The monitoring program further removed some attributes that were found not to produce useful or consistent information (Lanigan et al. 2007). The remaining attributes represent upslope, riparian, and inchannel processes (Table 2).

Table 2—Attributes included in the watershed condition assessments listed by ecological process.

The condition of a watershed is defined as “good” if the state of these attributes support a high diversity and abundance of aquatic and riparian species. Many of the physical indicators are chosen for their relevance to native or desired fish species because of these species’ roles in driving management policies (including the NWFP itself) and the availability of research related to their habitat needs. However, we attempt to assess indicators relative to the natural potential of the site to provide biotic habitat. A watershed that naturally does not support fish populations (because of elevation or other natural conditions) but has little vegetation disturbance, few roads, good pools, and wood should be evaluated positively. If this watershed loses significant vegetation, even from natural causes (e.g., fire), then the condition rating will go down (it is below its potential).

This simplified view of condition is a consequence of the fact that indicators taken at one point in time are imperfect measures for dynamic processes. Even a watershed with intact processes may not be in good condition in terms of providing quality fish habitat at any single assessment period. A fundamental principle underlying the monitoring program is that watersheds are naturally dynamic systems. Individual watersheds will cycle through conditions of high and low habitat quality, and not all watersheds can be expected to be in good condition at any one time (Naiman et al. 1992, Reeves et al. 1995). Therefore, the most important product of the monitoring program is the overall **distribution** of individual watershed ratings in the NWFP area. Implementing the ACS should result in an overall distribution of watershed condition scores that improves over time.

Assessment of Watershed Condition

To assess watershed condition for the NWFP, Reeves et al. (2004) proposed a strategy which used the watershed concept as a unit of analysis, integrated multiple indicators, and provided a representative sample of the NWFP area. Because the multiple aspects related to watershed condition are diffuse and many of the relationships between them not yet quantified statistically, they recommended a knowledge-based systems methodology. This type approach enables the integration of quantitative information and more qualitative expert knowledge into an explicit computer model, which then facilitates and documents the consistent and transparent application of a methodology for evaluation. An assessment of management options for federal lands in the interior Columbia River basin took a similar approach, supplementing a expert group approach (Sedell et al. 1997) with computer models that encapsulated expert knowledge and assessed each watershed unit in a consistent manner (Rieman et al. 2000, 2001).

To implement this knowledge-based systems approach, AREMP selected the Ecosystem Management Decision Support System software (EMDS; Reynolds et al. 2002) as the modeling platform. EMDS integrates knowledge-based and decision-modeling components into the ArcGIS software (ESRI, <http://www.esri.com>). The AREMP assessments use the Netweaver component (Rules of Thumb Inc., <http://rules-of-thumb.com/>), which combines a basic multi-attribute decision analysis framework with a variety of mathematical and logical operators. Multiple criteria evaluation is derived from multi-attribute utility theory, which encompasses a suite of techniques for integrating multiple metrics into a combined decision or index value (Keeney and Raiffa 1976). These techniques are designed to address two basic problems with combining indicators, which are often 1) measured in different units, and 2) of different importance to the overall evaluation objective. The sidebar “Overview of the Watershed Evaluation Modeling Process” (p. XX) provides details on how the individual indicators (or attributes) were normalized to a common scale and combined into an overall score.

Sidebar 1—Overview of the Watershed Evaluation Modeling Process

THE FINDINGS AND CONCLUSIONS IN THIS REPORT ARE IN PRESS AND SUBJECT TO CHANGE PRIOR TO FORMAL DISSEMINATION BY THE AGENCIES AND SHOULD NOT BE CONSTRUED TO REPRESENT AGENCY DETERMINATION OR POLICY.

For this assessment, models were developed for two scales, inchannel and upslope/riparian, following the process scales defined by the monitoring plan and the datasets available for the different scales.

Upslope and riparian evaluations were combined in one model because they are based on the same data sources and sampling design. Each model comprises three basic elements: a list of measurable watershed attributes to evaluate, criteria for rating each attribute, and a model structure, which defines how the attribute scores are aggregated into an overall score. Data from each watershed are run through the appropriate provincial models (inchannel and upslope/riparian) to produce scores on a standardized scale from +1 to -1, where +1 indicates excellent condition and -1 indicates poor condition.

For this assessment, decision-support models developed for each aquatic province (n = 6) for the 10-year report (Gallo et al. 2005) were refined during workshops attended by local agency professionals (Figure 4). Separate models were built for each aquatic province (Figure 2) to account for the ecological differences between provinces. The workshops consisted of a semi-structured group process through which participants reviewed the existing model structures, data attributes, and evaluation criteria, and came to consensus on changes needed. After the workshops, models were built and run and the results returned to the workshop participants. A second round of workshops was held to review the preliminary results. Participants compared the results of the models to their knowledge of the condition of the watersheds and suggested refinements to the models as necessary. These changes were made, resulting in the final models used. Generic model diagrams are presented in Figure 5 to illustrate how individual indicators were aggregated into overall condition scores; actual model diagrams and evaluation criteria for each provincial model are provided in appendix 4.

Figure 4—Program personnel from FS, BLM, NPS, Pacific Northwest Research Station, Washington Department of Fish and Wildlife, Washington Department of Ecology, Washington Forest Practices Board Cooperative Monitoring, Evaluation, and Research Committee, Environmental Protection Agency, National Oceanic and Atmospheric Administration Fisheries, Oregon Fish and Wildlife, and California Fish and Game provided technical expertise and local knowledge for decision-

THE FINDINGS AND CONCLUSIONS IN THIS REPORT ARE IN PRESS AND SUBJECT TO CHANGE PRIOR TO FORMAL DISSEMINATION BY THE AGENCIES AND SHOULD NOT BE CONSTRUED TO REPRESENT AGENCY DETERMINATION OR POLICY.

support model refinement during a series of aquatic province workshops. Specialists also provided feedback on how well the output from refined models matched up with their perspective of 'on-the-ground' conditions, so the models could be further fine-tuned.

Figure 5—Illustrative model diagrams showing how indicators are combined into overall condition scores (actual structures for each province detailed in appendix 4).

Monitoring Questions

Data from two scales were used to answer the key monitoring questions, which led to differences in specific study designs and attributes evaluated. The inchannel status evaluation (question 1) was based on sampling of stream data (e.g., water temperature, pool frequency, macroinvertebrates) collected in 193 watersheds by AREMP teams from 2002 to 2009. The assessments for questions 2 and 3 were based on watershed-wide mapped data (e.g., road density, canopy cover) derived from satellite imagery and other corporate data sets (circa 1994 and 2008, 1,379 watersheds). The following sections describe the specific study design and attributes evaluated for each of the monitoring questions.

1. *What is the status of inchannel conditions?*

Study Design

The AREMP study design identified 250 randomly selected watersheds (see app. 3) from the 1,379 watersheds in the sixth-field watershed coverage (version 1.1, dated 2002) for the NWFP area that met both the ownership criterion (i.e., more than 25 percent of the total stream length is located on federal land) and a set of criteria associated with safety and access concerns (see app. 5). Watersheds were selected using generalized random stratified tessellation survey, a process that guarantees a spatially balanced sample (Stevens and Olsen 2003, 2004). According to Reeves et al. (2004), 50 watersheds should be sampled each year for 5 years. On year 6, the watersheds sampled the first year will be revisited. Because of funding limitations, we were only able to sample inchannel attributes in an average of 24 watersheds per year for a combined total of 193 watersheds in 2002 to 2009. Since no repeat

sampling has yet occurred, the inchannel data from all years are used only to represent current status and not trend. Some indicators may be affected by natural year-to-year environmental variations (e.g. the effect of climate on water temperature), which is not accounted for in the current framework but may deserve consideration in future assessments.

Within each watershed selected, inchannel data were collected at multiple sites (stream segments referred to as reaches). These sites were selected by using generalized random stratified tessellation survey (Stevens and Olsen 2003, 2004), the same procedure used to select watersheds. The 1:100,000 National Hydrography Dataset stream layer was clipped to the boundary of each watershed. Sample points were placed on the stream layer within the watershed boundary at random (Figure 6).. These points represented the downstream starting point for the inchannel surveys. The length of the sample reach was determined as 20 times the average bankfull width, with minimum and maximum reach lengths of 175 and 525 yards, respectively. On average, 6 sites were sampled in each watershed, with a range from 4 to 10.

Figure 6—Example of randomly selected sample sites in a sixth-field watershed. The sampled stream reaches (red dots) were selected from 1:100,000 stream layers by using a generalized random stratified tessellation survey, a process that guarantees a spatially balanced sample (Stevens and Olsen 2004).

Attributes

A list of suggested physical, biological, and chemical attributes of stream reaches used for the inchannel evaluation was initially suggested in Reeves et al. (2004) and subsequently refined in the provincial expert workshops. A list of attributes used in this assessment is provided in Table 3; all of these attributes were based on data collected by AREMP field crews. Further details on the metrics used and evaluation criteria by province can be found in appendix 4 and details on data collection methods in appendix 5 and the AREMP field protocol (AREMP 2009)

Table 3—Inchannel attributes used in assessment and sources of metrics used

Figure 7—Bankfull width-to-depth ratios were calculated from cross-sectional profiles.

Figure 8—Large pieces of wood in the stream channel were counted in each sampled stream reach.

Figure 9—Fine sediment was measured at pool tail crests.

Sidebar 2—Macroinvertebrate Metric

Sidebar 3—Amphibian Metric

Data Analysis

Inchannel data are presented using descriptive statistics and graphical displays of the decision-support model scores. Inferential statistics were used in two cases to test the reliability of generalizing from our sample to the overall population of stream reaches in the NWFP area. First, inchannel scores were tested to determine whether an equal proportion of watersheds fell in each of the five status categories by using a Pearson chi-squared test for goodness of fit (Maindonald and Braun, 2003). Second, scores for the different land use categories were compared to determine if they differed significantly. Selecting the appropriate test was based on whether the classified scores met the assumption of homogeneity of variance between classes. Other test assumptions were accounted for due to the large randomly selected sample (193 watersheds) with no classification of scores resulting in any one class containing a small percentage of the total scores. The Levene test was selected to determine homogeneity of variance since it is relatively insensitive to departures from normality within the classes (Sheskin, 2004).

If the variances of the classified scores were determined to be homogenous, parametric tests were used to compare the mean scores. In these cases, the Student's t-test was employed to compare key and nonkey watersheds and a one-way analysis of variance F-test was used to compare land use allocations. When a significant difference was found within the land use allocations, it was investigated further using a Tukey honest significant difference multi-comparison test to identify which allocations had a significant probability of being different. In cases where the Levene test showed there was a significant chance that the variances of the classified scores were not homogeneous, nonparametric tests were used to test the

classifications. Under these conditions, Wilcoxon rank sum tests were used to test for significant differences between key and nonkey watersheds, while a Kruskal-Wallis test was used to compare differences between land use allocations. When a significant probability of a difference between land use allocations was identified, Behrens-Fisher generalized p-values were used to determine which allocations were different. In interpreting test results, we chose a significance level of 10 percent ($\alpha = 0.10$) as more appropriate than the commonly used 5 percent because of the high natural variability in stream habitats (Al-Chokhachy 2010, Bryant et al. 2004). All analyses were conducted using the R statistical package (R Development Core Team 2011).

2. What is the status of upslope and riparian conditions?

Study Design

The upslope and riparian status assessment used the (sub)watershed as the basic unit of analysis (as described under Study Area). The NWFP area contains 1,379 watersheds that met the sampling criteria of at least 25 percent of stream channels along the 1:100,000 stream layer in federal ownership.

Riparian reserves were defined in the NWFP to have variable widths based on a combination of 100-year flood plains, breaks in slope, riparian vegetation, and site potential trees (USDA and USDI 1994), but these boundaries have yet to be delineated. Therefore, riparian areas for road, vegetation, and landslide risk assessments were based on fixed-width buffers that were placed on the stream layer (see following sections for details). Upslope attributes (also referred to as “watershed-wide” attributes) were calculated for the entire federal portion of the watershed, including the riparian area. While this approach may count riparian areas twice, the upslope and riparian attributes are assessed as proxies for different processes and multicollinearity is not an issue because we are not statistically estimating the influence of explanatory factors. Watershed-wide metrics also avoid wide variation in the amount of nonriparian areas

in watersheds, and they tend to be consistent with available studies on watershed impacts (e.g. road density is typically measured as total watershed density).

Attributes

Each province used a somewhat different set of attributes and model structure to evaluate upslope and riparian conditions (see app. 4 for details); however, all the attributes fell into three basic categories: roads, vegetation, and landslide risk. Attributes were calculated for the federal portion of each watershed based on data from geographic information systems (GIS) and remote sensing data sets. The GIS layers used in the analyses were collected from various sources, including the FS, BLM, and NPS, and other state and federal agencies. Further details on the layers used in the analyses are provided in appendix 5.

Roads—The road layer comprises 2008 data from a variety of sources (see app. 5 for details). Road density in upslope and riparian areas and the frequency of road-stream crossings were determined for all the federal watersheds, except for the Oregon Coast province, which dropped road-stream crossings because of concerns about the effects of the variation in stream mapping density among watersheds (see “Road Data Challenges” sidebar on page xx in Chapter 5, Emerging Issues). All data were based exclusively on the GIS layers, with no field verification. For these analyses, road and 1:24,000 stream layers were clipped to watershed boundaries. A fixed buffer was placed over streams in the watershed to determine riparian area. The width of the buffer was determined during the decision-support model workshops and varied across the provinces. The buffer widths were based on what the participants believed was the relevant area for the riparian process of interest. For example, a narrow buffer was used in evaluating the extent that stream channels were constricted by the presence of a road. Wider buffers were used for evaluating wood and sediment input into streams. For riparian road density analyses, the road layer was laid over the riparian buffer, and miles of road inside the buffer were summarized by watershed. The number of road-stream crossings was calculated by performing a GIS intersection between the road and stream layers.

Vegetation—The analysis used GIS vegetation layers developed by the Interagency Mapping and Assessment Project (IMAP) in Oregon, Washington, and California to assess vegetation characteristics (Moeur et al., in press). Layers were built by using Landsat Thematic Mapper remote sensing data and forest inventory plot data using a gradient nearest neighbor approach (Ohmann and Gregory 2002). Using this method, the closest matching plot data is assigned to each map pixel, enabling a wide variety of vegetation attributes to be calculated. Expert groups from the different aquatic provinces chose somewhat different vegetation attributes, but most chose some combination of canopy cover and mean tree diameter metrics assessed for the whole watershed and for riparian corridors (see app. 4). In each of the federal watersheds, the vegetation layer and the 1:24,000 stream layer were clipped to the federal ownership boundary. A fixed-width buffer, which varied by province, was applied to the stream layer to designate the riparian area (app. 4).

Landslide Risk—As part of the natural disturbance regime, landslides play an important long-term role in the production and renewal of habitat conditions for salmonids (Figure 10; Reeves et al. 1995). However, timber harvesting and road construction can increase the amount of landslide activity beyond natural levels, overwhelming river systems and negatively impact aquatic organism populations (Montgomery et al. 2000, Swanson and Dyrness 1975, Reeves et al. 1995). The AREMP landslide risk attribute was based on an empirical model developed for the Oregon Coast Range (Miller and Burnett 2008) and expanded to the NWFP area using landslide data derived from aerial photographs in 14 AREMP sixth-field watersheds. The model used widely available digital elevation model (DEM) and land-cover data (10-m DEMs and 25-m satellite imagery) and found landslide frequencies varied by topography, forest cover, and proximity to roads: landslide frequency was 48% higher in areas with early seral vegetation (average diameter < 4 inches) and 170 percent higher in areas within 50 m of roads (Miller 2006). AREMP landslide risk attribute evaluates the effect of management activities on landslide susceptibility by comparing the “current” landslide susceptibility with “baseline” susceptibility by sixth-

field watershed. Baseline susceptibility was determined using the 10-m DEMs and forest cover class, assuming that trees greater than 4 inches in diameter were growing in all areas capable of supporting tree growth. We determined susceptibility in each cell in the DEM grid and then calculated the average susceptibility rating across all cells in the sixth-field watershed. This measurement was considered the best approximation of a watershed's natural or baseline susceptibility for landsliding, although we recognize that we are underestimating susceptibility because we do not account for loss of trees owing to natural processes. The "current" susceptibility rating was determined by multiplying the baseline rate of each cell in early seral areas by a factor of 1.48 and areas within 50 m of a road by a factor of 2.7. The effect of management on landslide susceptibility was calculated as a ratio of the current susceptibility to the baseline. Landslide risk was included for all aquatic provinces except the Klamath/Siskiyou /Franciscan province because model results did not match up well with a locally derived model, so we are continuing to work with local provincial experts to refine this attribute before including it in the decision-support model.

Figure 10—Naturally occurring landslides provide large wood and spawning gravels to rivers. However, excessive landslides caused by management activities can overwhelm river systems and negatively impact aquatic organisms.

Data Analysis

The condition of upslope and riparian processes was estimated for the federal portion of each sixth-field watershed in the NWFP area having greater than 25 percent federal ownership along the 1:100,000 stream layer. Condition scores are calculated by evaluation models designed for each aquatic province, which aggregate a number of attributes (principally road and vegetation) derived from GIS and remote sensing data (see "Methods" section and app. 4). The normalized watershed condition scores range from -1 to +1 and are positively related with the condition of watersheds: watersheds in good condition have higher scores than those in poor condition. Because data on every watershed in the target population were analyzed, inferential statistics are not needed to test the reliability of generalizing results from a sample to

THE FINDINGS AND CONCLUSIONS IN THIS REPORT ARE IN PRESS AND SUBJECT TO CHANGE PRIOR TO FORMAL DISSEMINATION BY THE AGENCIES AND SHOULD NOT BE CONSTRUED TO REPRESENT AGENCY DETERMINATION OR POLICY.

a larger population. All differences are effectively statistically significant, so what remains for judgment is whether differences are meaningful in terms of biology or management.

Sidebar 4—Distribution Functions

3. *What is the trend in watershed conditions?*

Study Design

Since no repeat sampling has yet occurred for the inchannel data, it was only possible to use the upslope and riparian data set to calculate trend. The same 1,379 watersheds as used for question 2 are used here. Assessment of trend estimates watershed condition status in 1994 and again in 2008 based on upslope and riparian process attributes.

Attributes

Attributes used for the trend analysis are the same as those described above under the status of upslope and riparian condition. Additional information related to the calculation of these metrics related to the two time periods is provided below (with further details in appendix 5).

Roads—Information on road building and decommissioning conducted since the NWFP was implemented is spotty and incomplete. Although most of the federal road layers contain attributes that describe whether specific road segments were decommissioned, dates of decommissioning and road building on Forest Service land were not available. Historical roads layers (e.g., from 1994 when the NWFP was implemented) also were not available. Therefore, to obtain road density data from 1994 (time 1) and 2008 (time 2) to analyze change, we used total road miles (existing + decommissioned) as the time 1 data point and the existing roads as the time 2 data point. We assumed that all the roads were decommissioned later than 1994. The Forest Service and the BLM rarely decommissioned roads before

1990.⁵ Although analyzing just the miles of roads that have been decommissioned may seem more straightforward than looking at miles of roads in time 1 and time 2, this approach would not allow us to use the decision-support model and determine the distribution of watershed conditions across the NWFP area in the two periods. For simplicity, we refer to the range of dates as 1994 to 2008 in subsequent figures to match our 15-year reporting period.

Vegetation—For change analysis, vegetation layers for the Plan area were developed (as described above under question 2) for two dates: 1996 and 2006 in Washington and Oregon, and 1994 and 2007 in California. The temporal consistency of the vegetation models was enhanced by using Landsat imagery that had been geometrically rectified and radiometrically normalized through time (i.e., 'temporally normalized') using the LandTrendr algorithms (Kennedy et al. 2010) and by using a single 'pool' of inventory plots for model development for both dates (Moeur et al, in press). For simplicity, we refer to the range of dates as 1994 to 2008 in subsequent figures to match our 15-year reporting period.

Landslide Risk—Since landslide risk is based on the intersection of roads and early seral vegetation with topographic risk, it was simply calculated for time 1 (1994) and time 2 (2008) based on the roads and vegetation layers developed for each period as described above. The underlying topographic risk factor was not changed between the two periods.

Data Analysis

The results were based on the amount of change that occurred; i.e., trends in watershed condition were calculated by comparing the 2008 to the 1994 upslope/riparian scores. Trend scores are simply the 2008 score minus the 1994 score, which results in a possible range between -2 and +2 (given that the maximum theoretical change is from a -1 score in 1994 to a +1 score in 2008, or vice-versa). Positive trend scores indicate an improvement in condition and negative scores a decline. No statistical tests were

⁵ Erkert, T. 2003. Personal communication. Road engineer, USDA Forest Service, Pacific Northwest Region, 333 SW First Ave., Portland, OR 97204.

needed to compare scores because trend was assessed based on a complete data set of the population of interest rather than a sample of a larger population.

Nevertheless, there is measurement error in the underlying data attributes. Of particular importance to the trend calculation is the fact that slight pixel-level differences in the Landsat imagery between dates can result in a change in the vegetation plot (and associated attributes) assigned to a pixel. While both plot assignments generally approximate on-the-ground conditions, the switch will cause slight changes in attributes resulting in “noisiness” in the model scores beyond what has occurred on the ground; however, the magnitude of this noise is unknown at the current time (see Moeur et al. 2010 for further discussion). As simple way to reduce the effect of these fluctuations and provide a conservative bound for estimated changes, we also calculated the percentage of watersheds trending up or down using an arbitrary minimum change threshold of greater than 5 percent (± 0.1 in model score).

To further understand what was driving the trends, we developed a method to assess the relative contributions from the roads, vegetation, and landslide risk components. Watershed were divided into seven bins by their overall trend scores (-2 to -0.3, -0.3 to -0.1, -0.1 to <0, 0, >0 to 0.1, 0.1 to 0.3, 0.3 to 2). For each watershed, the relative contributions were calculated using the following formula:

AttributeScore_i = trend subscores (-2 to +2) for roads, vegetation, and landslide risk

BinBoundary = boundary of the bin range closest to zero

The percentage contribution of attribute subscores for each bin was calculated by taking the average of the individual watershed contributions.

Model Validation

The reliability of our watershed condition assessments depends on the validity of the evaluation models used. Model validation, as understood in the natural sciences, means testing to see if a model produces empirically accurate results with respect to independent, real world observations (Oreskes 1994). However, expert-based systems, such as used here, are often built for situations in which such empirical tests are neither possible nor affordable – watershed condition is more of a concept than a measurable attribute in the environment. For this reason, validation in the expert systems sense is often done by comparing model processes and results back to the judgments of experts (Turban and Aronson 2001). We accomplished this type of validation by checking preliminary results with the provincial expert groups, identifying and researching discrepancies between results and experts opinions, and adjusting the models to better reflect their judgments.

We also employed two data-based analyses as a check on model results. First, we compared watershed model results to a few simple indicators measured consistently across the NWFP area, and, second, we compared inchannel to upslope and riparian results.

Uniform Indicators

The watershed evaluation models used for questions 2 and 3 differ somewhat by aquatic province. Attributes used in the watershed condition models for the different aquatic provinces were similar in nature (roads, vegetation, landslide risk), but the specific metrics (e.g. tree size, width of riparian buffers) and evaluation criteria varied by province. The rules for integrating the attributes into overall condition scores also varied. As an alternative view and general check on these results, this analysis looked at a few key indicators individually and in a consistent fashion across all provinces.

Study Design

The same set of 1,379 watersheds as used for questions 2 and 3 were evaluated for the same time periods, 1994 and 2008.

THE FINDINGS AND CONCLUSIONS IN THIS REPORT ARE IN PRESS AND SUBJECT TO CHANGE PRIOR TO FORMAL DISSEMINATION BY THE AGENCIES AND SHOULD NOT BE CONSTRUED TO REPRESENT AGENCY DETERMINATION OR POLICY.

Attributes

Road density—Road density has often been used in watershed assessments because of documented effects on a number of stream attributes, broadly including patterns of water and sediment delivery (Reid and Dunne 1984; Gucinski et al. 2001; Lee et al. 1997; Reiman et al. 2001). Road density was chosen as an indicator for most of the AREMP provincial models. The metric used was watershed road density, expressed as the number of road miles per square mile of watershed area, and it was calculated the same as described above under upslope and riparian status attributes. Figure 11 illustrates road density for a single watershed.

Late-seral vegetation— A number of the provincial models incorporated indicators of the amount of the watershed and/or riparian area in mid- to late-seral stages, as approximated by average tree diameter and canopy cover metrics (Figure 12). As an alternative indicator, we adopted the late-successional old-growth (LSOG) metric used by the NWFP Vegetation Monitoring Program, which was an average conifer diameter ≥ 20 inches and conifer canopy cover ≥ 10 percent (Moeur et al., In press). The metric was derived from the same data sets used for vegetation attributes for questions 2 and 3 (see app. 5).

Vegetation disturbance—The loss of natural vegetation cover can also be a major influence on water and sediment delivery (Reid 1993; Grant et al. 2008). A number of AREMP provincial models included indicators of early seral conditions as a proxy for disturbance (Figure 13). For this alternative metric, we made use of a new approach to extracting vegetation disturbance information from satellite data, called LandTrendr, developed in conjunction with the NWFP Late-successional Old-growth monitoring module (Kennedy, Yang and Cohen 2010; Cohen, Yang and Kennedy 2010). LandTrendr classifies disturbance into long- and short-duration events. Long-duration events were attributed to insect and disease agents. Short-duration events were attributed to fire and harvest. Fire-caused losses were identified by matching LandTrendr disturbance areas with digitized fire polygons from the national

interagency fire database, Monitoring Trends in Burn Severity (MTBS). Harvest was defined as any short-duration disturbance not characterized as caused by fires recorded in the MTBS fire layer. Since the MTBS layer was the only way to distinguish fire from other disturbances, this harvest metric also includes blowdown and small fires not captured in MTBS.

For the alternative AREMP metric percentage of equivalent disturbed area was derived from the LandTrendr disturbance layer. All LandTrendr disturbances have a magnitude value of 15 (minimum detection threshold) to 100 percent, so the equivalent acres were calculated by multiplying the magnitude by the disturbance area. Percent disturbed areas for each year were summed into five periods, 1985–1989, 1990–1994, 1995–1999, 2000–2004, and 2005–2008. Note that it's possible for an area to be more than 100 percent disturbed if it has multiple disturbances within a 5-year period. For example, if a watershed burned completely (100 percent) twice in a 5-year period, it would receive an aggregate fire disturbance value of 200 percent. The vegetation disturbance from insects and disease was less than 0.3 percent of the NWFP area over the 1994 to 2008 period and so was ignored for this analysis.

Data Analysis

These alternative indicators were analyzed individually using descriptive statistics and graphical displays of the original indicator units (e.g. mi/mi²) as opposed to scaled model scores. Again, no statistical tests are needed, given that the data are a complete census of the population of interest.

Figure 11—Road density (miles per square mile of watershed) was used as an indicator of watershed condition. In this example of a sixth-field watershed (purple line), the black lines are roads, blue lines are streams, and red dots are road-stream crossings.

Figure 12—Watershed condition evaluations included metrics of average tree diameter and percent canopy cover in riparian and upslope areas.

Figure 13—Disturbance from fire and harvest was used an alternative metric for assessing watershed condition.

Upslope Versus Inchannel Conditions

In the 2004 monitoring report (Gallo et al. 2005), we combined the upslope/riparian and inchannel scores to derive an overall watershed score. We chose not to make this combination for this iteration for a number of reasons. First, only 193 watersheds have been sampled for inchannel attributes, whereas we were able to generate upslope and riparian data for all 1,379 watersheds in the NWFP areas. Second, we lack trend data for the inchannel attributes. Third, the separation emphasizes the fact that they provide different information about watershed condition. Inchannel scores are our best estimate at current conditions for fish and other aquatic biota, whereas watershed-wide conditions represent more of a risk assessment: What is the likely impact of upslope and riparian conditions on the future state of aquatic organisms? Instead of combining the scores, we report the them separately. We also decided to look for correlation between the scores, in order to understand the extent to which they are measuring directly related aspects of watershed condition.

To assess correlation between the two data sources, we paired the inchannel model scores from the 193 watersheds (2002-2009) to the reduced set of upslope/riparian model status scores (2006-2008) for the same watersheds. We performed a simple linear regression on these paired data and tested against the null hypothesis that the slope of the linear relationship was zero based on a standard t-test.

Chapter 3: Results

Results are presented for each of the key monitoring questions, whose answers provide insight for evaluating the success of the aquatic conservation strategy in the entire Northwest Forest Plan (NWFP) area and by land use allocation. As described in the “Methods” section, standardized condition model scores range from -1 to +1, with watersheds in good condition having higher scores than those in poor condition. For much of the following display and analysis, we have grouped scores into the five equal condition categories very high (>0.6), high (>0.2 and ≤ 0.6), moderate (≤ 0.2 and ≥ -0.2), low (< -0.2 and ≥ -0.6), and very low (< -0.6).

1. What is the status of inchannel conditions?

The distribution of scores for the 193 subwatersheds with inchannel data ranged from -0.76 to +1.0, with a mean score of +0.2. Grouping scores into categories, shows that relatively few fell into the low (10 percent) and very low (1 percent) categories (Figure 14). The majority of inchannel attribute scores fell into the moderate (35 percent) and high (41 percent) ranges, with relatively few (12 percent) in the very high category. The number of watersheds in each category was compared to a theoretical uniform discrete distribution in which each category had an equal number. The actual distribution of scores is different from the equal distribution (Pearson chi-squared, p -value < 0.0001) indicating that the number of scores in each category is not similar to the result of random effects.

Figure 14—Inchannel condition scores by status category for the 193 randomly selected watersheds in the Northwest Forest Plan area that have been sampled for inchannel attributes as of 2009.

For the low and very low-scoring subwatersheds (scores < -0.2), water temperature was usually the most influential factor (mean temperature score = -0.77). In many of the provincial evaluation models, a poor water temperature score carried more weight than other attributes because it was only measured once for each subwatershed (at the lowest elevation on federal land), in contrast to the other attributes, which

THE FINDINGS AND CONCLUSIONS IN THIS REPORT ARE IN PRESS AND SUBJECT TO CHANGE PRIOR TO FORMAL DISSEMINATION BY THE AGENCIES AND SHOULD NOT BE CONSTRUED TO REPRESENT AGENCY DETERMINATION OR POLICY.

were averaged over 6 to 8 sites. Aquatic invertebrate scores (mean = -0.6) also appeared influential in producing the low scores.

Looking at the distributions of individual attribute scores, one can see that pools, substrate, and invertebrates were all negatively skewed, with more positive scores and a longer negative tail, whereas wood was more evenly distributed with a median near zero (Figure 15). Temperature actually fell into a bi-modal distribution with many scores at the extremes (-1 and +1, causing the lack of "whiskers" on the box-whisker chart), owing to temperatures either above or below the evaluation criteria range.

Figure 15—Distribution of attribute scores for aquatic pools, substrate, wood, temperature, and aquatic invertebrates and for the 193 randomly selected watersheds in the Northwest Forest Plan area that were sampled for inchannel attributes as of 2009.

Figure 16 shows the spatial distribution of inchannel scores. Low scores are found only in the southern half of the NWFP area, with over 70 percent of watersheds in the low and very low categories occurring in the Klamath/Siskiyou and Franciscan provinces.

Figure 16—Distribution of inchannel condition scores for the 193 randomly selected watersheds in the Northwest Forest Plan (NWFP) area that have been sampled for inchannel attributes as of 2009.

Inchannel Status by Land Category

Examining overall inchannel score results in the context of land use allocations (Figure 17) shows that congressionally reserved lands have the highest scores (mean +0.31, median +0.31), followed by late-successional reserve (LSR) (mean +0.21, median +0.22) and matrix (mean +0.14, median +0.16). Statistical analysis showed that the difference between median scores for reserved lands and matrix lands is significant (Behrens-Fisher generalized p-value = 0.036), whereas the difference between LSR lands and the other two land use allocations is not significant (LSR-matrix generalized p-value = 0.649, LSR-reserved generalized p-value = 0.230). The reserved class had the least variability, followed by increasing

distribution ranges for LSR and matrix lands. There was a significant difference between the means of the key and nonkey categories (Student's t-test, p-value = 0.042), with key watersheds scoring higher (mean +0.27, median +0.29) than nonkey watersheds (+0.16, median +0.15). Their distribution ranges were also different, with the range of key watershed scores more concentrated than for nonkey.

Figure 17—Distribution of inchannel scores by land use category for the 193 randomly selected watersheds in the Northwest Forest Plan area that were sampled for inchannel attributes as of 2009 (reserved = congressional reserves, LSR = late-successional reserves).

The distributions of individual attribute scores were more variable in regard to land use categories (Figure 18). Statistically significant differences between attribute scores by land use category are summarized in Table 4. Pool scores were high across all land use allocations, and the median score on matrix lands was found to be significantly greater than on congressionally reserved lands (Behrens-Fisher generalized p-value = 0.02); no difference was found between key and nonkey watersheds (Wilcoxon rank sum p-value = 0.13). In contrast, inchannel wood scores were found to be higher in key than nonkey watersheds but no differences were confirmed for the reserved/LSR/matrix categories. Examining the distribution of aquatic invertebrate scores by land use allocation showed that congressionally reserved lands had a higher mean value than for matrix and LSR lands. No statistically significant differences were found for substrate or temperature scores. Although the mean temperature score on reserved lands appeared considerably higher than on LSR or matrix lands, the difference was not statistically significant due to the high dispersion in scores across all allocations (min -1.0, max 1.0 for all three allocations).

Figure 18—Distribution of inchannel attribute scores by land use category (reserved = congressional reserves, LSR = late-successional reserves).

Table 4--Inchannel scores by land use category (CR = congressional reserves, LSR = late-successional reserves).

2. What is the status of upslope and riparian conditions?

The condition of upslope and riparian processes are estimated by scoring and integrating a variety of road and vegetation attributes derived from remote sensing and other mapped data sets (see “Methods” section and app. 4 for details). Data on every watershed in the target population were analyzed; therefore, inferential statistics are not needed to test the reliability of results. Measurement error inherent in the attributes is still an issue; however, it has not yet been fully quantified and propagated through the assessment models (see “Discussion” section and app. 6).

Overall watershed condition scores of the 1,379 watersheds ranged from a low of -0.99 to a high of +1, with a mean score of +0.17. Scores are clustered in the center of the distribution and skewed slightly positive, with 50 percent falling between approximately -0.25 and +0.5. Figure 19 presents a view of the distribution of scores by status category. The largest percentage fell into the moderate category (29 percent), followed by the high category (25 percent). Comparable amounts were classified as very high (22 percent) and low (21 percent), but relatively few watersheds scored in the very low (4 percent) category.

Figure 19—Watershed condition scores in the Northwest Forest Plan area by status category (2008).

The spatial distribution of watershed scores shows some noticeable patterns (Figure 20). High scores are found in the central Olympic Peninsula (Olympic National Park), the north central Cascades, the Oregon Coast Range, and scattered pockets along the Klamath-Siskiyou mountain range (mostly corresponding to designated wilderness areas). Low condition scores can be seen in the southern Olympic region, eastern Klamath-Siskiyou, and along the eastern and western flanks of the Cascade Range in Oregon and Washington. Breaking these scores down by aquatic province reveals some notable patterns (Table 5). First, no watersheds were rated low in the Oregon Coast or High Cascades, and very few in the Olympic and Klamath/Siskiyou/Franciscan. The Western Cascades had both the largest

percentage in the very low category and the lowest distribution overall. In contrast, the Oregon Coast, North Cascades, and Olympic all had the highest percentage of their watersheds in the very high category.

Figure 20—Watershed condition status scores (2008) in the Northwest Forest Plan (NWFP) area.

Table 5—Watershed condition status scores (2008) by aquatic province

Watershed Status by Land Category

Figure 21 presents the summary of watershed scores (using upslope and riparian data) by land category. Reserved lands having the highest scores (mean +0.56, median +0.62), followed by LSR (mean +0.14, median +0.15) and matrix (-0.12, median -0.13). No watershed that was predominately reserved land received the minimum score (-1). Conversely no watersheds in the matrix category received a perfect score (+1). Although the distributions of the ACS categories had similar ranges, key watersheds generally scored higher (mean +0.27, median +0.27) than nonkey watersheds (mean +0.11, median +0.09).

Figure 21—Distribution of watershed condition status scores (2008) by land use allocation (reserved = congressional reserves, LSR = late-successional reserves).

Figure 22 shows the frequency of score occurrences by land category, and it reveals a distinct distribution for each category. A preponderance of matrix scores are in the middle range (-0.4 to +0.4), with very few greater than +0.5. The greatest frequency of scores in the LSR category also occurs in the middle range (+0.15) but there is a greater frequency of higher scores across the range and especially at the upper end. The reserved class shows very few watersheds at the lower end and a steadily increasing number of higher scores, until a small dropoff just before +1. Differences between key and nonkey watersheds are less pronounced (Figure 23). Both distributions have their maximum frequencies around

+0.25 but nonkey watersheds have a greater frequency of scores below this value, whereas key watersheds have greater frequencies above it.

Figure 22—Frequency of watershed status scores (2008) by land use allocation (reserved = congressional reserves, LSR = late-successional reserves). These curves show the data in a continuous manner, rather than by data bins (i.e., histograms).

Figure 23—Frequency of watershed status scores (2008) for key and nonkey watersheds. These curves show the data in a continuous manner, rather than by data bins (i.e., histograms).

Looking at the spatial distribution of watershed scores in the various categories (Figure 24), an obvious pattern is that many of the higher scoring watersheds on congressionally reserved lands lie near the Cascade Crest, where higher elevation lands are naturally less amenable to human activities. The contiguous nature of the Olympic National Park, which makes up a significant portion of reserved lands, also may help to explain the high watershed quality on these lands. The spatial distribution of scores by key and nonkey watersheds does not reveal any further patterns by this designation (Figure 25).

Figure 24—Watershed condition status (2008) by land use allocation in the Northwest Forest Plan (NWFP) area.

Figure 25—Watershed condition status (2008) for key and nonkey watersheds in the Northwest Forest Plan (NWFP) area.

3. What is the trend in watershed conditions?

Overall, there was a positive change in watershed scores, from a mean score of +0.14 in 1994 to +0.17 in 2008. Scores increased for 69 percent of watersheds versus 23 percent showing declines. Since a portion of these shifts was likely due to errors inherent in the satellite imagery classification process, we also calculated a more conservative estimate looking at only condition score changes of greater than 5 percent (± 0.1); using this threshold, 10 percent of watersheds increased versus the 4 percent which decreased (Table 6).

THE FINDINGS AND CONCLUSIONS IN THIS REPORT ARE IN PRESS AND SUBJECT TO CHANGE PRIOR TO FORMAL DISSEMINATION BY THE AGENCIES AND SHOULD NOT BE CONSTRUED TO REPRESENT AGENCY DETERMINATION OR POLICY.

Table 6—Percent of watersheds (n = 1,379) that decreased, increased, or had no change in watershed scores between 1994 and 2008.

Figure 26 shows a general increase in scores (a shift to the right), especially for watersheds in the low (centered around -0.75) and mid to high (+0.25 to 0.75) ranges. The number of watersheds in the very high range (+0.9) actually decreased slightly, primarily as a result of fire on reserved lands.

Figure 26—Frequency of watershed condition scores in 1994 and 2008. These curves show the data in a continuous manner, rather than by data bins (i.e., histograms).

Figure 27 and Table 7 examine the watershed condition score changes in more detail. Figure 27 reveals that most score changes (81%) were relatively small (± 0.1 in model score or 5% of possible change from -1 to +1). Fifty-five percent of the watersheds had score increases between 0 and +0.1; increases in vegetation scores were the dominant driver (66%) for this range, but with some contribution from roads (24%) and landslide risk (11%; Table 7). Scores declined by 0.1 or less in 18% of watersheds, and in this portion of the distribution the relative influence of vegetation scores (83%) to road scores (2%) was much greater. The landslide risk contribution remained comparable (15%), but it was likely driven more by vegetation changes in landslide prone areas than road changes. Eight percent of watersheds showed no change in score (dot on Figure 27). There were two reasons why a score might not change, either 1) there were no changes in any of the underlying attributes between the sampling dates, or 2) an increase in one or more attributes was cancelled out by declines in others (generally these increases/decreases were quite small).

Figure 27—Distribution of changes in watershed condition scores between 1994 and 2008.

Table 7—Attribute influences on watershed condition score changes between 1994 and 2008

The proportion of changes in watershed scores due to roads, vegetation, and landslide risk diverges further between the positive and negative tails of the distribution. Vegetation losses (and associated landslide risk increases) were responsible for all of the negative trend scores beyond -0.3. Five watersheds decreased by more than -0.5; all were located in the Klamath/Siskiyou and Franciscan provinces and experienced heavy vegetation losses from the 2002 Biscuit Fire. Positive trends in the +0.1 to +0.3 range were mostly due to road decommissioning (42%) but with a fair contribution from vegetation (30%) and landslide risk (28%). Reduced landslide risk was the dominant driver of improvement about the +0.3 level (54%), apparently due to road decommissioning in landslide prone areas, since road improvements contributed considerably more (38%) than vegetation (8%) at this level. Only four watersheds increased by 0.5 or more, and three of these four were driven primarily by road decommissioning. Fish Creek in the Western Cascades had over 100 miles of road decommissioned and realized the largest score increase overall (+1.5). In contrast, the increase in the Olympic Province's Deep Creek (+0.5) was entirely due to vegetation and its associated effects on landslide risk.

For simplification and display, we have again grouped scores into categories. Figure 28 shows the change in the distribution of watersheds by category. Most notable are the decline in watersheds in the very low category and an increase in the high category.

Figure 28—Watershed condition scores by status category in 2008.

Looking at a display of these changes by status categories in 1994 and 2008 (Figure 29), one sees that most changes occurred in the middle score categories (low, moderate, high) and that these changes were predominantly positive. Some watersheds from all three of these classes improved to the very high category, demonstrating that large changes appear possible in a relatively short period (15 years). The categories at the extremes (very low and very high) showed the least amount of change. This lower mobility was likely due to many of their indicator values being either considerably above or below the

evaluation thresholds, so even moderate changes in the indicator values would not change the indicator scores (e.g. a change in road density from six to four mi/mi² would not change the evaluated score from -1).

Figure 29—Number of watersheds with category scores from 1994 and 2008. The bubble sizes range from few watersheds (smallest bubbles) to many watersheds (largest watersheds).

The watershed condition trend map (Figure 30) uses seven categories instead of five, and the central categories also have a smaller interval (0.2) than the extremes (0.5) for better discrimination because changes in scores tended to be more tightly grouped than the status scores. The largest negative changes can be seen in southwest Oregon and are due to loss of vegetation from the Biscuit Fire. Other declines can be seen scattered mostly along the east side of the Cascade crest, again principally because of large fires.

Figure 30—Change in watershed condition score 1994 to 2008.

Watershed Trend by Land Use Category

All land use categories had more watersheds with score increases than decreases between 1994 and 2008 (Table 6). The proportion of increases to decreases was much higher (almost double) in late-successional reserves and matrix lands than in congressional reserves. The mean score of the reserved class actually declined slightly (-0.01), while the LSR (+0.05) and matrix (+0.04) both showed slight increases on average. Key and nonkey watersheds both had approximately three times more watersheds with score increases as compared to decreases. There was also little difference in the average trend between the key (mean +0.04) versus nonkey (+0.03) watersheds

The magnitude of changes did differ somewhat by land use allocation (Figure 31). While the majority of changes were small ($< \pm 0.1$) for all categories, the LSR experienced more of the larger positive changes ($> +0.1$) than the other two classes. Figure 32 shows where in the overall score distributions the

THE FINDINGS AND CONCLUSIONS IN THIS REPORT ARE IN PRESS AND SUBJECT TO CHANGE PRIOR TO FORMAL DISSEMINATION BY THE AGENCIES AND SHOULD NOT BE CONSTRUED TO REPRESENT AGENCY DETERMINATION OR POLICY.

largest changes occurred. Matrix watersheds had the most positive score changes in the lower (-0.9 to -0.6) and middle-upper (+0.2 to +0.6) ranges. Changes in the LSR category were similarly distributed at the lower end and more broadly at the middle-upper range (-0.2 to +0.7). The reserved category showed a shift from the upper end (+0.8 to +1.0) to slightly lower scores (+0.5 to +0.8).

Figure 31—Distribution of changes in watershed scores from 1994 to 2008 by land use allocation.

Figure 32—Frequency of 1994 and 2008 watershed scores by land use allocation. These curves show the data in a continuous manner, rather than by data bins (i.e., histograms) (reserved = congressional reserves, LSR = late-successional reserves).

Differences between key and nonkey watersheds were less pronounced. There was little difference in the average trend between the key (mean +0.04) versus nonkey (+0.03) watersheds. The change magnitude distributions were quite similar in form, and again the majority of changes were small ($< \pm 0.1$) for both categories (Figure 33). Looking at the continuous score distributions (Figure 32), both categories had positive score changes in the lower (-0.9 to -0.6) and middle-upper (+0.1 to +0.6) ranges (Figure 34). Key watersheds experienced more positive changes in the middle range (-0.2 to +0.2).

On average, more roads were decommissioned in key watersheds (4.1 mi) than in nonkey (1.6 mi); however, in terms of watershed road density (miles road per square mile of watershed) the difference in changes are smaller: an average of -0.14 mi/mi^2 in key and -0.07 mi/mi^2 nonkey watersheds. Given that the evaluation criteria thresholds for road density ranged from 0.5 mi/mi^2 (+1 score) to 4 mi/mi^2 (-1 score), a decline in density on the order of -0.14 mi/mi^2 would not have a large effect on the overall score. Two examples show the possible range of effects. Lower Fish Creek in the western Cascades had the largest mileage of roads decommissioned (118 mi) and the highest decline in road density (-2.5 mi/mi^2), which dropped road density from 3.3 mi/mi^2 in 1994 to 0.8 mi/mi^2 in 2008 with a high corresponding score change from -0.9 to +0.8. In contrast, the lower South Fork Skokomish River in the

Olympic province had the second highest decline in road density (-1.6 mi/mi²), which dropped road density from 4.6 mi/mi² in 1994 to 3.0 mi/mi² in 2008 but produced no score change because it was still at or above the evaluation criteria of 3 mi/mi² that defines poor condition.

Figure 33—Distribution of changes in watershed condition scores between 1994 and 2008 for key and nonkey watersheds.

Figure 34—Frequency of 1994 and 2008 watershed condition scores for key and nonkey watersheds. These curves show the data in a continuous manner, rather than by data bins (i.e., histograms).

Model validation

The watershed evaluation models used to address questions 1 through 3 are complex in that they integrate a number of different indicators and do so somewhat differently for each province. For an alternative view and check on our results, we looked at three single indicators measured consistently across the NWFP area: road density, the percentage of area in large trees, and the amount of disturbance from harvest and fires.

Road Density

The density of roads in watersheds ranges from 0 to 7.1 miles of road per square mile in 1994 and from 0 to 6.8 mi/mi² in 2008. Our data set registered 3,406 miles of roads decommissioned between 1994 and 2008. However, the effect of this removal on the overall distribution of road densities was small, from a median of 2.42 mi/mi² (mean of 2.35) in 1994 to 2.31 mi/mi² (mean of 2.25) in 2008 (Figure 35). Looking at the distribution of changes reveals that only 5 percent of watersheds showed any increase in road density, 45 percent had no change, and 50 percent decreased (Figure 36). Increases were small, a maximum of 0.2 mi/mi², and decreases ranged to the -2.5 mi/mi² in Fish Creek in the western Cascades.

Figure 35—Watershed road density (mi/mi²) in 1994 and 2008.

Figure 36—Changes in watershed road density (mi/mi²) from 1994 to 2008.

For context, the watershed model evaluation criteria for most of the provinces rate less than 0.5 to 1 mile of road per square mile of watershed as good condition and from 2.4 to 4 mi/mi² as poor condition. Using these rough guides and looking at Figure 35, one can see that approximately 25 to 50 percent of watersheds (the top half and/or whisker in the box charts) would be rated as poor condition using this single indicator. This result is considerably lower than the distribution of our model evaluation scores under question 1, where only 4 percent of the watersheds fell into the poor category. However, these results are roughly consistent with what happens inside the evaluation model, with roads often scoring poorly but with vegetation scoring well, bringing up the overall scores. The result is also consistent with discussions in our evaluation workshops, where experts have identified road density as the major problem in many watersheds.

By Land Use Category

Road density distributions by land use category (Figure 37) are consistent with the overall watershed model results (Figure 21). The lowest road densities were found in congressionally reserved lands (mean 0.76 mi/mi²), higher densities in the LSR category (mean 2.6 mi/mi²), and the highest densities on matrix lands (mean 3.1 mi/mi²). Although road building is generally prohibited on congressionally reserved lands, some contain legacy roads, and our classification by entire watersheds means that nonreserved lands will often be included in watersheds classified as reserved. Key watersheds also tend to have lower densities (Figure 37D, mean 1.9 mi/mi²) than nonkey watersheds (Figure 37E, mean 2.4 mi/mi²).

Figure 37—Road density by land category, 2008 (reserved = congressional reserves, LSR = late-successional reserves).

In terms of trend, the LSR category had a larger average reduction in density (mean -0.15 mi/mi^2) than the matrix (mean -0.10 mi/mi^2) or reserved (mean -0.03 mi/mi^2) categories. Road density reduction in key watersheds (mean -0.15 mi/mi^2) was average approximately double that in nonkey watersheds (mean -0.07 mi/mi^2). Small reductions (between 0 and 0.1 mi/mi^2) were the most frequent, occurring in 20 to 30 percent watersheds across all categories (Figure 38).

Figure 38—Change in road density (mi/mi^2) by land use category, 1994 to 2008.

Late-successional old-growth

A number of the provincial models incorporate indicators of the amount of the watershed and riparian area in different seral stages, as approximated by average tree diameter and canopy cover metrics. For a simple, uniform comparison metric, we used percent of watershed area in late-successional old-growth (LSOG) vegetation, as defined by the NWFP Vegetation Monitoring Program (average tree diameter ≥ 20 inches and canopy cover ≥ 10 percent).

The percentage of watershed area meeting this large tree definition ranges from 0 to 89 percent over the 1,379 watersheds, and this range remained constant from 1994 to 2008. The mean percentage decreased from 30.1 to 29.5 in this period, and the median also dropped from 30 to 29.7 (Figure 39). Large tree coverage between 1994 and 2008 decreased in more watersheds (57 percent) than it increased (42 percent) (Figure 40). Changes were generally small, and therefore somewhat uncertain, given the classification error inherent in the satellite data. Only 1.3 percent of watersheds experienced a more definite increase of greater than 5 percent in LSOG, while 4.3 percent of watersheds showed a loss in LSOG of greater than 5 percent.

Figure 39—Percentage of watershed area in late-successional old-growth vegetation (LSOG; conifer quadratic mean diameter ≥ 20 inches and canopy cover ≥ 10 percent) in 1994 and 2008.

Figure 40—Change in percentage of watershed area in late-successional old-growth vegetation (LSOG; conifer quadratic mean diameter > 20 inches) from 1994 to 2008.

Comparing these LSOG results to vegetation scores from our watershed models is problematic because the watershed vegetation metrics vary considerably between and even within provinces. The variations in the watershed models are designed to adjust for local growth potential, as influenced by factors such as high elevations and low precipitation regimes. For example, the High Cascades model has both riparian and upslope vegetation evaluations. In the riparian zone, the metric is the percent of the watershed where average tree diameter is ≥ 14 inches, except in lodgepole pine dominated areas where a ≥ 8 inch diameter threshold is used. The metric is scored by comparing it to a target range of 30 percent (-1 model score) to 70 percent (+1 score). The upslope evaluation uses the area where average tree diameter is ≥ 8 inches and canopy cover is ≥ 40 percent. Target ranges vary, depending on whether the watershed falls within the rain-on-snow zone or not (see Appendix 4 for model details by province).

Despite these differences, a comparison of trend results between the watershed model vegetation scores and the simpler LSOG metric should help better understand both measures. The overall distributions of the two metrics moved in opposite directions. In contrast to the declines in LSOG area, as described above, AREMP model vegetation scores increased in more watersheds (58%) than decreased (28%). There were also larger changes in the model scores ($> 5\%$ or $< -5\%$, equivalent to ± 0.1 in model score), and more of these changes were positive than negative (19% versus 12%). No overall pattern to these differences between LSOG and watershed model vegetation scores was evident; watersheds with divergent trends were scattered across all provinces. Reasons for the disparities varied by province and by individual watershed attributes. Some of the most divergent scores occurred in the High Cascades in the area of the 2003 B & B Fire, where modeled vegetation scores trended down sharply but LSOG scores actually increased. These watersheds had very little vegetation classified as LSOG (< 3 percent) to begin

with, so small increases in LSOG area translated into relatively large gains when measured as a percentage change. The fixed minimum and maximum area targets used to derive the watershed vegetation scores (e.g. 30 to 70 percent for riparian vegetation) are more robust to such differences in areas. High trend differences between LSOG and watershed vegetation scores also appear in burned areas of the Klamath/Siskiyou and Franciscan province. In those watersheds, however, both LSOG and model metrics declined, but LSOG fell much more sharply. The divergence was due to the fact that this provincial evaluation model used indicators based on canopy cover and early seral (< 5 inch diameter) area, both of which rebound much more quickly after severe fires than the LSOG large tree metric.

By Land Use Category

The distribution of LSOG by land use category (Figure 41) does not follow the same pattern as the watershed condition model results (Figure 21). In particular, congressionally reserved lands had a lower average of area in large trees (mean 28 percent) than late-successional reserves (36 percent) and only slightly higher than matrix lands (26 percent). This lower average is likely due to the fact that a large proportion of congressionally reserved watersheds are in higher elevations where the natural vegetation is less likely to attain an average tree diameter of 20 inches (ideally we would stratify size expectations by site vegetation potential, a task we have prioritized for future development—see chapter 5 Emerging Issues). The key watershed classification does not appear to have this same elevation correlation, so key watersheds exhibit the expected pattern having more large tree area than nonkey watersheds.

Figure 41—Percent of watershed area in late-successional (LSOG; conifer quadratic mean diameter at breast height \geq 20 inches and canopy cover \geq 10 percent) by land use allocation, 2008 (Reserved = Congressional reserves, LSR = late-successional reserves).

The average trend change in LSOG for all classes was small (1 percent or less) and positive, except for a slight decline (-0.1 percent) on matrix lands. The trend ranges were also quite similar, except for matrix lands, which did not have the higher (>10 percent) increases shown by some watersheds in all the

other classes (Figure 42). As with road density, changes in the overall distributions were slightly more visible when broken out by land use class, but no other significant pattern is apparent.

Figure 42—Change in percentage of watershed in late-successional old-growth (LSOG; diameter at breast height \geq 20 inches and canopy cover \geq 10 percent) from 1994 to 2008 by land category.

Disturbance

An alternative to judging vegetation by its current condition, and one that avoids the need for setting "natural" size or cover thresholds, is to look at the amount of vegetation disturbance in a watershed. A new disturbance data set for the full NWFP area based on satellite imagery changes only recently became available (Kennedy et al. 2010), so none of the provincial models currently incorporate disturbance metrics. The attributes we derived for this analysis are the percentage of watershed area disturbed by harvest and fire, summed over 5-year periods and averaged over the 1,379 watersheds (since the latest data available was 2008, the last period comprises only 4 years; see "Methods" section for further details).

By individual watershed, the cumulative percentage area disturbed during the periods ranged from 0 to 22 percent for fire and 0 to 18 percent for harvest. Because disturbance area in most watersheds was zero or very low, the average area disturbed over all 1,379 watersheds was considerably smaller: 0.03 to 1.79 percent for fire and 0.21 to 1.41 percent for harvest (Figure 43). As a regional average, harvest declined dramatically from the first period (1985-1989) to the second (1990-1994), even before the NWFP initiation. There was a moderate uptick to 0.41 percent harvested in the latest period (2004-2008); however, the level is less than one-third of the first period. Fire was more variable. A smaller average percentage was lost to fire than harvest in the first two periods, whereas fire losses exceeded harvest in periods three and four (with a notable jump in period four owing to the Biscuit Fire and others).

Figure 43—Average percentage of watershed area affected by harvest and fire disturbances 1985 to 2008.

For comparison, the watershed evaluation models of a number of the provinces use a measure of percentage of forest land in early seral stage (as represented by average tree diameter less than 5 inches) to approximate disturbance. The evaluation criteria in these models rate 5 percent or less disturbance as good condition and greater than 25 percent as poor. Comparing these thresholds to the sum of harvest and fire disturbance percentages shows that only 45 watersheds (3 percent) experienced a greater than 5 percent disturbance and only 11 watersheds (0.8 percent) would score in the very low category (model score ≤ -0.6). In contrast to road density and large tree alternative metrics, evaluating this disturbance metric using the thresholds in our watershed models would produce a considerably more positive distribution of condition scores than was produced by the vegetation component of our provincial watershed evaluation models, where 6 percent of the watersheds fell into the poor category.

By Land Use Category

Disturbance levels, averaged over all watersheds, show some distinct patterns by land use class (Figure 44). On a per watershed basis, pre-NWFP (prior to 1994) vegetation losses were primarily due to harvest on matrix (timber production) lands. In contrast, the greatest post-NWFP losses have been mainly due to fire and mostly on congressionally-reserved lands and late-successional reserves (LSR). Note that harvest associated with reserved watersheds does not indicate harvest in reserved areas, since the harvest category includes small fires and blow down, and watersheds classified as reserve may have portions of non-reserve lands (see “Methods” section for further details). Average harvest levels are low in both periods in the reserved category, but decline dramatically on the LSR class and somewhat less on matrix lands. Sharp declines are also seen on key and nonkey watersheds, but the difference between the two is slight. Average losses from fire went up in all land categories from the pre-NWFP to post-NWFP period, but the increase was noticeably less on matrix lands.

Figure 44—Average yearly pre- and post-Northwest Forest Plan (NWFP) fire and harvest disturbance levels by land use allocation, and for key and nonkey

watersheds (Reserved = congressionally reserved, LSR = late-successional reserve).

Upslope/Riparian Versus Inchannel Conditions

The shapes of the overall score distributions for upslope/riparian and inchannel conditions are shown in (Figure 45). The majority of watersheds scored in the moderate and high condition categories. Inchannel scores were more tightly grouped around the center of the distribution, with fewer scores in the very low, low, and very high categories.

Figure 46 displays the inchannel values paired with the corresponding upslope/riparian scores by watershed, along with the best fit regression line. The test of whether the slope of the line was zero had a two-sided p-value of 0.12, indicating a possible relationship, but the strength of this relationship was extremely weak ($R^2 = 0.01$).

Figure 45—Watershed versus inchannel current status scores. These curves show the data in a continuous manner, rather than by data bins (i.e., histograms).

Figure 46—Upslope/riparian versus inchannel current status scores for 193 watersheds with both data types available.

Chapter 4: Discussion

Watershed Trend

Did the Northwest Forest Plan (NWFP) succeed in maintaining and improving watershed condition? Examining the trend data expressed as a continuous distribution (Figure 27) and score categories (Figure 29), both suggested the answer is yes. The overall trend was clearly positive with 69 percent of watersheds trending up versus 23 percent showing declines (Table 6). Although a portion of these shifts was likely due to errors inherent in the satellite imagery classification process, a more conservative estimate still showed a similarly positive ratio, with 10 percent of watersheds increasing more than this threshold versus 4 percent decreasing. In most cases, these larger positive changes depended on the combined effects of natural vegetation growth and road decommissioning efforts, and especially road decommissioning in landslide-prone areas.

A greater proportion of the positive changes in watershed condition occurred on late-successional reserve (LSR) and matrix lands than on reserved lands, and the mean score for reserved lands actually declined slightly. Considering that the reserved class is already generally in good condition with respect to roads and vegetation, it is not surprising that the larger increases were found in the other allocations. Our analysis of contributing factors revealed that declining watershed condition scores were driven almost exclusively by vegetation losses, and the alternative disturbance indicator showed that fire was the main driver of these losses. Parallel to a similar finding in the late-successional old-growth monitoring report (Moeur et al., in press), vegetation losses due to fire were much higher on reserved lands and on key watersheds than on the other land use categories. These disproportionate vegetation losses were one reason average improvement on key watersheds was only slightly greater than on nonkey watersheds. Another reason key watersheds did not show more improvement was that, although more roads were

decommissioned on key watersheds, these numbers were often too small in terms of overall density to raise condition scores.

Upslope/Riparian and Inchannel Current Status

In addition to the satellite vegetation classification and geographic information system (GIS)-based road data used to evaluate upslope and riparian condition, this report also assessed current watershed condition status using field sampling of multiple inchannel attributes. The shapes of the overall distributions of these scores (Figure 45), showed the majority of watersheds scoring in the moderate and high condition categories. In terms of land use categories, both upslope and instream condition scores generally followed a pattern consistent with the amount of allowable management (i.e., timber harvest). Matrix lands had the lowest upslope/riparian and inchannel scores. LSR scores were higher, followed by reserved lands with the highest scores. Key watersheds also were in considerably better condition on average than nonkey watersheds.

Model Validation

We looked at a number of simple metrics, i.e., road density, late-successional old-growth, and vegetation disturbance, to provide an alternative view and partial check on our results. None of these metrics closely matched the results from our more complex provincial evaluation models, but this was not unexpected given that each metric provides only a partial picture of watershed influences and does not account for the natural variability in conditions across the broad monitoring area. Nevertheless, these metrics provided some additional insights relevant to watershed conditions.

When viewed as a regional aggregate, road density has changed little over the life of the NWFP. However, dramatic changes have been accomplished in targeted watersheds, for example Lower Fish Creek in the western Cascades dropped road density from 3.3 mi/mi² in 1994 to 0.8 mi/mi² in 2008 through decommissioning 118 miles of roads. This targeting of road decommissioning appeared to follow

the intent of the Aquatic Conservation Strategy, in that the rate in key watersheds was double that in nonkey watersheds. Similarly, in broad aggregate the area in late-successional old-growth has changed little over the 15-year period, although the latest estimates suggest a slight decline in area (Moeur et al., in press). In contrast, the AREMP model vegetation attributes scores generally increased; however, this was not inconsistent because many of the AREMP indicators were based on earlier seral stages. The utility of LSOG as a metric for aquatic monitoring appeared low because of this mismatch in seral stage and the natural variability of vegetation potential across the NWFP area. In contrast, the vegetation disturbance metric showed considerable promise for contributing to aquatic assessment. Many of the expert groups wished to assess disturbance but had to rely on a simple mean tree diameter metrics (cover, quadratic mean diameter) as a proxies, and these measures are difficult to calibrate to the variability in natural vegetation potential. The disturbance metric also revealed that average losses from fire went up in all land categories from the pre-NWFP to post-NWFP period, especially in reserved lands and key watersheds. Average timber harvest levels declined sharply from pre-NWFP levels on both key and nonkey watersheds, but the decline in both classes was similar, suggesting that the key watershed designation did not have a major influence on harvest levels at the regional scale.

Reeves et. al. (2004 p. 6) stated, “To be meaningful, a monitoring program should provide insights into cause-and-effect relations between environmental stressors and anticipated ecosystem responses.” Although the shapes of the overall score distributions looked similar (Figure 45), we found little correspondence at the individual watershed level (Figure 46). Reid and Furniss (1998) summarized a number of past efforts which illustrate the difficulty of showing strong relationships based on a short period:

Gilbert (1917), for example, demonstrated that the aggradational front for sediment from long-discontinued hydraulic mining in the Sierra Nevada foothills was still progressing through the Central Valley. Similarly, Trimble (1983) found that the form of low-order

channels in the Coon Creek watershed of Wisconsin responded quickly to agricultural impacts of the last century, while downstream channels are still adjusting today. In Redwood Creek, California, logging-related impacts on channel morphology only became evident when major storms occurred, and parts of the channel are only now responding to sediment introduced by the 1964 flood (Madej and Ozaki 1996). Similarly, Frissell et al. (1997) note that it may take another major flood to allow channel morphology to recover, even if upslope practices have indeed improved to the point that recovery is possible.

Other reasons for difficulty in finding relationships between stressors and response are a lack of direct upslope-inchannel links, and upstream influences (see sidebar “Modeling Inchannel Variables,” pg. XX). However, a few recent studies have found statistical relationships between a few simple measures of upslope/riparian conditions and corresponding inchannel attributes. Lee et al. (1997) found a relationship between road density and the status of resident salmonids in the interior Columbia basin. Kaufmann and Hughes (2006) were able to factor out major natural landscape differences (basin size, stream gradient, lithology) and found relationships between road density, riparian condition, and an index of stream macroinvertebrates. Al-Chokhachy et al. (2010) have further developed statistical methods for filtering out natural landscape variability, and they found a relationship between upslope road density and an index of inchannel physical conditions.

Sidebar 5--Modeling Inchannel Variables

Evaluating Dynamic Processes and the Role of Fire

Fire has surpassed timber harvest as a dominant influence on vegetation on federal lands in the NWFP area (Figure 44; Healy et al. 2008; Moeur et al., in press). The NWFP watershed condition models currently do not distinguish between vegetative disturbances: fire and harvest (and other disturbances) affect condition simply through their impacts on canopy cover and average tree size

metrics. Reviewers of this report have argued that fire, as a natural process, should not be considered in the same way, and we acknowledge that different types of disturbances have different associated effects and ecological legacies. New metrics based on disturbance types (Kennedy et al. 2010) rather than vegetation status may enable further distinction among disturbances in future assessments. However, fire does often have short- to medium-term negative impacts on fish habitat from increases in mass wasting, sedimentation, solar radiation, and water temperatures (Burton 2005, Dunham et al. 2003), although it also plays a longer term role in renewing habitat attributes (Reeves et al. 1995). Given the possibility of a changing climate, along with changes in disturbance regimes (Dale et al. 2001), we believe it is important to capture all disturbance effects on watershed conditions, rather than to assume that some are natural and therefore just part of a natural baseline or reference condition.

Management Implications

We can draw some management implications from this type of broad-scale monitoring and assessment, but it must be realized that restoration actions are planned and implemented with finer-grained information that is much more sensitive to the local context. At the regional level under current management practices, watershed condition appears to be improving, with widespread, small score increases due to maturing vegetation and larger but localized increases from restoration actions. According to our assessment methods, road decommissioning in landslide prone areas is the most effective action for raising watershed scores. Such decommissioning has the dual benefit of reducing more direct road impacts, such as altered hydrology and erosion, as well as the indirect effect on reducing landslide risks. Removing roads in riparian buffers, especially those with associated stream crossings, is another strategy that yields dual benefits. Reducing overall road density, even in lower risk areas, can be effective since roads are weighted more heavily than vegetation in a number of the provincial expert models. However, because these models incorporate certain thresholds, changes in density that remain above the upper threshold (which varies by province from 2.4 to 4 mi/mi²) will not affect evaluation

THE FINDINGS AND CONCLUSIONS IN THIS REPORT ARE IN PRESS AND SUBJECT TO CHANGE PRIOR TO FORMAL DISSEMINATION BY THE AGENCIES AND SHOULD NOT BE CONSTRUED TO REPRESENT AGENCY DETERMINATION OR POLICY.

scores. Based on this evaluation technique, there is an advantage to targeting restoration to watersheds at or below the threshold, where each mile removed will benefit scores, rather than those watersheds that are the most heavily roaded.

Similar management implications can be drawn from the assessment relative to vegetation. Riparian areas are generally accorded more weight in the models, so restoration activities in these areas will have greater positive effects on evaluation scores. Thinning in riparian reserves is one restoration strategy (Chan et al. 2004), and, if targeted to smaller diameter trees, it will increase scores by raising the mean tree diameter. However, two of the provinces also include riparian canopy cover metrics, and reducing cover below these thresholds could have a corresponding negative effect. While upslope vegetation generally carries less weight, it also tends to have lower thresholds in terms of mean tree diameter expectations. Thus revegetating burned or harvested areas can yield results more quickly than in riparian areas, again especially in landslide prone areas. Reducing stand-replacing fires, via fuel treatments or suppression, will also have positive effects under our evaluation approach; however, suppression alone can have negative effects by reducing mean tree diameters and increase longer-term risks of more severe fires. Finally, similar to the concern expressed in Moeur et al. (in press) about old-growth reserves, the unpredictable nature and dynamic role of fire may have implications for the static reserves approach that lies behind the designated set of key watersheds.

Improving the Assessment

We anticipate working on the following issues to improve confidence in these assessment models: improving the data and error estimates, improving the validity of the evaluation criteria, and comparison to other assessments.

Improving data quality and quantity and better understanding and communicating measurement error are fundamental to advancing these assessments. We continue to work on obtaining attributes important

to salmonid habitat, such as fish passage barriers, and broader aquatic processes, which are currently missing from these models owing to lack of regional data sets (see chapter 5 Emerging Issues).

Quantifying and communicating the effect of measurement errors on watershed condition estimates is also needed. Appendix 6 presents information on data quality estimates for a number of attributes, but more detail is clearly desirable, especially for the upslope indicators. Estimating uncertainty levels in the results of this type of multimetric, nonstatistical model is challenging, but there are feasible methods. One such method is a Monte Carlo approach, which runs the model multiple times drawing inputs from the expected distributions of the attribute values each time (Refsgaard et al. 2007).

Evaluation criteria, the standards against which the input data are judged, are perhaps the most critical and sensitive piece of the assessment process. The criteria used are derived from and validated by expert workshops. It is often challenging to link these criteria back to specific empirical studies, but future work in this area would help to improve model validity (Gordon and Gallo 2011). Reeves and Duncan (2009) caution against using fixed standards because of natural variability owing to environmental gradients and even within sites in the same ecological space. The NWFP models include broad-scale criteria adjustments by ecological province and some finer scale gradients; however, a more empirical approach would be desirable. As discussed above, a few recent studies have factored in environmental gradients by using multiple regression techniques (Al-Chokhachy et al. 2010; Kaufmann and Hughes 2006).

Comparison to results from other assessments also provides an opportunity for further validation. New biotic indices, which combine both abundance and diversity measures, show promise, and there are a number of available data sets in the Pacific Northwest with the necessary information (Hubler 2008, Hughes et al. 2004, Whittier et al. 2007). Other recent assessments have used these biotic indices, although they have not attempted correlations with the other biophysical indicators (Hubler et al. 2009,

Mulvey et al. 2009, Oregon DEQ 2004). Testing for relationships between our expert-based watershed assessments and these biotic data sets is another potential approach to better model validation.

DRAFT

THE FINDINGS AND CONCLUSIONS IN THIS REPORT ARE IN PRESS AND SUBJECT TO CHANGE PRIOR TO FORMAL DISSEMINATION BY THE AGENCIES AND SHOULD NOT BE CONSTRUED TO REPRESENT AGENCY DETERMINATION OR POLICY.

Chapter 5: Emerging Issues

Issues related to the condition of watersheds are expected to emerge as our understanding of processes affecting watershed condition evolves. These issues will be incorporated into future iterations of decision-support models as needed information becomes available. The following highlight current emerging issues.

- **Fish passage**—Millions of dollars have been spent by the Forest Service (FS) and Bureau of Land Management (BLM) over the past 15 years restoring fish passage where it was blocked by culverts at road stream crossings. The FS spent \$2,090,000 in 2009 alone for improving fish passage in Oregon and Washington (USDA FS 2010). However we had no way to consider the effect of blocked fish passage on watershed condition, or improved watershed conditions when fish passage was improved or restored because fish passage assessment data were unavailable (see “Fish Passage at Culverts” sidebar page XX). When culvert locations and culvert fish passage assessment data become available we intend to use the amount of fish habitat available to fish as part of our watershed condition assessment.

Sidebar 6—Fish Passage

- **Complete road layer**—Road data are currently stored separately by the Forest Service and BLM. The two agencies manage their road information with different data structures and attributes which make combining the data very challenging. Also, non-system roads on federal lands are often not included and private lands have very poor road information (see sidebar page XX). Although roads are known to have a major affect on watershed condition (see Daigle 2010), it will remain difficult to analysis watershed conditions across ownership boundaries until agencies utilize more compatible information systems.

Sidebar 7—Road Data Challenges

- **All lands approach**—Recent Forest Service policy statements have emphasized looking at agency actions in the context of the broader landscape (Tidwell 2010). The use of geographic information system (GIS) and satellite imagery should eventually allow for a consistent assessment of watershed condition across all land ownerships. For example, vegetation data are now consistent for all ownerships through the use of satellite imagery. However, road data are not consistent, and

including nonfederal lands could affect watershed condition scores in two ways—with opposite results:

4. If nonfederal lands within a mixed ownership watershed (classified by the majority landowner) has lots of roads shown (Figure 47), this could result in an overall low (poor) watershed condition score even if the majority of the watershed consists of federal lands with few or no roads.
5. Nonfederal land road layers are known to underrepresent the number of roads in watersheds. Gallo et al. 2005 found that 37 percent of the roads on nonfederal lands were missing on the agency layer used, compared to 10 percent on BLM lands and 15 percent on FS lands. If a large number of roads on nonfederal lands are missing on a watershed's road layer, the watershed could receive an artificially high (good) watershed condition score.

Figure 47—Federal lands in the headwaters of the Clearwater River watershed (Mount Baker-Snoqualmie National Forest) are within a wilderness. Nonfederal lands in the upper half of the same watershed are managed for timber production, and that area is heavily roaded. Including the nonfederal lands would misrepresent the watershed score if the watershed is classified as a federal watershed.

- **Inadequate flow**—Dams and irrigation practices have altered flow regimes in many watersheds. Altered flow regimes include changes in timing and magnitude of peak flow and dewatering that result in insufficient flow during summer (Gallo et al. 2005).
- **Invasive aquatic species**—The effects of invasive and exotic aquatic species on the aquatic biotic community are usually detrimental to native species. However, an overall, accurate assessment of invasive species distributions is currently unavailable. Although invasive species data bases currently exist for Oregon, Washington, California, U.S. Geological Services, U.S. Fish and Wildlife Service, National Park Service, and FS, the ability to easily and quickly share data among agencies is still being developed.
- **Fire**— Fire condition class, which identifies how vegetation conditions have deviated from historical fire conditions could be included in our watershed condition models. However, the effect of fire suppression and stand-replacing burns on vegetation and stream reach conditions is still largely unknown, especially for forests that tend to have catastrophic fires several centuries apart (Franklin and Dyrness 1973), which is common throughout the Northwest Forest Plan (NWFP) area. Fire condition class is currently being mapped for all federal lands in the NWFP area.

THE FINDINGS AND CONCLUSIONS IN THIS REPORT ARE IN PRESS AND SUBJECT TO CHANGE PRIOR TO FORMAL DISSEMINATION BY THE AGENCIES AND SHOULD NOT BE CONSTRUED TO REPRESENT AGENCY DETERMINATION OR POLICY.

- **Climate change**—Climate change is likely to affect our measures of watershed condition in a number of ways, from stream temperatures to morphology changes from increased storms to vegetation changes. Further thought and discussion are needed to decide whether the program should attempt to distinguish between direct management and climate change effects in the future.
- **Restoration projects**—A link between restoration projects and the monitoring program has yet to be established. For example, the decision-support models are sensitive only to road decommissioning, because road improvements are not tracked consistently in a database. Assigning and tracking other road attributes (such as water bar installation and culvert replacement) is needed to better reflect how roads affect the condition of watersheds (Gallo et al. 2005).
- **Unified model**—Aquatic province watershed condition models are similar in the attributes being evaluated but their model structures differ (see app. 4). A “unified model structure” may be possible, but still allow evaluation criteria to vary based on contextual information, e.g., geology, precipitation zone, elevation. This would have the advantage of simplifying the effort needed to keep track of different model structures.
- **Distribution of watershed condition scores**—The desired distribution of watershed condition scores was not identified by the strategy. This information would allow field unit specialists to determine if planned disturbances will move watershed conditions outside the range of natural variation (Gallo et al. 2005).
- **Model validation and links between upslope, inchannel and biological indicators**—The alternative metrics we looked at in this report do not appear well-suited for validating our models. New biotic indices, which combine both abundance and diversity measures, show promise, and there are a number of available data sets in the Pacific Northwest with the necessary information (Hubler 2008; Hughes et al. 2004, Whittier et al. 2007). Testing for relationships between our expert-based watershed assessments and these biotic data sets could be a more promising approach to model validation and linking upslope and inchannel models.
- **National watershed condition assessment coordination**—The FS is developing a consistent national process for watershed condition assessment to improve the system for rating watershed condition. The Aquatic and Riparian Effectiveness Monitoring Program (AREMP) already evaluates some, but not all, of the indicators proposed for use in a national assessment (USDA

2010). One option being considered is to have AREMP include additional attributes identified in national assessment guidelines as part of regional watershed condition assessments.

- **Management review**—An AREMP management review is proposed for 2011 to determine if any changes are appropriate after 10 years of implementing the program. Suggested questions to be addressed include:

- What are the relevant management questions today?
- The watershed condition monitoring program uses a fairly sophisticated decision-support tool; is it a good fit?
- Is the ongoing amount of effort still needed to answer the current management questions of status and trend? Is it too rigorous?
- What new technologies exist that can help answer management questions (e.g., satellite imagery, and LIDAR)? Can these be utilized in lieu of instream sampling (to reduce program costs and safety risks to field crews)?
- What are AREMP's strengths and weaknesses?
- What can or do we use AREMP information for in addition to NWFP monitoring questions (e.g., project support, forest plan revisions, Endangered Species Act consultation, impaired water listings, watershed assessments, watershed condition framework [FS new national effort], and Survey and Manage riparian species detection)?
- What are program costs and what is the current staffing level?

Sidebar 8—Support to Local Units – Protocol comparison test and aquatic invasive species

Sidebar 9—Support to Local Units – Restoration project survey support and bull trout reintroduction

References

- Al-Chokhachy, R.; Roper, B.B.; Archer, E.K. 2010.** Evaluating the status and trends of physical stream habitat in headwater streams within the interior Columbia River and upper Missouri River basins using an index approach. *Transactions of the American Fisheries Society*. 139(4): 1041-1059.
- Alegria, J.; Hyzer, M.; Mulder, B. [et al.]. 1995.** Guidance for implementation monitoring for management of habitat for late-successional and old-growth related species within the range of the northern spotted owl. Draft. On file at: Regional Ecosystem Office, 333 SW First Avenue, Portland, OR 97208.
- Andersen, H.; Lanigan, S.H. 2009.** Aquatic and riparian invasive species survey. <http://www.reo.gov/monitoring/reports/watershed-reports-publications.shtml>. October 2010.
- AREMP 2009.** Field protocol manual: Aquatic and riparian monitoring program. <http://www.reo.gov/monitoring/reports/watershed/2009FieldProtocolFinal.pdf> (Jan 10, 2010)
- Bailey, R.G.; Avers, P.E.; King, T.; McNab, W.H., eds. 1994.** Ecoregions and subregions of the United States [map 1:7,500,000]. With supplementary table of map unit descriptions by McNab, W.H.; Bailey, R.G., comps., eds. Washington, DC: U.S. Department of Agriculture, Forest Service.
- Baker, D.; Ferguson, G.; Palmer, C.; Tolle, T. 2005.** Northwest Forest Plan—the first 10 years (1994–2003): implementation monitoring, summary of regional interagency monitoring results. [R6-RPM-TP-04-2005](#). U.S. Department of Agriculture, Forest Service, Pacific Northwest Region. 24 p.
- Benda, L.E.; Miller, D.J.; Dunne, T. [et al.]. 1998.** Dynamic landscape systems. In: Naiman, R.J.; Bilby, R.E., eds. *River ecology and management: lessons from the Pacific coastal ecoregion*. New York: Springer-Verlag: 261–288.

- Berg, L. 1982.** The effect of exposure to short-term pulses of suspended sediment on the behavior of juvenile salmonids in Hartman, G.G.; Oswald E.T.; Hetherington, E.D. [et al.], [eds.] Proceedings of the Carnation Creek workshop: a ten-year review. Department of Fisheries and Oceans, Pacific Biological Station Nanaimo, Canada: 177-196.
- Blaustein, A.R.; Wake, D.B.; W.P. Sousa. 1994.** Amphibian declines: judging stability, persistence, and susceptibility of populations to local and global extinctions. *Conservation Biology*, 8: 60–71.
- Bown, K. [and others]. 2002 (4 October).** Letter to USDA Forest Service Forest Supervisors and USDI Bureau of Land Management District Managers and Field Managers. Recommended Monitoring Interview Agenda. On file with: Resource Planning and Monitoring, Pacific Northwest Regional Office, 333 SW First Ave., Portland, OR 97208.
- Bryant, M.D.; Caouette, J.P.; and Wright, B.E. 2004.** Evaluating stream habitat survey data and statistical power using an example from southeast Alaska. *North American Journal of Fisheries Management*. 24: 1353– 1362.
- Bryce, S.A.; Omernick, J.; Larsen, D. 1999.** Ecoregions: a geographic framework to guide risk characterization and ecosystem management. *Environmental Practice*. 1: 141–153.
- Burton, T.A. 2005.** Fish and stream habitat risks from uncharacteristic wildfire: Observations from 17 years of fire-related disturbances on the Boise National Forest, Idaho. *Forest Ecology and Management*. 211(1-2): 140-149.
- Chan, S.; Anderson, P.; Cissel, J.; Lateen, L.; Thompson, C. 2004.** Variable density management in riparian reserves: lessons learned from an operational study in managed forests of western Oregon, USA. *Forest Snow and Landscape Research*. 78(1/2): 151-172.
- Charnley, S., tech. coord. 2006.** Northwest Forest Plan—the first 10 years (1994–2003):

socioeconomic monitoring results. Gen. Tech. Rep. PNW-GTR-649. Portland, OR: U.S. Department of Agriculture, Forest Service, Pacific Northwest Research Station. 6 vol.

Cohen, W.B.; Yang, Z.; Kennedy, R. 2010. Detecting trends in forest disturbance and recovery using yearly Landsat time series: 2. TimeSync -- Tools for calibration and validation. *Remote Sensing of Environment*. 114(12): 2911-2924.

Consulting Science Team. 2001. Umpqua land exchange project multi-resource land allocation model handbook. Portland, OR: Foundation for Voluntary Land Exchanges.
<http://www.or.blm.gov/umpqua/documents.htm>. (August 28, 2010).

Daigle, P. 2010. A summary of the environmental impacts of roads, management responses, and research gaps: literature review. *British Columbia Journal of Ecosystems and Management* 10(3): 65-89.

Dale, V.H.; Joyce, L.A.; McNulty, S.; Neilson, R.P.; Ayres, M.P.; Flannigan, M.D.; Hanson, P.J.; Irland, L.C.; Lugo, A.E.; Peterson, C.J.; Simberloff, D.; Swanson, F.J.; Stocks, B.J.; Michael Wotton, B. 2001. Climate change and forest disturbances. *BioScience*. 51(9): 723-734.

Davis, R.J., tech. coord. 2011. Northwest Forest Plan—Status and Trend of Northern Spotted Owl Populations and Habitats from 1994 to 2008. Gen. Tech. Rep. Portland, OR: U.S. Department of Agriculture, Forest Service, Pacific Northwest Research Station.

Dunham, J.B.; Young, M.K.; Gresswell, R.E.; Rieman, B.E. 2003. Effects of fire on fish populations: landscape perspectives on persistence of native fishes and nonnative fish invasion. *Forest Ecology and Management*. 178(1-2): 183-196.

Dunham, J.; Gallo, K., 2008. Assessing the Feasibility of Native Fish Reintroductions: A Framework and Example Applied to Bull Trout in the Clackamas River, Oregon: Reston, Virginia, U.G. Geological Survey, Open-File Report 2008-1007, 23 p.

THE FINDINGS AND CONCLUSIONS IN THIS REPORT ARE IN PRESS AND SUBJECT TO CHANGE PRIOR TO FORMAL DISSEMINATION BY THE AGENCIES AND SHOULD NOT BE CONSTRUED TO REPRESENT AGENCY DETERMINATION OR POLICY.

Dwyer, W.L. 1994. Seattle Audubon Society, et al. v. James Lyons, Assistant Secretary of Agriculture, et al. Order on motions for Summary Judgment RE 1994 Forest Plan. Seattle, WA: U.S. District Court, Western District of Washington.

Forest Ecosystem Management Assessment Team [FEMAT]. 1993. Forest ecosystem management: an ecological, economic, and social assessment. Portland, OR: U.S. Department of Agriculture; U.S. Department of the Interior [et al.]. [Irregular pagination].

Franklin, J.F.; Dyrness, C.T. 1973. Natural vegetation of Oregon and Washington. Gen. Tech. Rep. PNW-8. Portland, OR: USDA Forest Service Pacific Northwest Research Station.

Frissell, C.A.; Liss, W.J.; Greswell, R.E.; Nawa, R.K.; Ebersole, J.L. 1997. A resource in crisis: changing the measure of salmon management in Strouder, D.J.; Bisson, P.A.; Naiman, R.J., Pacific salmon and their ecosystems. New York, Chapman and Hall: 411-444.

Gallo, K.; Lanigan, S.H.; Eldred P.; Gordon S.N.; Moyer C. 2005. Northwest Forest Plan – the first 10 years (1994-2003): preliminary assessment of the condition of watersheds. Gen. Tech. Rep. PNW-GTR-647. Portland, OR: U.S. Department of Agriculture, Forest Service, Pacific Northwest Research Station. 133 p.

Gallo, K.; Moyer, C.; Lanigan, S. 2002. Interagency regional monitoring Northwest Forest Plan aquatic and riparian effectiveness-monitoring program. 2002 annual report.

http://www.reo.gov/monitoring/report_show.php?show=watershed. (October 25, 2010)

Gilbert, G.K. 1917. Hydraulic-mining debris in the Sierra Nevada. Professional Paper 105. Department of Interior, Geological Survey. Washington, DC: 154p.

Grant, G.E.; Lewis, S.L.; Swanson, F.J.; Cissel, J.H.; McDonnell, J.J. 2008. Effects of forest practices on peak flows and consequent channel response: a state-of-science report for western Oregon

and Washington. Gen. Tech. Rep PNW-GTR-760. Portland, OR: U.S. Department of Agriculture, Forest Service, Pacific Northwest Research Station. 76 p.

Gucinski, H.; Furniss, M.J.; Ziemer, R.R.; Brookes, M.H. 2001. Forest roads: a synthesis of scientific information. Portland, OR: USDA Forest Service Pacific Northwest Research Station. ~-**Haynes, R.W.; Bormann, B.T.; Lee, D.C.; Martin, J.R., tech. eds. 2006.** Northwest Forest Plan—the first 10 years (1994–2003): synthesis of monitoring and research results. Gen. Tech. Rep. PNW-GTR-651. Portland, OR: U.S. Department of Agriculture, Forest Service, Pacific Northwest Research Station. 292 p.

Haynes, R.W.; Bormann, B.T.; Lee, D.C.; Martin, J.R.; tech. eds. 2006. Northwest Forest Plan—the first 10 years (1994–2003): synthesis of monitoring and research results. Portland OR: USDA Forest Service, Pacific Northwest Research Station. 292 pp.

Healey, S.; Cohen, W.; Spies, T.; Moer, M.; Pflugmacher, D.; Whitley, M.; Lefsky, M. 2008. The relative impact of harvest and fire upon landscape-level dynamics of older forests: lessons from the Northwest Forest Plan. *Ecosystems*. 11(7): 1106-1119.

Hemstrom, M.; Spies, T.; Palmer, C. [et al]. 1998. Late-successional and old-growth forest effectiveness monitoring plan for the Northwest Forest Plan. Gen. Tech. Rep. PNW-GTR-438. Portland, OR: U.S. Department of Agriculture, Forest Service, Pacific Northwest Research Station. 37p.

Herlihy A. 2009. Development of an index of biotic integrity for forested Pacific Northwest stream macroinvertebrates. Unpublished report. On file with: AREMP/RPM, USDA Forest Service R6, 333 SW First Ave, Portland OR 97204. 17 pp.

Hilsenhoff, W.L. 1988. Rapid field assessment of organic pollution with a family level biotic index. *The Journal of the North American Benthological Society*. 7: 65-68.

THE FINDINGS AND CONCLUSIONS IN THIS REPORT ARE IN PRESS AND SUBJECT TO CHANGE PRIOR TO FORMAL DISSEMINATION BY THE AGENCIES AND SHOULD NOT BE CONSTRUED TO REPRESENT AGENCY DETERMINATION OR POLICY.

- Hubler, S. 2008.** PREDATOR: development and use of RIVPACS-type macroinvertebrate models to assess the biotic condition of wadeable Oregon streams. Hillsboro, OR: Oregon Department of Environmental Quality, Laboratory and Environmental Assessment Division. 51p.
- Hubler, S.; Miller, S.; Merrick, L.; Leferink, R.; Borisenko, A. 2009.** High level indicators of Oregon's forested streams. Hillsboro, OR: Oregon Dept. Environmental Quality, Laboratory and Environmental Assessment Division.
- Huff, M.H.; Raphael, M.G.; Miller, S.L. [et al.], tech. coords. 2006 .** Northwest Forest Plan—the first 10 years (1994–2003): status and trends of populations and nesting habitat for the marbled murrelet. Gen. Tech. Rep. PNW-GTR-650. Portland, OR: U.S. Department of Agriculture, Forest Service, Pacific Northwest Research Station. 149 p.
- Hughes, R.M.; Howlin, S.; Kaufmann, P.R. 2004.** A biointegrity index (IBI) for coldwater streams of western Oregon and Washington. *Transactions of the American Fisheries Society*. 133(6): 1497-1515.
- Karr, J.R., Chu, E.W. 1999.** Restoring life in running waters: better biological monitoring. Washington DC: Island Press.
- Kaufmann, P.R.; Hughes, R.M. 2006.** Geomorphic and anthropogenic influences on fish and amphibians in Pacific Northwest coastal streams. In: Hughes, R.M.; Wang, L.; Seelbach, P.W., eds. *American Fisheries Society symposium: landscape influences on stream habitats and biological assemblages*. Bethesda, MD: American Fisheries Society: 429-455.
- Kaufmann, P.R.; Levine, P.; Robison, E.G. [et al.] 1999.** Quantifying physical habitat in wadeable streams. EPA/620/R-99/003. Washington, DC: U.S. Environmental Protection Agency. 102 p.
- Keeney, R.L.; Raiffa, H. 1976.** Decisions with multiple objectives : preferences and value tradeoffs. New York: Wiley. xxviii, 569 p. pp.
- Kennedy, R.E.; Cohen, W.B.; Schroeder, T.A.; 2007.** Trajectory-based change detection for automated characterization of forest disturbance dynamics. *Remote Sensing of Environment*. 110: 370-386.

- Kennedy, R.E.; Yang, Z.; Cohen, W.B. 2010.** Detecting trends in forest disturbance and recovery using yearly Landsat time series: 1. LandTrendr -- Temporal segmentation algorithms. *Remote Sensing of Environment*. 114(12): 2897-2910.
- Kershner, J.L.; Roper B.B.; Bouwes, N.; Henderson, R.C.; Archer, E. 2004.** An analysis of stream habitat conditions in reference and managed watersheds on some federal lands within the Columbia River basin. *North American Journal of Fisheries Management*. 24: 1363-1375
- Knighton, A.D. 1999.** Downstream variation in stream power. *Geomorphology*. 29: 293–306.
- Lanigan, S.H.; Moyer, C.; Gallo, K.; Gruendike, P. 2007.** Aquatic and Riparian Effectiveness Monitoring Program. Interagency Monitoring Program – Northwest Forest Plan Area. Annual technical report. On file with: Resource Planning and Monitoring, USDA Forest Service, Pacific Northwest Regional Office, 333 SW First Avenue, Portland, OR 97208.
- Lee, D.C.; Sedell, J.R.; Rieman, B.E.; Thurow, R.F; Williams, J.E. 1997.** BROADSCALE ASSESSMENT OF AQUATIC SPECIES AND HABITATS. An assessment of ecosystem components in the interior Columbia Basin and portions of the Klamath and Great basins. Gen. Tech. Rep. PNW-GTR-405, Portland, OR: U.S. Department of Agriculture, Forest Service, Pacific Northwest Research Station. Vol. 3, chapter 4.
- LEMMA. 2010.** GNN Accuracy Assessment Report. USDA Forest Service, PNW Research Station, Landscape Ecology, Modeling, Mapping and Analysis Program. Internet:
http://www.fsl.orst.edu/lemma/main.php?project=common&id=mr&model_region=200&ref=nwfp
(Mar 20, 2011)
- Lint, J., tech. coord. 2005.** Northwest Forest Plan—the first 10 years (1994–2003): status and trends of northern spotted owl populations and habitat. Gen. Tech. Rep. PNW-GTR-648. Portland, OR: U.S. Department of Agriculture, Forest Service, Pacific Northwest Research Station. 176 p.

- Lint, J.; Noon, B.; Anthony, R. [et al.]. 1999.** Northern spotted owl effectiveness monitoring plan for the Northwest Forest Plan. Gen. Tech. Rep. PNW-GTR-440. Portland, OR: U.S. Department of Agriculture, Forest Service, Pacific Northwest Research Station. 43 p.
- Madej, M.A.; Ozaki, V. 1996.** Channel response to sediment wave propagation and movement, Redwood Creek, California, USA. *Earth Surface Processes and Landforms*. 21: 911-927.
- Madsen, S.; Evans, D.; Hamer, T. [et al.]. 1999.** Marbled murrelet effectiveness monitoring plan for the Northwest Forest Plan. Gen. Tech. Rep. PNW-GTR-439. Portland, OR: U.S. Department of Agriculture, Forest Service, Pacific Northwest Research Station. 51 p.
- Maindonald, J.; Braun, J. 2003.** Data analysis and graphics using R an example based approach. New York: Cambridge University Press: 84-91.
- Merritt, R.W.; Cummins, K.W.; eds. 1996.** An introduction to the aquatic insects of North America. Dubuque, IA: Kendall/Hunt Publishing Company. 862p.
- Miller, D.J. 2003.** Programs for DEM analysis. In: Landscape dynamics and forest management. Gen. Tech. Rep. RMRS-GTR-101CD. Fort Collins, Co.: USDA Forest Service, Rocky Mountain Research Station.
- Miller, D. 2006.** Landslides in watershed condition assessments. Portland, OR: Unpublished report. On file with: AREMP/RPM, USDA Forest Service R6, 333 SW First Ave, Portland OR 97204. 17 pp.
- Miller, D.J.; Burnett, K.M. 2008.** Probabilistic model of debris-flow delivery to stream channels, demonstrated for the Coast Range of Oregon, USA. *Geomorphology*. 94: 184–205.
- Miller, D.J.; Burnett, K.M. 2007.** Effects of forest cover, topography, and sampling extent on the measured density of shallow, translational landslides. *Water Resources Research* .43, W03433. DOI: 10.1029/2005WR004807.

Moeur, M.; Ohmann, J.L.; Kennedy, R.; [et al.]. [In press]. Northwest Forest Plan—The First 15 Years (1994-2007): status and trend of late-successional and old-growth forests. Gen. Tech. Rep. Portland, OR: U.S. Department of Agriculture, Forest Service, Pacific Northwest Research Station.

Moeur, M.; Spies, T.A.; Hemstrom, M. [et al.]. 2005. Northwest Forest Plan—the first 10 years (1994–2003): status and trend of late-successional and old-growth forest. Gen. Tech. Rep. PNW-GTR-646. Portland, OR: U.S. Department of Agriculture, Forest Service, Pacific Northwest Research Station. 142 p.

Montgomery, D.R.; Schmidt, K.M.; Greenberg, H.M.; Dietrich, W.E. 2000. Forest clearing and regional landsliding. *Geology*. 28: 311-314.

Moore, K.; Jones, K.; Dambacher, J. 1999. Methods for stream habitat surveys. Corvallis, OR: Aquatic Habitat Inventory Project, Natural Production Program, Oregon Department of Fish and Wildlife. 59 p.

Moyer, C.D.; Chambers, J.L. [N.d.]. Results from the field data quality assessment program for the Aquatic and Riparian Effectiveness Monitoring Program 2001–2003. Draft on file with: Resource Planning and Monitoring, USDA Forest Service, Pacific Northwest Regional Office, 333 SW First Avenue, Portland, OR 97208.

Moyer, C.; Lanigan, S.; Isley, M. [et al.]. 2010. Results of the Aquatic and Riparian Effectiveness Monitoring Program Quality Assurance Program 2001-2008. Draft on file with: Resource Planning and Monitoring, USDA Forest Service, Pacific Northwest Regional Office, 333 SW First Avenue, Portland, OR 97208.

Mulder, B.; Alegria, J.; Czaplewski, R. [et al.]. 1995. Effectiveness monitoring: an interagency program for the Northwest Forest Plan. Report of the Monitoring Design Group, Intergovernmental Advisory Committee. On file with: Resource Planning and Monitoring, USDA Forest Service, Pacific Northwest Regional Office, 333 SW First Avenue, Portland, OR 97208.

THE FINDINGS AND CONCLUSIONS IN THIS REPORT ARE IN PRESS AND SUBJECT TO CHANGE PRIOR TO FORMAL DISSEMINATION BY THE AGENCIES AND SHOULD NOT BE CONSTRUED TO REPRESENT AGENCY DETERMINATION OR POLICY.

- Mulder, B.; Noon, B.; Spies, T. [et al]. 1999.** The strategy and design of the effectiveness monitoring program for the Northwest Forest Plan. Gen. Tech. Rep. PNW-GTR-437. Portland, OR: U.S. Department of Agriculture, Forest Service, Pacific Northwest Research Station. 138 p.
- Mulvey, M.; Leferink, R.; Borisenko, A. 2009.** Willamette basin rivers and streams assessment. Hillsboro, OR: Oregon Dept. Environmental Quality, Laboratory and Environmental Assessment Division. 68.
- Naiman, R.J.; Beechie, T.J.; Benda, L.E. [et al.]. 1992.** Fundamental elements of healthy watersheds in the Pacific Northwest coastal ecoregion. In: Naiman, R.J., ed. Watershed management: balancing sustainability and environmental change. New York: Springer-Verlag: 127–188.
- Noon, B.R.; Raphael, M.G.; Spies, T.A. 1999.** Conceptual basis for designing an effectiveness monitoring program. In: Mulder, B.S.; Noon, B.R.; Olsen, A.R. [et al.]. The strategy and designing of the effectiveness monitoring program for the Northwest Forest Plan. Gen. Tech. Rep. PNW-GTR-437. Portland, OR: U.S. Department of Agriculture, Forest Service, Pacific Northwest Research Station: 17–36.
- Ohmann, J.L.; Gregory, M.J. 2002.** Predictive mapping of forest composition and structure with direct gradient analysis and nearest-neighbor imputation in coastal Oregon, USA. Canadian Journal of Forest Research. 32: 725-741
- Ohmann, J.L.; Gregory, M.J., Spies, T.A. 2007.** Influence of environment, disturbance, and ownership on forest vegetation of Coastal Oregon. Ecological Applications. 17(1): 18-33.
- Olson, D.H.; Anderson, P.D.; Frissell, C.A.; Welsh, H.H.; Bradford, D.F. 2007.** Biodiversity management approaches for stream-riparian areas: perspectives for Pacific Northwest headwater forests, microclimates, and amphibians. Forest Ecology and Management. 246: 81-107.

Oregon Department of Environmental Quality. 2004. Oregon's 2004 water quality assessment section 305(b) report. Hillsboro, OR: Water Quality Division. 55.

Oreskes, N., Shrader-Frechette, K., & Belitz, K. 1994. Verification, validation, and the confirmation of numerical models in the earth sciences. *Science*, 263, 641-646.

Pabst, R.J.; Spies, T.A. 1999. Structure and composition of unmanaged riparian forests in the coastal mountains of Oregon, U.S.A. *Canadian Journal of Forest Resources*. 29: 1557–1573.

Peck, D.V.; Lazorchak, J.M.; Klemm, D.J., eds. 1999. Environmental monitoring and assessment program—surface waters: western pilot study field operations manual for wadeable streams. Unpublished draft. Washington, DC: U.S. Environmental Protection Agency. 226 p. On file with: <http://www.epa.gov/>.

Pierce K.B., Jr.; Ohmann, J.L.; Wimberly, M.C.; [et al.]. 2009. Mapping wildland fuels and forest structure for land management: a comparison of nearest-neighbor imputation and other methods. *Canadian Journal of Forest Research*. 39(10): 1901-1916.

R Development Core Team 2011. R: A Language and Environment for Statistical Computing. Vienna, Austria: R Foundation for Statistical Computing. <http://www.R-project.org>

Reeves, G.H.; Benda, L.E.; Burnett, K.M.; [et al.]. 1995. A disturbance based ecosystem approach to maintaining and restoring freshwater habitats of evolutionarily significant units of anadromous salmonids in the Pacific Northwest. In *Evolution and the aquatic ecosystem: defining unique units in population conservation*. Am. Fisheries Soc., Bethesda, MD: Nielson J.L.; Powers D.A.; eds. American Fisheries Society: 334-349.

Reeves, G.H.; Duncan, S. 2009. Ecological history vs. social expectations: managing aquatic ecosystems. *Ecology and Society*. 14(2): Article 8.

- Reeves, G.H.; Hohler, D.B.; Larsen, D.P. [et al.]. 2004.** Effectiveness monitoring for the aquatic and riparian component of the Northwest Forest Plan: conceptual framework and options. Gen. Tech. Rep. PNW-GTR-577. Portland, OR: U.S. Department of Agriculture, Forest Service, Pacific Northwest Research Station. 71 p.
- Refsgaard, J.C.; van der Sluijs, J.P.; Højberg, A.L.; Vanrolleghem, P.A. 2007.** Uncertainty in the environmental modelling process - a framework and guidance. *Environmental Modelling and Software*. 22(11): 1543-1556.
- Reid, L.M. 1993.** Research and cumulative watershed effects. Gen. Tech. Rep. PSW-GTR-141. Albany, CA: Pacific Southwest Research Station, Forest Service, U.S. Department of Agriculture; 118 p.
- Reid, L.M.; Dunne T. 1984.** Sediment production from forest road surfaces, *Water Resources Research*., 20(11): 1753–1761.
- Reid, L.M.; Furniss, M.J. 1998.** On the use of regional channel-based indicators for monitoring. Unpublished report. USDA Forest Service, Redwood Sciences Laboratory, Arcata, CA. 24 p.
- Rieman, B.E.; Hessburg, P.F.; Lee, D.C.; Thurow, R.F.; Sedell, J.R. 2000.** Toward an integrated classification of ecosystems: defining opportunities for managing fish and forest health. *Environmental Management*. 25(4): 425-444.
- Rieman, B.E.; Hessburg, P.F.; Luce, C.; Dare, M.R. 2010.** Wildfire and Management of Forests and Native Fishes: Conflict or Opportunity for Convergent Solutions? *BioScience*. 60(6): 460-468.
- Rieman, B.E.; Peterson, J.T.; Clayton, J.; Howell, P.; Thurow, R.; Thompson, W.; Lee, D. 2001.** Evaluation of potential effects of federal land management alternatives on trends of salmonids and their habitats in the interior Columbia River basin. *Forest Ecology and Management*. 153(1-3): 43-62.

- Roper, B.B.; Buffington, J.M.; Bennett; [et al.]. 2010.** A comparison of the performance and compatibility of protocols used by seven monitoring groups to measure stream habitat in the Pacific Northwest. *North American Journal of Fisheries Management*. 30(2): 565-587.
- Sedell, J.R.; Lee, D.C.; Rieman, B.E.; Thurow, R.F.; Williams, J.E. 1997.** Effects of proposed alternatives on aquatic habitats and native fishes. In: Quigley, T.M.; Lee, K.M.; Arbelbide, S.J., eds. Evaluation of EIS alternatives by the science integration team. Portland, OR: USDA Forest Service Pacific Northwest Research Station: 435-535.
- Sheskin, D.J. 2004.** Handbook of Parametric and Nonparametric Statistical Procedures, 3rd edition. Chapman & Hall/CRC. 706 p.
- Standford, J.A. 1998.** Rivers in the landscape: introduction to the special issue on riparian and groundwater ecology. *Freshwater Biology*. 40: 402-406.
- Stevens, D.L., Jr.; Olsen, A.R. 2003.** Variance estimation for spatially balanced samples of environmental resources. *Environmetrics*. 14: 593-610.
- Stevens, D.L., Jr.; Olsen, A.R. 2004.** Spatially-balanced sampling of natural resources. *Journal of the American Statistical Association*. 99: 262-278.
- Stewart, C; Martine, K.; tech eds. 2006.** Northwest Forest Plan—the first ten years (1994–2003): effectiveness of the federal-tribal relationship. Tech. Paper R6-RPM-TP-02-2006. Portland, OR: U.S. Department of Agriculture, Forest Service, Pacific Northwest Region. 51 p.
- Trimble, S.W. 1983.** A sediment budget for Coon Creek basin in the driftless area, Wisconsin, 1853-1977. *American Journal of Science*. 283: 454-474.

Tuchmann, E.T.; Connaughton, K.P.; Freedman, L.E.; Moriwaki, C.B. 1996. The Northwest Forest Plan: a report to the President and Congress. Portland, OR: U.S. Department of Agriculture, Forest Service, Pacific Northwest Research Station. 253 p.

Turban, E., & Aronson, J.E. 2001. Decision support systems and intelligent systems. Upper Saddle River, NJ: Prentice Hall.

U.S. Department of Agriculture, Forest Service [USDA FS]. [2010]. Watershed program assessment (PART) – implementation guide for assessing and tracking changes to watershed condition (Implementation Guide). Draft on file with: Resource Planning and Monitoring, USDA Forest Service, Pacific Northwest Regional Office, 333 SW First Avenue, Portland, OR 97208.

U.S. Department of Agriculture, Forest Service [USDA FS]. 1998. Stream condition inventory guidebook. Version 4.0. [Place of publication unknown]: Pacific Southwest Region. 57 p.

U.S. Department of Agriculture, Forest Service [USDA FS]. 2002. Landscape dynamics and forest management. Gen. Tech. Rep. RMRS-GTR-101-CD. [1 CD-ROM]. Fort Collins, CO: U.S. Department of Agriculture, Forest Service, Rocky Mountain Research Station.

U.S. Department of Agriculture, Forest Service [USDA FS]. 2010. Watershed Program Assessment (PART) –Implementation guide for assessing and tracking changes to watershed condition. Unpublished document. On file with: USDA Forest Service, Washington Office, 1400 Independence Ave., SW, Washington, D.C. 20250.

U.S. Department of Agriculture, Forest Service [USDA FS]. 2010. FY 2009 Legacy roads and trails, accomplishment report. Portland, OR: Pacific Northwest Region. 20 p.

U.S. Department of Agriculture, Forest Service; U.S. Department of the Interior, Bureau of Land Management [USDA and USDI]. 1994. Record of decision for amendments to Forest Service and

THE FINDINGS AND CONCLUSIONS IN THIS REPORT ARE IN PRESS AND SUBJECT TO CHANGE PRIOR TO FORMAL DISSEMINATION BY THE AGENCIES AND SHOULD NOT BE CONSTRUED TO REPRESENT AGENCY DETERMINATION OR POLICY.

Bureau of Land Management planning documents in the range of the northern spotted owl and standards and guidelines for management of habitat for late-successional and old-growth forest related species in the range of the northern spotted owl. [Place of publication unknown]. 74 p. [plus Attachment A: standards and guidelines].

van Wagtendonk, J.W.; Root, R.R.; Key, C.H. 2004. Comparison of AVIRIS and Landsat ETM+ detection capabilities for burn severity. *Remote Sensing of Environment*. 92: 397-408.

Whittier, T.R.; Hughes, R.M.; Lomnicky, G.A.; Peck D.V. 2007. Fish and amphibian tolerance values and an assemblage tolerance index for streams and rivers in the western USA. *Transactions of the American Fisheries Society*. 136: 254–271. ~~

Appendix 1: Objectives of the Aquatic Conservation Strategy¹

Forest Service and Bureau of Land Management-administered lands within the range of the northern spotted owl will be managed to:

- Maintain and restore the distribution, diversity, and complexity of watershed and landscape-scale features to ensure protection of the aquatic systems to which species, populations, and communities are uniquely adapted.
- Maintain and restore spatial and temporal connectivity within and between watersheds. Lateral, longitudinal, and drainage network connections include flood plains, wetlands, upslope areas, headwater tributaries, and intact refugia. These network connections must provide chemically and physically unobstructed routes to areas critical to fulfilling life history requirements of aquatic and riparian-dependent species.
- Maintain and restore the physical integrity of the aquatic system, including shorelines, banks, and bottom configurations.
- Maintain and restore water quality necessary to support healthy riparian, aquatic, and wetland ecosystems. Water quality must remain within the range that maintains the biological, physical, and chemical integrity of the system and benefits survival, growth, reproduction, and migration of individuals composing aquatic and riparian communities.
- Maintain and restore the sediment regime under which aquatic ecosystems evolved. Elements of the sediment regime include the timing, volume, rate, and character of sediment input, storage, and transport.
- Maintain and restore in-stream flows sufficient to create and sustain riparian, aquatic, and wetland habitats and to retain patterns of sediment, nutrient, and wood routing. The timing, magnitude, duration, and spatial distribution of peak, high, and low flows must be protected.
- Maintain and restore the timing, variability, and duration of flood-plain inundation and water table elevation in meadows and wetlands.

THE FINDINGS AND CONCLUSIONS IN THIS REPORT ARE IN PRESS AND SUBJECT TO CHANGE PRIOR TO FORMAL DISSEMINATION BY THE AGENCIES AND SHOULD NOT BE CONSTRUED TO REPRESENT AGENCY DETERMINATION OR POLICY.

- Maintain and restore the species composition and structural diversity of plant communities in riparian areas and wetlands to provide adequate summer and winter thermal regulation, nutrient filtering, appropriate rates of surface erosion, bank erosion, and channel migration and to supply amounts and distributions of coarse woody debris sufficient to sustain physical complexity and stability.
- Maintain and restore habitat to support well-distributed populations of native plant, invertebrate, and vertebrate riparian-dependent species.

¹From page B-11 of USDA and USDI 1994.

DRAFT

Appendix 2: Project-scale restoration efforts.

Seven local unit watershed restoration projects are highlighted from throughout the Northwest Forest Plan area. Some of the projects may have had immediate effects, such as opening up habitat to fish by replacing poorly designed culverts that previously blocked fish passage. But most restoration projects should be viewed as a critical first step in restoring natural watershed processes.

Sidebar 10—Cummins and Tenmile Watershed

Sidebar 11—Elkhorn Creek Watershed

Sidebar 12—Legacy roads and Trails Program

Sidebar 13—Methow Valley Subbasin

Sidebar 14—Paradise Creek

Sidebar 15—Scott River

Sidebar 16—Six Rivers National Forest

Appendix 3: Watersheds Selected for Inchannel Monitoring

The 250 watersheds shown in Table 8 were randomly chosen and sampled as part of a panel design to determine the status and trend of inchannel processes. Approximately 25 to 30 watersheds have been sampled each year.

Table 8—List of the 250 watersheds randomly chosen selected to determine the status and trend of inchannel processes. Includes the U.S. Geological Survey hydrologic unit code (USGS HUC), the aquatic province, the national forest (NF), Bureau of Land Management (BLM) unit, or national park (NP) that manages the land, the fifth-field watershed name, the sixth-field watershed name, and the percentage of the total watershed area that is federally owned.

Appendix 4: Model Structures, Evaluation Criteria, and Status and Trend Maps for each of the Provincial Decision-Support Models

The model structure, evaluation criteria, and watershed condition status and trend maps are shown for each aquatic province with the Northwest Forest Plan area.

Model Structures

The AREMP team worked with local experts from each province to develop evaluation models for the upslope/riparian and the inchannel data. These models hierarchically aggregate a number of attributes into broader indices of inchannel and watershed condition. A graphical depiction of the model structures for each province is contained in this appendix (Figure 48 through Figure 53). Some model sections were “turned off” in this iteration because the corresponding data were not available. These unused portions of the models are indicated by gray text.

Attribute-level Scoring

A model begins by reading in a set of data observations, which we call “attributes,” for a watershed. These attributes are the rightmost nodes in the model structure diagrams. For example, attributes for the High Cascades province (Figure 48) watershed condition model includes “watershed road density”. Details about the attributes for each provincial model are shown in Table 9 (watershed condition) and Table 10 (inchannel condition). The “Attribute and measure” column contains the attribute name, units of measure, and qualifiers if any, (e.g., temperature is evaluated differently in watersheds depending on whether bull trout, *Salvelinus confluentus*, are present).

As part of the model-building process, the provincial experts developed evaluation criteria for each attribute. These evaluation criteria determine how any particular data value is scored on a common scale from +1 to -1 according to its relationship to watershed condition. As the attribute data for each watershed

THE FINDINGS AND CONCLUSIONS IN THIS REPORT ARE IN PRESS AND SUBJECT TO CHANGE PRIOR TO FORMAL DISSEMINATION BY THE AGENCIES AND SHOULD NOT BE CONSTRUED TO REPRESENT AGENCY DETERMINATION OR POLICY.

are read into the model, they are each compared to their respective evaluation criteria to produce an evaluation score for each between +1 and -1. For the Olympic Peninsula province, if there are no riparian roads (density = 0), then the evaluated score would be +1; if road density was 0.1 mi/mi² of riparian area or greater, the score would be -1; and if the density falls between 0 and 0.1 mi/mi², the attribute receives a score that is a linear interpolation between +1 and -1 (e.g., 0.05 mi/mi² would evaluate to 0). Note that there is an important difference between a data value of “zero” and “no data.” Data values of zero (e.g., riparian roads example above) are compared to their evaluation curve the same as all other data values. However, if data for a particular attribute are lacking in a particular watershed, then that attribute is given an *evaluated score* of zero, representing a neutral value that does not indicate either good or poor condition. The “Data value” and “Evaluated score” columns show how the raw data values correspond to evaluated scores, and the “Curve shape” column gives a graphical depiction of this (generally linear) function, with data values represented on the x-axis and corresponding evaluation scores on the y-axis. The “Source” column gives the basis on which the curve was constructed, most often the professional judgment of workshop participants but also including data sets and published reports or standards.

Aggregation of Attribute Scores

After each attribute datum is evaluated, the model begins to aggregate these scores together in a hierarchical fashion. The combined score is passed up to the next level in the model hierarchy, where it is combined again with results from other parts of the model. The modeling software enables a number of different aggregation functions, but we limited choice to the three simplest:

- MIN: take the minimum score from those being aggregated.
- AVE: take the average of the aggregated scores.
- MAX: take the maximum score from those being aggregated.

These functions determine whether the situation is of a “limiting factor” type, where the worst condition score determines the combined score, a “partially compensatory” situation, where scores are all counted equally, or a “fully compensatory” situation, where the best score determines the combined score.

In addition to operators, each node in the model can also be assigned a weight. These weights are listed on the model structure diagrams. For example, the North Cascades model (Figure 50) weighted riparian tree size at 0.7 and watershed-wide vegetation at 0.3, so the overall vegetation score comes 70 percent from riparian value and 30 percent from the watershed value. These weights are only relevant under the AVE operator.

Evaluation Model Structures for each Aquatic Province

Figure 48. High Cascades province evaluation model structures for watershed and inchannel conditions.

Figure 49. Klamath-Siskiyou -Franciscan province evaluation model structures for watershed and inchannel conditions.

Figure 50. North Cascades province evaluation model structures for watershed and inchannel conditions.

Figure 51. Olympic Peninsula province evaluation model structures for watershed and inchannel conditions.

Figure 52. Oregon Coast province evaluation model structures for watershed and inchannel conditions.

Figure 53. West Cascades province evaluation model structures for watershed and inchannel conditions.

Decision-Support Model Attributes and Evaluation Criteria by Province

Table 9—Evaluation criteria used in watershed-level decision-support model for aquatic province models.

Table 10—Evaluation criteria used in inchannel-level decision-support models by aquatic province.

Status and Trend Maps for each Province

Figure 54—Distribution of inchannel condition scores for the randomly selected watersheds in the High Cascade and West Cascade Provinces in the Northwest Forest Plan (NWFP) area that have been sampled for inchannel attributes as of 2009.

Figure 55—Watershed condition status scores (2008) for the High Cascade and West Cascade Provinces in the Northwest Forest Plan (NWFP) area.

Figure 56—Watershed condition trend scores (1994 to 2008) for the High Cascade and West Cascade Provinces in the Northwest Forest Plan (NWFP) area.

Figure 57—Distribution of inchannel condition scores for the randomly selected watersheds in the Klamath/Siskiyou and Franciscan Provinces in the Northwest Forest Plan (NWFP) area that have been sampled for inchannel attributes as of 2009.

Figure 58—Watershed condition status scores (2008) for the Klamath/Siskiyou and Franciscan Provinces in the Northwest Forest Plan (NWFP) area.

Figure 59—Watershed condition trend scores (1994 to 2008) for the Klamath/Siskiyou and Franciscan Provinces in the Northwest Forest Plan (NWFP) area.

Figure 60—Distribution of inchannel condition scores for the randomly selected watersheds in the North Cascades Province in the Northwest Forest Plan (NWFP) area that have been sampled for inchannel attributes as of 2009.

Figure 61—Watershed condition status scores (2008) for the North Cascades Province in the Northwest Forest Plan (NWFP) area.

Figure 62—Watershed condition trend scores (1994 to 2008) for the North Cascades Province in the Northwest Forest Plan (NWFP) area.

Figure 63—Distribution of inchannel condition scores for the randomly selected watersheds in the Olympic Peninsula Province in the Northwest Forest Plan (NWFP) area that have been sampled for inchannel attributes as of 2009.

Figure 64—Watershed condition status scores (2008) for the Olympic Peninsula Province in the Northwest Forest Plan (NWFP) area.

Figure 65—Watershed condition trend scores (1994 to 2008) for the Olympic Peninsula Province in the Northwest Forest Plan (NWFP) area.

Figure 66—Distribution of inchannel condition scores for the randomly selected watersheds in the Oregon Coast Province in the Northwest Forest Plan (NWFP) area that have been sampled for inchannel attributes as of 2009.

Figure 67—Watershed condition status scores (2008) for the Oregon Coast Province in the Northwest Forest Plan (NWFP) area.

Figure 68—Watershed condition trend scores (1994 to 2008) for the Oregon Coast Province in the Northwest Forest Plan (NWFP) area.

Appendix 5: Additional Information on Inchannel, Upslope and Riparian Data Inputs

Inchannel Data

Study Design

Watershed Selection

We excluded watersheds from sampling if any of the following conditions were met:

1. A minimum of sampling four stream reaches could not be completed within 6 days (the length of time available each sampling trip) because of time constraints, accessibility issues, or stream reach constraints (see below).
2. Watersheds that didn't have at least four sites that met the site selection criteria.

Watersheds that were not assessed but remain in the sample include watersheds that were originally evaluated and dropped based on selection criteria that did not include the possible selection of dry channels and those that experienced a temporal disturbance preventing the crew from sampling during the correct panel. Watersheds originally dropped for not having enough water were reevaluated based on photos and reconnaissance to determine whether they would likely meet the current site selection protocol or be permanently dropped. Disturbances that prevented field crews from accessing watersheds include:

3. Watersheds deemed dangerous for a survey crew to be working in the area (i.e., law enforcement personnel identified a watershed as currently having prevalent drug growing operations).
4. Fire activity that blocked or limited road/trail access to the watershed or had potential to spread, endangering the crew while working in the stream.

Skipped watersheds have not been excluded from the sample. They are expected to be sampled in subsequent years, since excluding them would violate the assumption of random selection.

Inchannel Site Selection

Given the list of randomly selected sites, crews sampled individual sites until an 8-day sampling period expired. In each watershed, sites were sampled in the order they were selected. The number of sites sampled was typically a function of access (i.e., more sites were sampled in areas that were easily accessed). We excluded individual stream reaches from sampling if:

5. The stream reach was not safely accessible; i.e., it could not be reached without putting the crew in danger. (A long hike into a steep canyon did not qualify as a dangerous situation for the crew.)
6. The stream reach was not wadeable because of depth or current.
7. Travel time (round trip) from road camp or wilderness camp was over 4 hours to get to and from the stream reach. The wilderness camp cannot be more than 1 day hike from the trailhead.
8. The geographic positioning system (GPS) point (used to identify the beginning of a stream reach) was located on private land.
9. The GPS point for a stream reach was located in a lake, wetland or marsh, or on a dam or glacier.
10. The stream reach was an artificial stream or irrigation canal.

Upslope/Riparian Data

Common Geographic Information System (GIS) Layers Used for All Provinces

Watershed boundaries

The boundaries used were from the first draft of the sixth-field hydrologic unit code boundaries developed by the Regional Ecosystem Office dated 2002. The first draft (version 1.1) was used because it was the version available when the 250 watersheds were selected.

Streams and lakes

The Oregon and Washington streams and water bodies were taken from the BLM 1:24,000 stream geodatabase. The BLM created regional geodatabases by joining the Pacific Northwest hydrography

framework (HUC) watershed data, so the line work is identical to the northwest hydrography framework layer. This stream layer has highly variable densification, but most of our metrics are calculated on a per stream mile basis and so are relatively insensitive to stream density (e.g. more road crossings would be mapped on a denser stream layer but the per stream mile metric would not necessarily be higher). The California layer was pieced together by the FS Region 5 remote sensing laboratory from a combination of FS, BLM, and cartographic feature file data. The resulting layer also had varying stream density depending on the source. Some attributes were applied only to perennial streams, which were identified as PERIODICITY [field] = “per” (which excluded streams coded as unknown or null).

Riparian Areas / Stream buffers

Riparian areas differed by province, and different riparian buffer widths were sometimes used for road-stream interactions and vegetation. Buffer widths are expressed here as one-sided distance from the stream, so a 50 m buffer creates a riparian area that is 100 m wide. Riparian areas were created by buffering the stream lines by the prescribed width. Streams wide enough to be represented by a double line were buffered separately and the resulting buffer added to the single-line stream buffers. Lakes were removed from the final buffered layer.

Digital elevation models (DEM)

30-m (98.4-ft) DEMs were obtained in 2001 from the National Elevation Data set (NED) compiled by the U.S. Geological Survey (USGS).

Average Precipitation

Average annual precipitation from 1961 to 1990 downloaded from the Oregon State University PRISM Climate Group (<http://www.prism.oregonstate.edu/products/>).

Roads

Road layers used in Oregon and Washington were a combination of USDA Forest Service (FS) road layers with the USDI Bureau of Land Management (BLM) ground transportation layer (GTRN). The FS

layers were obtained from each of the national forests in the NWFP area. The FS layers included attributes from the Infrastructure (INFRA) application attached to the road segments. The road layers were clipped to the ownership boundary of the forest. The FS ownership areas were removed from the BLM layer, and the FS road layers were pasted in using a “cookie cutter” process. No edge matching was done. The BLM and FS layers are from 2008. The decommissioning and year-built attributes were used to determine the change in the road system since 1994. The FS Pacific Southwest Region (Region 5) remote sensing laboratory constructed the California road layer.

Vegetation

Detailed attributes of forest composition and structure were mapped for all forests in the NWFP area for two “bookend” dates using gradient nearest neighbor (GNN) imputation (Ohmann and Gregory 2002). The bookend dates were 1996 and 2006 in Washington and Oregon, and 1994 and 2007 in California. The vegetation mapping for NWFP monitoring marks the first application of GNN imputation to multiple imagery dates. The primary challenge was to develop GNN model-based maps for the two bookend dates that were temporally consistent with each other and with maps of forest disturbance. A full description of the GNN bookends methodology can be found in Moeur et al. (in press).

Gradient nearest neighbor is one of many variations of nearest neighbor imputation methods. The GNN method was developed in the Pacific Northwest, and has been applied to broad-scale vegetation mapping across a wide range of forest ecosystems (Ohmann et al. 2007, Pierce et al. 2009). In GNN, forest attributes from regional inventory plots are assigned to map pixels where data are missing on the basis of a modeled relationship between the detailed forest attributes from plots and a combination of spatial predictor variables derived from Landsat satellite imagery, climate variables, topographic variables, and soil parent materials. The assigned plot data allows generation of thematic maps for any detailed attribute (or combination of attributes) of forest composition or structure measured on the plots. Ground data for GNN models were basal area by tree species and size class from Forest Inventory and

Analysis (FIA) periodic inventories on nonfederal lands, FIA annual inventory on all ownerships, and current vegetation survey inventories on FS and BLM lands. The assumption behind GNN methods is that two locations with similar combined spatial “signatures” should also have similar forest structure and composition.

For the bookends analysis, the GNN models used Landsat imagery that had been geometrically rectified and radiometrically normalized through time using the LandTrendr algorithms (Kennedy et al. 2007). The goal was to minimize differences in forest characteristics between the GNN bookend models that were caused by differences in the imagery that did not reflect real changes on the ground.

Vegetation Disturbance

A new approach to monitoring landscape change was implemented to map forest disturbance in the NWFP. Landsat-based detection of trends in disturbance and recovery (LandTrendr) produces yearly maps of forest disturbance using new analysis of annual Landsat Thematic Mapper satellite imagery. The LandTrendr approach improves the temporal frequency of disturbance maps, better separates subtle change from background noise, and can detect new phenomena that cannot be captured by older technologies that compare only two images at a time (Kennedy et al. 2010). A full description of the LandTrendr methodology can be found in Moeur et al. (2010).

In LandTrendr, annual time series of Landsat imagery were assembled for the entire NWFP area and then processed using basic atmospheric correction, cloud screening, and radiometric normalization. After image preparation, the time series of the Normalized Burn Ratio (NBR) (van Wagtenonk et al. 2004) for each 30-m pixel was extracted, and temporal segmentation algorithms were used to identify periods of both stability and change in each pixel’s NBR trajectory (Kennedy et al. 2007).

Maps were created by evaluating each pixel’s NBR segmentation results. Disturbance segments were identified as those experiencing declines in NBR over time, and pre- and post-disturbance percentage of vegetation cover was predicted using a statistical model of cover developed from photo interpreted plots

THE FINDINGS AND CONCLUSIONS IN THIS REPORT ARE IN PRESS AND SUBJECT TO CHANGE PRIOR TO FORMAL DISSEMINATION BY THE AGENCIES AND SHOULD NOT BE CONSTRUED TO REPRESENT AGENCY DETERMINATION OR POLICY.

(Cohen et al 2010). Relative cover loss was calculated as the change in cover divided by pre-disturbance cover. Pixels with less than 15 percent relative vegetation loss over 1 year duration were removed, and groups of remaining adjacent pixels with the same disturbance year and with size greater than about 1 ha (11 pixels) were retained as disturbance patches. Each pixel remaining in a patch was labeled with the magnitude of change (relative cover), duration of the loss process (in years), year of disturbance onset, and likely cause of the disturbance (fire, insect, or harvest). Allowing for slight mismatch in timing of segmentation (typically +/- 1 year for harvest and fire), the LandTrendr segmentation algorithms were found to capture and correctly time 90 and 86 percent of the high-intensity harvest and fire events, respectively. Medium-intensity fire and harvest were also captured with high accuracy (about 75 percent), and only lost significant sensitivity at low intensity level.

Areas Excluded from Vegetation Analysis

Subalpine areas, along with nonforested areas, were excluded from the vegetation analysis. Nonforested areas were based on a mask defined by The Landscape Ecology, Modeling, Mapping and analysis group (<http://www.fsl.orst.edu/lemma>) for the IMA data in Oregon and Washington, and a nonforested mask developed by the Northern Spotted owl and Marbled Murrelet NWFP habitat groups (Davis et al., in press). Subalpine areas were defined as from zones 29 to 33 in the 2009 version of the modeled potential natural vegetation zones for Washington and Oregon provided by the USDA Forest Service (Jan A. Henderson, USDA Forest Service, 2009). For California, the subalpine layer developed by the NWFP northern spotted owl group was used (Davis et al., in press).

Landslide Risk

Landslide risk was estimated using a model developed by the Earth Systems Institute (Miller 2003). The model was calibrated for the NWFP area using landslide data derived from aerial photographs in 14 AREMP sixth-field watersheds (Miller 2006). The model calculated a topographic risk factor, based on characteristics such as slope and concavity, from a 10-meter digital elevation model. This baseline

susceptibility was multiplied by the calibrated vegetation and road risk factors. The vegetation multiplier was based on the quadratic mean diameter of all dominant and codominant trees (QMDA_DOM) in the plot assigned to the map pixel: greater than 4" = multiply by 0.5, less than 4" = multiply by 1.48. A road proximity multiplier of 2.73 was applied to all pixels within 50 meters of a road.

Additional Details on Provincial Models

This section provides additional attribute details not covered by the generic descriptions above and the attribute tables in appendix 4.

High Cascades Province

- Road crossings in high-landslide-hazard areas: High hazard areas were defined by topographic landslide risk scores greater than 0.28 from the landslide model. The value of 0.28 was determined using an existing map of hazard roads from the Olympic Province. A 20 m buffer was applied to the Olympic hazard roads, and this was overlaid on the Miller (2006) topographic landslide risk layer. Eighty percent of these hazard roads buffers were captured by a topographic risk threshold of 0.28. This risk factor was applied to the High and West Cascades Province after examining their distributions of risk values were similar.
- Vegetation–riparian: Map pixels were classified as either lodgepole pine or nonlodgepole, based on dominant species. A lower diameter threshold was applied to lodgepole stands (8 inches) than to nonlodgepole stands (14 inches).
- Vegetation–watershed: Different canopy cover thresholds were applied to the wetter (70%) and drier (40%) areas of the province, as defined by the 40 inch precipitation contour from the PRISM GIS layer. Watersheds in the rain on snow hydrologic zones are considered to be more sensitive to increases in peak flow from harvest and road building (Grant et al. 2008), so different evaluation criteria were applied to these zones. Zones were defined using a combination of the rain and snow bands from the WOPR (BLM Western Oregon Plan revisions) in Oregon and elevation classes in Washington. The elevation classes are less than 457 m for rain, 457 to 1219 m for rain on snow, and greater than 1219 m for snow on snow.

Klamath/Siskiyou and Franciscan Province

- Road density on the lowest one-third of slopes: The lowest one-third of the slope was determined by running a slope position Arc Macro Language (AML) script that uses DEMs and creates a grid with the slope numbered from 1 to 100, with 1 being the bottom and 100 the top. The numbers 1 to 33 were used to define the lower one-third of the slope. The lower one-third area was turned into a polygon layer and intersected with the roads.
- Off- road vehicle (OHV) trail density. The officially designated OHV trails were extracted from the FS INFRA layer and the BLM GTRN layer and miles summed by watershed.
- Vegetation–riparian & watershed–oak woodlands: Map pixels were classified as either oak or non-oak, based on dominant species. Lower canopy cover evaluation criteria were applied to oak woodlands.

North Cascades Province

- Road density in hazard areas: Hazard areas were defined as having a topographic landslide risk value of greater than 1.02. The value of 1.02 was derived by comparing the current landslide model results with hazard areas determined in the AREMP 2004 report using the Shalstab model and land type associations (Gallo et al. 2005).
- Vegetation–east/west: Different vegetation metrics and evaluation criteria were applied on the east and west sides of the province, based on the Cascade crest (incorporating the Pickett crest).
- Vegetation–watershed–east–dry/wet areas: Different metrics were used for the wet and dry areas on the eastern side of the Cascades. Map pixels were assigned to dry areas using the following plant association groups: All ponderosa pine and Douglas-fir plant association groups or zones, Grand fir-pinegrass, Grand fir-pinegrass-lupine, Grand fir-pinemat manzanita, Grand fir-oceanspray-pinegrass, Grand fir-mountain snowberry, Grand fir-spirea-bracken fern, Grand fir-snowberry-pinegrass, Shrub-steppe.

Olympic Peninsula

- Vegetation–riparian: Different metrics were applied to the wetter (southwest) and drier (northeast) zones of the province as defined by the 110 in precipitation contour in the PRISM GIS layer. In the wetter zone, the metric was percentage of area where tree quadratic mean diameter ≥ 20 inches or

stand age was ≥ 100 years old. In the drier zone, the metric was percentage of area where tree quadratic mean diameter ≥ 12 inches or stand age was ≥ 100 years old. Riparian area was defined as 160-ft buffer around streams for both zones.

- Vegetation–urban/agriculture: The percentage of watershed covered by urban and agricultural land was calculated from the designations in the base vegetation dataset (Moeur et al., in press).

Oregon Coast

- Vegetation–urban/agriculture: The percentage of watershed covered by urban and agricultural land was calculated from the designations in the base vegetation dataset (Moeur et al., in press)

West Cascades

- Road crossings in high-landslide-hazard areas: High hazard areas were defined by topographic landslide risk scores greater than 0.28 from the landslide model. The value of 0.28 was determined using an existing map of hazard roads from the Olympic Province. A 20 m buffer was applied to the Olympic hazard roads, and this was overlaid on the Miller (2006) topographic landslide risk layer. Eighty percent of these hazard roads buffers were captured by a topographic risk threshold of 0.28. This risk factor was applied to the High and West Cascades Province after examining their distributions of risk values were similar.
- Vegetation–riparian: Map pixels were classified as either lodgepole pine or nonlodgepole, based on dominant species. A lower diameter threshold was applied to lodgepole stands (8 inches) than to nonlodgepole stands (14 inches).
- Vegetation–watershed: Different canopy cover thresholds were applied to the wetter (70%) and drier (40%) areas of the province, as defined by the 40 inch precipitation contour from the PRISM GIS layer. Different evaluation criteria were applied to different hydrologic zones: rain on rain, rain on snow, and snow on snow bands.
- Rain on snow bands: A combination of the rain and snow bands from the WOPR (BLM Western Oregon Plan revisions) in Oregon and elevation classes in Washington. The elevation classes are less than 457 m. for rain, 457 to 1219 m for rain on snow, and greater than 1219 m for snow on snow. Watersheds in the rain on snow bands are considered to be more sensitive to increases in peak flow from harvest and road building (Grant et al. 2008).

Appendix 6: Quality Assessment of Upslope/Riparian and Inchannel Data Inputs

Land Use Categories

Each subwatershed was categorized according to a NWFP land use allocation and a key/nonkey designation based on the categories covering the largest share of its land area. Table 11 provides summary statistics on the percentage of watershed area actually falling within its assigned categories.

Table 11—Accuracy of land use category assignments to individual subwatersheds.

Upslope and Riparian Attributes

Roads

The 10-year AREMP report (Gallo et al. 2005) assessed the accuracy of the roads GIS layer by comparing it to roads digitized from digital aerial photography data for 38 watersheds. Since the creation of a roads layer from aerial photography is resource intensive, and we have not heard of any major changes to the corporate GIS roads layer, we have relied on a re-analysis of this past dataset for assessing road attribute accuracy for the current report. Our re-analysis reduced the scope from all lands to federal lands only and changed the road crossings metric from a per road mile to a per stream mile measure. Summary statistics for the three primary road attributes used are presented in Table 12.

Table 12--Percent difference in aerial photography-based road attributes compared to corporate GIS road attributes.

Vegetation

The accuracy of the vegetation attributes used in the AREMP watershed assessment models are reported in LEMMA (2010) in terms of correlation coefficients (R) and root mean squared error (RMSE)

by modeling region. We have summarized the range of each accuracy metric across all the regions in Table 13

Table 13—Summary of vegetation attribute accuracy assessment metric ranges.

Landslide Risk

No error assessment has been done for the landslide risk attribute.

Inchannel Attributes

Methods

AREMP implemented a Quality Assurance (QA) Plan to ensure that all data collected were scientifically sound and of known quality and includes activities such as training field crews, capturing errors in data collection procedures, and conducting secondary surveys at randomly selected sites. Analysis of the remeasured subset of sample sites (generally referred to as the paired set of initial-survey and secondary-survey data) was conducted to determine the consistency of the sample data (Table 1). These paired survey results were examined to distinguish between environmental and measurement effects. Environmental effects are the differences that occur naturally between watersheds and between sites in watersheds; these effects are considered uncontrollable. Measurement effects include differences in measurements between crews and unexplained error; these effects are considered controllable through training, refinements in field protocols (to reduce subjectivity), and improvement in equipment.

Results

Highlights of a 2010 review of 9 years of the QA program (and changes we are making in response to the review) include the following:

- Additional training time is needed to ensure crews fully understand how to implement field protocols.
 - An additional week of training was added during the 2010 field season.

THE FINDINGS AND CONCLUSIONS IN THIS REPORT ARE IN PRESS AND SUBJECT TO CHANGE PRIOR TO FORMAL DISSEMINATION BY THE AGENCIES AND SHOULD NOT BE CONSTRUED TO REPRESENT AGENCY DETERMINATION OR POLICY.

- We should continue to develop and implement automated data checking processes such as data entry forms and data validation rules using the latest technology for efficiency.
 - New applications for our data loggers to ensure accurate data entry were added during the 2010 field season.
- Annually review the training program and evaluate all aspects for possible improvement.
 - Changes in our 2010 field season included adding more laser training and more practice reaches.
- Continue surveying some watersheds a second time during the same year as the initial survey to determine the between-crew variability.
 - Eight watersheds will have reaches resampled by independent crews in 2010 to determine between-crew variability.
- The variation due to measurement by survey crews across all years was low for average bankfull width, average bankfull width:depth, gradient, the median particle size (D_{50}) without bedrock, pool tail crest fines, DSM (Decision Support Model) wood pieces and frequency, and EPT (Ephemeroptera, Plecoptera, Trichoptera) richness (Table 14). This indicates these field measurements are more consistent than the other field attributes (more accurately measured). No change is recommended for protocols associated with these attributes.
- The variation due to measurement across all years in average bankfull depth, stream channel sinuosity, median particle size (D_{50}), number of pools, pool frequency, average residual pool depth, dissolved oxygen, specific conductance, intolerant richness, and terrestrial amphibian Simpsons index is high relative to the environmental variation (Table 14). Further examination of how the protocols are being implemented is warranted.
- On an annual basis, some field attributes such as the frequency of DSM wood pieces and pool frequencies show marked improvement in performance (measurement accuracy) after 2004, which corresponds to when an overhaul of the training program occurred (i.e., new training sites, development of standardized lesson plans, and an overall training document as a component of the QA program) and the standardization of field protocols with the PACFISH/INFISH Biological Opinion Effectiveness Monitoring Program.

THE FINDINGS AND CONCLUSIONS IN THIS REPORT ARE IN PRESS AND SUBJECT TO CHANGE PRIOR TO FORMAL DISSEMINATION BY THE AGENCIES AND SHOULD NOT BE CONSTRUED TO REPRESENT AGENCY DETERMINATION OR POLICY.

- Data completeness (the proportion of sites with measurements present for all attributes) was 92, 92, and 86 percent for channel morphology, physical habitat, and water chemistry attributes, respectively.

The amount of effort invested in conducting secondary surveys of sites as part of the quality assurance program was equivalent to about 2 years worth of work spread across the 9 years examined. In-depth findings and recommendations resulting from examining all years of data collected by the program will be posted at: <http://www.reo.gov/monitoring/reports/watershed-reports-publications.shtml>.

Table 14—Field attributes evaluated in the Quality Assurance Program report. The environment column represents the percent of total variation in the data due to the differences between watersheds plus the differences between sites within watersheds. The Observer column represents the percent of the total variation in the data due to measurement by crews and unexplained error. The two columns do not total 100% because other terms related to time (differences through time) account for the remained of the total variation.

Appendix 7: Contact Information

Want to know more? Please contact:

Steve Lanigan, Team Leader	503.808.2261	slanigan@fs.fed.us
Sean Gordon, Research Associate	503.808.2698	seangordon@fs.fed.us
Peter Eldred, GIS Analyst	541.750.7078	peldred@fs.fed.us
Chris Moyer, Fisheries Biologist	541.750.7017	cmoyer@fs.fed.us
Mark Isley, Database Manager	541.750.7081	markisley@fs.fed.us
Steve Wilcox, GIS Cartographer	541.750.7122	sewilcox@fs.fed.us
Heidi Andersen, Fisheries Biologist	541.750.7067	hvandersen@fs.fed.us

Please visit our Web site for more information on publications, presentations, reports, and summer employment:

<http://www.reo.gov/monitoring/watershed-overview.shtml>

The U.S. Department of Agriculture (USDA) prohibits discrimination in all its programs and activities on the basis of race, color, national origin, age, disability, and where applicable, sex, marital status, familial status, parental status, religion, sexual orientation, genetic information, political beliefs, reprisal, or because all or part of an individual's income is derived from any public assistance program. (Not all prohibited bases apply to all programs.) Persons with disabilities who require alternative means for communication of program information (Braille, large print, audiotape, etc) should contact USDA's TARGET Center at 202.720.2600 (voice and TDD). To file a complaint of discrimination, write USDA, Director, Office of Civil Rights, 1400 Independence Avenue, Washington, D.C. 20250-9410, or call 800.795.3272 (voice) or 202.720.6382 (TDD). USDA is an equal opportunity provider and employer.

THE FINDINGS AND CONCLUSIONS IN THIS REPORT ARE IN PRESS AND SUBJECT TO CHANGE PRIOR TO FORMAL DISSEMINATION BY THE AGENCIES AND SHOULD NOT BE CONSTRUED TO REPRESENT AGENCY DETERMINATION OR POLICY.

Tables

Table 1—Collapsed land use and watershed categories used in this analysis.

Category	Number of watersheds	Land use allocation ^a or ACS designation
Congressional reserves	406	Congressional reserves Administratively withdrawn areas
Late-successional reserves	464	Late-successional reserve 1 Late-successional reserve 2 Late-successional reserve 3 Managed late-successional reserves Adaptive management – late successional reserve ^b
Matrix	509	Matrix lands Riparian reserves Adaptive management areas – non late-successional reserves ^c
Key	469	Tier 1 key watersheds Tier 2 key watersheds
Nonkey	910	All federal lands not designated as key watershed

^a Described by Tuchmann et al. (1996)

^b These areas follow the general guidelines associated with late-successional reserves

^c These areas follow the general guidelines associated with matrix lands

Table 2—Attributes included in the watershed condition assessments listed by ecological process.

Ecological processes		Attributes Assessed
General process	Key process	
Upslope subsystem:		
Vegetative succession, growth, and mortality	Wood production and transport	Vegetation seral stage
Hydrologic cycle	Peak flow and timing/amount	Vegetation hydrologic maturity and road density
Soil cycle	Sediment production and transport	Road density, landslide risk, vegetation seral stage
Riparian subsystem:		
Vegetative succession, growth, and mortality	Community structural development	Vegetation seral stage and association
Vegetative succession, growth, and mortality	Wood delivery	Vegetation seral stage, riparian road density
Soil cycle	Sediment production and transport	Road density, off-highway vehicle (OHV) trail density, stream-crossing density
Hydrologic cycle	Flood-plain loss	Riparian road density
Hydrologic cycle	Connectivity for fish and wood passage	Stream-crossing density
Inchannel subsystem:		
Hydrologic cycle	Water storage and yield, off-channel habitat	Channel connectivity with flood plain
Channel structural dynamics	Sediment and wood delivery	Bankfull width:depth ratio, channel pools, wood, substrate composition
Energy exchange	Heat delivery	Water temperature
Chemical and nutrient turnover	Chemical and nutrient delivery	Water quality
Biotic community dynamics	Biotic integrity	Amphibians and macroinvertebrate indices

Adapted from Reeves et al. (2004).

Table 3—Inchannel attributes used in assessment and sources of metrics used

Attribute	Source of metric
Bankfull width-to-depth ratio	Calculated from cross-sectional profiles (Peck et al. 1999)
Pool frequency	Calculated from longitudinal profiles (Peck et al. 1999)
Pool depth	Maximum pool depth (AREMP 2009, p. 29)
Flood-plain connectivity	AREMP protocol (AREMP 2009, p. 31)
Wood frequency	From protocol developed by the Oregon Department of Fish and Wildlife (Moore et al. 1999)
Substrate fines (percentage of fine sediments)	From protocol developed by USDA Forest Service (1998)
Substrate D50 (median particle size)	Based on a modification of Peck et al. (1999)
Macroinvertebrates	See “Macroinvertebrate Metric” side bar (p. XX)
Terrestrial and aquatic amphibians	See “Amphibian Metric” side bar (p. XX)
Dissolved oxygen	Data were collected with either with a YSI Professional or YSI 556 MPS ¹
Water temperature	Data were collected with either with a YSI Professional or YSI 556 MPS by using Onset tidbits

¹ The use of trade or firm names in this publication is for reader information and does not imply endorsement by the U.S. Department of Agriculture of any product or service.

Table 4—Inchannel attribute scores by land use category (CR = congressional reserves, LSR = late-successional reserves)

Attribute	Statistic	CR	LSR	Matrix	p-value	Statistic	Key	Nonkey	p-value
Inchannel score	median	+0.31	+0.22	+0.16	0.04	mean	+0.27	+0.15	0.04
Pools	median	+0.66	+0.82	+0.94	0.02	median	+0.82	+0.82	0.13
Substrate	median	+0.45	+0.47	+0.65	0.69	mean	+0.43	+0.39	0.65
Wood	mean	+0.10	-0.02	+0.02	0.67	median	+0.26	-0.11	0.03
Temperature	mean	+0.31	+0.05	-0.01	0.17	median	+0.32	+0.15	0.94
Invertebrates	mean	+0.38	+0.13	+0.13	0.08	mean	+0.25	+0.15	0.28

Key: bold indicates significant differences between italicized and non-italicized values.

Table 5—Percent of watersheds (n = 1379) falling within each status condition class (2008) by aquatic province

Province	Very Low	Low	Moderate	High	Very High
High Cascades		21%	36%	32%	12%
Klamath-Siskiyou-Franciscan	<1%	23%	31%	27%	18%
North Cascades	5%	11%	20%	27%	37%
Olympic	2%	8%	5%	30%	56%
Oregon Coast		8%	25%	32%	35%
West Cascades	13%	34%	27%	17%	9%

THE FINDINGS AND CONCLUSIONS IN THIS REPORT ARE IN PRESS AND SUBJECT TO CHANGE PRIOR TO FORMAL DISSEMINATION BY THE AGENCIES AND SHOULD NOT BE CONSTRUED TO REPRESENT AGENCY DETERMINATION OR POLICY.

Table 6—Percent of watershed score trend changes (1994 to 2008) due to changes in roads, vegetation, and landslide risk attributes.

Score Change Categories	Roads	Vegetation	Landslide Risk	All Watersheds (n = 1379)	
				Total Count	Total Percent
-2 to -0.3	0%	94%	6%	18	1%
-0.3 to -0.1	0%	84%	16%	52	4%
-0.1 to <0	2%	83%	15%	246	18%
0				108	8%
>0 to 0.1	24%	66%	11%	753	55%
0.1 to 0.3	42%	30%	28%	175	13%
0.3 to 2	38%	8%	54%	27	2%

Table 7—Percent of watersheds (n = 1,379) that decreased, increased, or had no change in watershed scores between 1994 and 2008.

	Decrease	No change	Increase
All watersheds	23%	8%	69%
Changes > ± 0.1	4%	86%	10%
Congressional reserve	38%	19%	43%
Late-successional reserve	17%	3%	80%
Matrix	16%	3%	81%
Key	23%	9%	68%
Nonkey	23%	7%	70%

THE FINDINGS AND CONCLUSIONS IN THIS REPORT ARE IN PRESS AND SUBJECT TO CHANGE PRIOR TO FORMAL DISSEMINATION BY THE AGENCIES AND SHOULD NOT BE CONSTRUED TO REPRESENT AGENCY DETERMINATION OR POLICY.

Table 8—List of the 250 subwatersheds randomly chosen selected to determine the status and trend of inchannel processes. Includes the U.S. Geological Survey hydrologic unit code (USGS HUC), the aquatic province, the national forest (NF), Bureau of Land Management (BLM) unit, or national park (NP) that manages the land, the fifth-field watershed name, the sixth-field subwatershed name, and the percentage of the total subwatershed area that is federally owned.

USGS HUC	Province	Administrative unit	Watershed name	Subwatershed name	Year surveyed	Percent Federal
171003030503	Oregon Coast	Coos Bay BLM	Lake Creek	Upper Camp Creek	2003	90.13
171003030401	Oregon Coast	Coos Bay BLM	Middle Umpqua River	Paradise Creek	2007	49.47
171003050405	Oregon Coast	Coos Bay BLM	East Fork Coquille	Elk Cr	2008	36.25
171003050404	Oregon Coast	Coos Bay BLM	East Fork Coquille	Brewster Canyon	2003	45.01
171003050501	Oregon Coast	Coos Bay BLM	North Fork Coquille	North Coquille	2002	25.73
171003030704	Oregon Coast	Coos Bay BLM	Lower Smith River	Upper Lower Smith River		40.75
180102030101	High Cascades	Crater Lake NP	Wood River	E Fork Annie	2002	100.00
170703010207	High Cascades	Deschutes NF	Deschutes River / Browns Creek	Browns Creek	2009	100.00
170703010104	High Cascades	Deschutes NF	Deschutes River / Charleton Creek	Snow Creek	2002	100.00
170703020203	High Cascades	Deschutes NF	Crescent Creek	Summit Lake	2002	100.00
170703020204	High Cascades	Deschutes NF	Crescent Creek	Crescent Lake		100.00
170703010803	High Cascades	Deschutes NF	Squaw Creek	Upper Trout Creek	2005	100.00
170703010907	High Cascades	Deschutes NF	Upper Metolius River	Canyon Creek	2003	99.64
171002060301	Oregon Coast	Eugene BLM	Wildcat Creek	Upper Wildcat Creek	2005	33.35
170900060607	Western Cascades	Eugene BLM	South Santiam River	Owl Creek	2009	31.96
170900020304	Western Cascades	Eugene BLM	Upper Coast Fork Willamette River	Cottage Grove Reservoir	2006	29.04
170900020201	Western Cascades	Eugene BLM	Mosby Creek	Upper Mosby Creek	2005	30.29
170701051004	High Cascades	Gifford Pinchot NF	Little White Salmon River	Middle Little White Salmon River	2007	38.13
170800020202	Western Cascades	Gifford Pinchot NF	Muddy River	Clearwater Creek	2008	100.00

THE FINDINGS AND CONCLUSIONS IN THIS REPORT ARE IN PRESS AND SUBJECT TO CHANGE PRIOR TO FORMAL DISSEMINATION BY THE AGENCIES AND SHOULD NOT BE CONSTRUED TO REPRESENT AGENCY DETERMINATION OR POLICY.

USGS HUC	Province	Administrative unit	Watershed name	Subwatershed name	Year surveyed	Percent Federal
170800040402	Western Cascades	Gifford Pinchot NF	Upper Cispus River	Walupt Creek	2008	100.00
170800020503	Western Cascades	Gifford Pinchot NF	East Fork Lewis River	Copper Creek	2006	89.77
170800040205	Western Cascades	Gifford Pinchot NF	Upper Cowlitz River	Johnson Creek	2006	99.03
170800020203	Western Cascades	Gifford Pinchot NF	Muddy River	Elk Creek	2003	100.00
170800020108	Western Cascades	Gifford Pinchot NF	Upper Lewis River	Alec Creek	2003	100.00
170800020102	Western Cascades	Gifford Pinchot NF	Upper Lewis River	Twin Falls Creek	2003	100.00
170800040307	Western Cascades	Gifford Pinchot NF	Cowlitz Valley Frontal	Siler Creek		65.18
171100150110	Western Cascades	Gifford Pinchot NF	Upper Nisqually River	Little Nisqually River	2005	88.38
170800020401	Western Cascades	Gifford Pinchot NF	Yale Reservoir	Upper Siouxon Creek	2006	100.00
170800020404	Western Cascades	Gifford Pinchot NF	Yale Reservoir	Cougar Creek		43.73
170800040409	Western Cascades	Gifford Pinchot NF	Upper Cispus River	Blue Lake / Cispus River		100.00
170800040302	Western Cascades	Gifford Pinchot NF	Cowlitz Valley Frontal	Willame Creek	2002	98.96
170701051002	Western Cascades	Gifford Pinchot NF	Little White Salmon River	Big Lava Bed Frontal	2002	89.29
180102090203	Klamath / Siskiyou	Klamath NF	Indian Creek	East Fork Indian Creek	2005	96.22
180102090303	Klamath / Siskiyou	Klamath NF	Elk Creek	Lower Elk Creek	2007	97.46
180102090703	Klamath / Siskiyou	Klamath NF	Lower Salmon River	Somes Creek	2009	98.46
180102100401	Klamath / Siskiyou	Klamath NF	Lower Salmon River	Crapo Creek		97.08
180102060802	Klamath / Siskiyou	Klamath NF	Empire Creek	Humbug Creek		46.11
180102060803	Klamath / Siskiyou	Klamath NF	Empire Creek	Vesa Creek		60.73
180102080103	Klamath / Siskiyou	Klamath NF	East Fork Scott River	Noyes Valley	2008	38.94
180102090204	Klamath / Siskiyou	Klamath NF	Indian Creek	South Fork Indian Creek	2009	97.91
180102080402	Klamath / Siskiyou	Klamath NF	Moffett Creek	Indian Creek		46.20

THE FINDINGS AND CONCLUSIONS IN THIS REPORT ARE IN PRESS AND SUBJECT TO CHANGE PRIOR TO FORMAL DISSEMINATION BY THE AGENCIES AND SHOULD NOT BE CONSTRUED TO REPRESENT AGENCY DETERMINATION OR POLICY.

USGS HUC	Province	Administrative unit	Watershed name	Subwatershed name	Year surveyed	Percent Federal
180102050102	Klamath / Siskiyou	Klamath NF	Mount Shasta Woods	Horsethief Creek		48.05
180102090501	Klamath / Siskiyou	Klamath NF	Ukonom Creek	Oak Flat Creek	2006	98.06
180102090402	Klamath / Siskiyou	Klamath NF	Clear Creek	Tenmile Creek	2003	100.00
180102080203	Klamath / Siskiyou	Klamath NF	South Fork Scott River	Haynes Lake Creek	2003	27.77
180102100102	Klamath / Siskiyou	Klamath NF	South Fork Salmon River	Summerville	2002	98.42
180102100106	Klamath / Siskiyou	Klamath NF	South Fork Salmon River	Crawford Creek	2003	97.23
180102060903	Klamath / Siskiyou	Klamath NF	West Fork Beaver Creek	Bear Creek	2007	43.88
180102080601	Klamath / Siskiyou	Klamath NF	Bridge Flat	Emigrant Creek		35.78
180102080101	Klamath / Siskiyou	Klamath NF	East Fork Scott River	Upper East Fork Scott River	2006	30.82
180102090302	Klamath / Siskiyou	Klamath NF	Elk Creek	Upper Elk Creek	2006	100.00
180102090603	Klamath / Siskiyou	Klamath NF	Dillon Creek	Jackass Creek	2009	100.00
180102060502	High Cascades	Medford BLM	Klamath River / Iron Gate	Scotch Creek	2004	56.22
180102060405	High Cascades	Medford BLM	Jenny Creek	Keene Creek		44.55
171003090203	Klamath / Siskiyou	Medford BLM	Applegate River / Mckee Bridge	Applegate River / Star Gulch	2004	71.56
171003020804	Klamath / Siskiyou	Medford BLM	West Fork Cow Creek	West Fork Cow Creek / Bear Creek	2003	50.43
171003020603	Klamath / Siskiyou	Medford BLM	Upper Cow Creek	Upper Cow Cree / Galesville	2002	45.64
171003110502	Klamath / Siskiyou	Medford BLM	Deer Creek	Middle Deer Creek		33.50
171003100403	Klamath / Siskiyou	Medford BLM	Rogue River / Horseshoe Bend	Rogue River / Big Windy Creek	2003	95.39
171003080301	Klamath / Siskiyou	Medford BLM	Evans Creek	Upper Evans Creek	2005	30.46
171003020801	Oregon Coast	Medford BLM	West Fork Cow Creek	Upper West Fork Cow Creek	2002	44.44

THE FINDINGS AND CONCLUSIONS IN THIS REPORT ARE IN PRESS AND SUBJECT TO CHANGE PRIOR TO FORMAL DISSEMINATION BY THE AGENCIES AND SHOULD NOT BE CONSTRUED TO REPRESENT AGENCY DETERMINATION OR POLICY.

USGS HUC	Province	Administrative unit	Watershed name	Subwatershed name	Year surveyed	Percent Federal
171003070602	Western Cascades	Medford BLM	Trail Creek	West Fork Trail Creek	2003	42.84
171003110504	Klamath / Siskiyou	Medford NF	Deer Creek	Mc Mullin Creek	2009	30.27
171003100405	Klamath / Siskiyou	Medford NF	Rogue River / Horseshoe Bend	Kelsey Creek	2006	96.05
171003070809	Klamath / Siskiyou	Medford NF	Little Butte Creek	Little Butte / Lick	2009	36.45
171003070802	Klamath / Siskiyou	Medford NF	Little Butte Creek	Lower North Fork Little Butte Creek	2006	28.87
171003110304	Klamath / Siskiyou	Medford NF	Sucker Creek	Lower Sucker Creek	2006	35.11
171003090403	Klamath / Siskiyou	Medford NF	Middle Applegate River	Applegate River / Humbug Creek	2008	36.78
171003020803	Oregon Coast	Medford NF	West Fork Cow Creek	West Fork Cow Creek / Elk Valley Creek	2007	53.56
171003070504	Western Cascades	Medford NF	Elk Creek / Rogue River	Elk Creek / Flat Creek	2007	28.38
180101030202	Franciscan	Mendocino NF	Rice Fork	Lower Rice Fork		80.46
180201160202	Franciscan	Mendocino NF	North Fork Cache Creek	Bartlett Creek		57.20
180101030105	Klamath / Siskiyou	Mendocino NF	Lake Pillsbury	Anderson Creek		88.78
180101040201	Klamath / Siskiyou	Mendocino NF	Black Butte River	Upper Black Butte River	2002	78.15
180101040204	Klamath / Siskiyou	Mendocino NF	Black Butte River	Lower Black Butte River		66.81
180101040106	Klamath / Siskiyou	Mendocino NF	Wilderness	Howard Creek	2005	93.23
180101040103	Klamath / Siskiyou	Mendocino NF	Wilderness	Balm Of Gilead Creek		100.00
171100050805	North Cascades	Mt. Baker-Snoqualmie NF	Baker River	Lower Baker Lake	2009	98.29
171100060101	North Cascades	Mt. Baker-Snoqualmie NF	Upper Sauk River	Sloan Creek	2006	100.00
171100040301	North Cascades	Mt. Baker-Snoqualmie NF	South Fork Nooksack River	Upper South Fork Nooksack River	2006	93.34
171100050604	North Cascades	Mt. Baker-Snoqualmie NF	Cascade River	Middle Cascade River	2009	99.97

THE FINDINGS AND CONCLUSIONS IN THIS REPORT ARE IN PRESS AND SUBJECT TO CHANGE PRIOR TO FORMAL DISSEMINATION BY THE AGENCIES AND SHOULD NOT BE CONSTRUED TO REPRESENT AGENCY DETERMINATION OR POLICY.

USGS HUC	Province	Administrative unit	Watershed name	Subwatershed name	Year surveyed	Percent Federal
171100090201	North Cascades	Mt. Baker-Snoqualmie NF	Skykomish River Forks	Upper North Fork Skykomish River	2004	100.00
171100060106	North Cascades	Mt. Baker-Snoqualmie NF	Upper Sauk River	Lower White Chuck River	2003	100.00
171100050702	North Cascades	Mt. Baker-Snoqualmie NF	Skagit River / Illabot Creek	Skagit River At Corkindale		63.17
171100050806	North Cascades	Mt. Baker-Snoqualmie NF	Baker River	Lower Baker River / Lake Shannon	2007	49.84
171100100102	North Cascades	Mt. Baker-Snoqualmie NF	North Fork Snoqualmie River	North Fork Snoqualmie River / Sunday Creek		55.70
171100080102	North Cascades	Mt. Baker-Snoqualmie NF	North Fork Stillaguamish River	North Fork Stillaguamish River At Squire Creek	2005	27.96
171100090206	North Cascades	Mt. Baker-Snoqualmie NF	Skykomish River Forks	Lower South Fork Skykomish River	2007	65.23
171100040104	North Cascades	Mt. Baker-Snoqualmie NF	Upper North Fork Nooksack River	Glacier Creek	2005	78.04
171100100303	North Cascades	Mt. Baker-Snoqualmie NF	Middle Fork Snoqualmie River	Taylor River		97.91
171100090107	North Cascades	Mt. Baker-Snoqualmie NF	Tye And Beckler Rivers	Lower Beckler River	2007	74.45
171100090104	North Cascades	Mt. Baker-Snoqualmie NF	Tye And Beckler Rivers	Lower Tye River		72.25
171100130104	Western Cascades	Mt. Baker-Snoqualmie NF	Upper Green River	Upper Green River / Twin Camp Creek	2009	48.41
171100140202	Western Cascades	Mt. Baker-Snoqualmie NF	Lower White River	Clearwater River	2003	43.18
171100140105	Western Cascades	Mt. Baker-Snoqualmie NF	Upper White River	Upper Greenwater River	2007	100.00
171100140104	Western Cascades	Mt. Baker-Snoqualmie NF	Upper White River	Upper White River / Silver Creek	2002	86.09
171100130101	Western Cascades	Mt. Baker-Snoqualmie NF	Upper Green River	Green River Headwaters	2005	43.80
170701050601	High Cascades	Mt. Hood NF	East Fork Hood River	Upper East Fork Hood River	2008	100.00

THE FINDINGS AND CONCLUSIONS IN THIS REPORT ARE IN PRESS AND SUBJECT TO CHANGE PRIOR TO FORMAL DISSEMINATION BY THE AGENCIES AND SHOULD NOT BE CONSTRUED TO REPRESENT AGENCY DETERMINATION OR POLICY.

USGS HUC	Province	Administrative unit	Watershed name	Subwatershed name	Year surveyed	Percent Federal
170701050201	High Cascades	Mt. Hood NF	Fifteenmile Creek	Headwaters Fifteenmile Creek	2003	49.81
170703060901	High Cascades	Mt. Hood NF	Tygh Creek	Upper Badger Creek	2005	96.99
170900110101	Western Cascades	Mt. Hood NF	Collawash River	Upper Hot Springs Fork Collawash	2007	100.00
170800010102	Western Cascades	Mt. Hood NF	Salmon River	Draw Creek	2003	100.00
170900110304	Western Cascades	Mt. Hood NF	Oak Grove Fork Clackamas River	High Rock Creek	2006	100.00
170900110401	Western Cascades	Mt. Hood NF	Upper Clackamas River	Pot Creek	2009	99.35
170800010201	Western Cascades	Mt. Hood NF	Zigzag River	Still Creek	2002	98.23
170800010504	Western Cascades	Mt. Hood NF	Bull Run River	Cedar Creek	2003	96.22
170900110201	Western Cascades	Mt. Hood NF	Upper Clackamas River	Cub Creek	2005	100.00
170800010501	Western Cascades	Mt. Hood NF	Bull Run River	Blazed Alder Creek	2008	99.16
170900110301	Western Cascades	Mt. Hood NF	Oak Grove Fork Clackamas River	Upper Oak Grove Fork Clackamas River		42.05
170800010506	Western Cascades	Mt. Hood NF	Bull Run River	Middle Bull Run River	2009	100.00
171100150101	Western Cascades	Mt. Rainier NP	Upper Nisqually River	Nisqually Headwaters	2006	98.24
171100050401	North Cascades	North Cascades NP	Skagit River / Gorge Lake	Fisher Creek	2002	97.54
170200090111	North Cascades	North Cascades NP	Stehekin	Boulder Creek	2003	98.87
170200080203	North Cascades	Okanogan NF	Upper Methow River	Rattlesnake Creek		89.01
170200080703	North Cascades	Okanogan NF	Lower Methow River	Mainstem Lower Methow River	2003	93.15
170200080204	North Cascades	Okanogan NF	Upper Methow River	Cedar Creek		100.00
170200080102	North Cascades	Okanogan NF	Lost River	South Fork Lost River		100.00
170200080103	North Cascades	Okanogan NF	Lost River	Lower Lost River		99.54

THE FINDINGS AND CONCLUSIONS IN THIS REPORT ARE IN PRESS AND SUBJECT TO CHANGE PRIOR TO FORMAL DISSEMINATION BY THE AGENCIES AND SHOULD NOT BE CONSTRUED TO REPRESENT AGENCY DETERMINATION OR POLICY.

USGS HUC	Province	Administrative unit	Watershed name	Subwatershed name	Year surveyed	Percent Federal
170200080502	North Cascades	Okanogan NF	Twisp River	South Creek		99.10
171100180601	Olympic Peninsula	Olympic NF	Big Quilcene River	Upper Big Quilcene River	2004	99.78
171100180302	Olympic Peninsula	Olympic NF	Hamma Hamma River	Hamma Hamma River	2002	75.08
171001010501	Olympic Peninsula	Olympic NF	Calawah River	North Fork Calawah River		52.00
171001020107	Olympic Peninsula	Olympic NF	Queets River	Salmon River	2005	29.27
171100200304	Olympic Peninsula	Olympic NF	Dungeness River	Lower Gray Wolf River	2007	99.30
171100180701	Olympic Peninsula	Olympic NF	Upper West Hood Canal Frontal	Spencer / Marple Creek		36.59
171001020104	Olympic Peninsula	Olympic NF	Queets River	Sams River		98.36
171001010602	Olympic Peninsula	Olympic NP	Bogachiel River	Middle Bogachiel River		100.00
171001010402	Olympic Peninsula	Olympic NP	Sol Duc River	Headwaters Sol Duc River	2008	99.70
171001010401	Olympic Peninsula	Olympic NP	Sol Duc River	North Fork Sol Duc River	2007	100.00
171001020402	Olympic Peninsula	Olympic NP	Upper Quinalt River	Graves Creek	2009	100.00
180102110603	Klamath/Siskiyou	Redding BLM	Weaver / Rush	Grass Valley Creek	2007	59.75
180102110604	Klamath/Siskiyou	Redding BLM	Weaver / Rush	Indian Creek		33.83
171003100601	Franciscan	Rogue River-Siskiyou NF	Rogue River / Illahe Creek	Shasta Costa Ck	2004	99.67
171003120106	Franciscan	Rogue River-Siskiyou NF	Chetco River	Boulder Creek	2004	100.00
171003100801	Franciscan	Rogue River-Siskiyou NF	Rogue River / Illahe Creek	Rogue / Illahe	2009	92.28
171003111101	Franciscan	Rogue River-Siskiyou NF	Illinois River / Lawson Creek	Lawson Creek		98.27
171003110804	Franciscan	Rogue River-Siskiyou NF	Illinois River / Klondike Creek	Florence Creek	2009	100.00
171003120501	Franciscan	Rogue River-Siskiyou NF	Hunter Creek	Upper Hunter Creek	2007	74.58
171003070402	High Cascades	Rogue River-Siskiyou NF	Big Butte Creek	Clarks Fork Creek / Fourbit Creek	2004	79.00
171003070403	High Cascades	Rogue River-Siskiyou NF	Big Butte Creek	Willow Creek	2006	60.27

THE FINDINGS AND CONCLUSIONS IN THIS REPORT ARE IN PRESS AND SUBJECT TO CHANGE PRIOR TO FORMAL DISSEMINATION BY THE AGENCIES AND SHOULD NOT BE CONSTRUED TO REPRESENT AGENCY DETERMINATION OR POLICY.

USGS HUC	Province	Administrative unit	Watershed name	Subwatershed name	Year surveyed	Percent Federal
171003070203	High Cascades	Rogue River-Siskiyou NF	South Fork Rogue River	Upper Middle Fork Rogue River	2005	97.55
171003070105	High Cascades	Rogue River-Siskiyou NF	Upper Rogue River	Rogue River / Foster Creek	2008	100.00
171003070803	High Cascades	Rogue River-Siskiyou NF	Little Butte Creek	Upper South Fork Little Butte Creek	2005	100.00
171003070113	High Cascades	Rogue River-Siskiyou NF	Upper Rogue River	Rogue River / Barr Creek		40.19
171003070112	High Cascades	Rogue River-Siskiyou NF	Upper Rogue River	Mill Creek	2004	84.50
171003070110	High Cascades	Rogue River-Siskiyou NF	Upper Rogue River	Abbott Creek		97.03
171003110603	Klamath / Siskiyou	Rogue River-Siskiyou NF	Illinois River / Josephine Creek	Sixmile Creek	2003	97.44
171003110102	Klamath / Siskiyou	Rogue River-Siskiyou NF	East Fork Illinois River	Dunn Creek	2009	96.45
171003110303	Klamath / Siskiyou	Rogue River-Siskiyou NF	Sucker Creek	Grayback Creek	2007	79.80
171003090107	Klamath / Siskiyou	Rogue River-Siskiyou NF	Upper Applegate River	Lower Carberry	2005	71.94
171003090106	Klamath / Siskiyou	Rogue River-Siskiyou NF	Upper Applegate River	Steve Fork Carberry Creek	2002	93.72
171003110604	Klamath / Siskiyou	Rogue River-Siskiyou NF	Illinois River / Josephine Creek	Baker Creek	2005	97.94
171003080106	Klamath / Siskiyou	Rogue River-Siskiyou NF	Bear Creek	Ashland Creek	2003	73.08
171003090103	Klamath / Siskiyou	Rogue River-Siskiyou NF	Upper Applegate River	Elliott Creek / Dutch Creek	2008	67.21
171003110104	Klamath / Siskiyou	Rogue River-Siskiyou NF	East Fork Illinois River	Lower East Fork Illinois River	2003	34.59
171003050101	Oregon Coast	Rogue River-Siskiyou NF	Coquille S Fk, Lwr	Headwaters South Fork Coquille River	2002	72.23
171003050104	Oregon Coast	Rogue River-Siskiyou NF	Coquille S Fk, Lwr	Elk Creek	2009	48.09
171003020504	Klamath / Siskiyou	Roseburg BLM	South Umpqua River	Stouts Creek	2009	50.20
171003020506	Klamath / Siskiyou	Roseburg BLM	South Umpqua River	Upper Shively Oshea	2005	30.27
171003020901	Klamath / Siskiyou	Roseburg BLM	Lower Cow Creek	Middle Creek	2004	32.96
171003030106	Oregon Coast	Roseburg BLM	Upper Umpqua River	Yellow Creek	2008	40.98

THE FINDINGS AND CONCLUSIONS IN THIS REPORT ARE IN PRESS AND SUBJECT TO CHANGE PRIOR TO FORMAL DISSEMINATION BY THE AGENCIES AND SHOULD NOT BE CONSTRUED TO REPRESENT AGENCY DETERMINATION OR POLICY.

USGS HUC	Province	Administrative unit	Watershed name	Subwatershed name	Year surveyed	Percent Federal
171003010903	Western Cascades	Roseburg BLM	Canton Creek	Pass Creek	2009	79.59
171002050104	Oregon Coast	Salem BLM	Upper Alsea River	Upper South Fork Of Alsea River	2009	66.90
171002030201	Oregon Coast	Salem BLM	Nestucca River	Upper Nestucca River		52.81
171002040402	Oregon Coast	Salem BLM	Upper Siletz River	Lower North Fork Of Siletz River	2008	36.05
170900090503	Western Cascades	Salem BLM	Upper Molalla River	Molalla River / Pine Creek	2002	41.96
170900110601	Western Cascades	Salem BLM	Lower Clackamas River	Upper Clear Creek		25.18
180102120406	Klamath / Siskiyou	Shasta-Trinity NF	Lower Hayfork Creek	Grassy Flat Creek	2007	94.31
180102120302	Klamath / Siskiyou	Shasta-Trinity NF	Upper Hayfork Creek	North Fork Hayfork Creek		46.29
180200040106	Klamath / Siskiyou	Shasta-Trinity NF	Ash Creek	Lower Ash Creek	2008	90.03
180102120304	Klamath / Siskiyou	Shasta-Trinity NF	Upper Hayfork Creek	Gurley Gulch	2009	87.50
180200031103	Klamath / Siskiyou	Shasta-Trinity NF	Squaw Creek	Lower Squaw Creek		88.51
180102120402	Klamath / Siskiyou	Shasta-Trinity NF	Lower Hayfork Creek	Philpot Creek	2003	67.70
180102110403	Klamath / Siskiyou	Shasta-Trinity NF	Stuart Fork	Stoney Creek	2004	71.90
180102110605	Klamath / Siskiyou	Shasta-Trinity NF	Weaver / Rush	Weaver Creek	2008	49.81
180102120204	Klamath / Siskiyou	Shasta-Trinity NF	Middle South Fork Trinity River	Indian Valley Creek	2005	96.82
180200040102	Klamath / Siskiyou	Shasta-Trinity NF	Ash Creek	Upper Ash Creek		64.97
180102120103	Klamath / Siskiyou	Shasta-Trinity NF	Upper South Fork Trinity River	Upper South Fork Trinity River		94.98
180200040303	Klamath / Siskiyou	Shasta-Trinity NF	Squaw Valley Creek	Panther Creek	2007	40.11
180102110301	Klamath / Siskiyou	Shasta-Trinity NF	East Fork Trinity River	Upper East Fork Trinity River		30.89
180200040103	Klamath / Siskiyou	Shasta-Trinity NF	Ash Creek	Horse Creek		28.03

THE FINDINGS AND CONCLUSIONS IN THIS REPORT ARE IN PRESS AND SUBJECT TO CHANGE PRIOR TO FORMAL DISSEMINATION BY THE AGENCIES AND SHOULD NOT BE CONSTRUED TO REPRESENT AGENCY DETERMINATION OR POLICY.

USGS HUC	Province	Administrative unit	Watershed name	Subwatershed name	Year surveyed	Percent Federal
180102110102	Klamath / Siskiyou	Shasta-Trinity NF	Main Trinity River	Little Trinity River	2004	55.33
180102111102	Klamath / Siskiyou	Shasta-Trinity NF	Lower Trinity River	Little French Creek		95.83
180200050401	Klamath / Siskiyou	Shasta-Trinity NF	Sacramento Arm Shasta Lake	Middle Salt Creek		41.67
180102111101	Klamath / Siskiyou	Shasta-Trinity NF	Lower Trinity River	Sailor Bar Creek		94.16
180200050103	Klamath / Siskiyou	Shasta-Trinity NF	Box Canyon	South Fork Sacramento River		74.26
180102110503	Klamath / Siskiyou	Shasta-Trinity NF	Trinity Reservoir	Lower Trinity Reservoir		74.37
180102111103	Klamath / Siskiyou	Shasta-Trinity NF	Lower Trinity River	Big French Creek	2007	100.00
180200031202	Klamath / Siskiyou	Shasta-Trinity NF	Pit Arm Shasta Lake	Potem Creek	2009	65.56
171002050405	Oregon Coast	Siuslaw NF	Lower Alsea River	Alsea River / Eckman Creek	2008	31.35
171002050302	Oregon Coast	Siuslaw NF	Drift Creek	Middle Drift Creek / Alsea River		36.59
171002060501	Oregon Coast	Siuslaw NF	Deadwood Creek	Upper Deadwood Creek	2006	55.43
171002050704	Oregon Coast	Siuslaw NF	Cummins Creek/tenmile Creek / Mercer Lake Frontal	Mercer Lake		58.62
171002030204	Oregon Coast	Siuslaw NF	Nestucca River	Nestucca River / Niagara Creek	2005	75.74
171003030706	Oregon Coast	Siuslaw NF	Lower Smith River	Lower North Fork Smith River		32.91
171002050202	Oregon Coast	Siuslaw NF	Five Rivers / Lobster Creek	Upper Five Rivers	2003	77.30
171002060602	Oregon Coast	Siuslaw NF	Indian Creek / Lake Creek	Lower Indian Creek	2005	53.66
180101050201	Franciscan	Six Rivers NF	Upper North Fork Eel River	Headwaters North Fork Eel River	2002	84.57
180101010301	Klamath / Siskiyou	Six Rivers NF	South Fork Smith River	Prescott Fork		99.02
180101010104	Klamath / Siskiyou	Six Rivers NF	North Fork Smith River	Peridotite Canyon		95.84

THE FINDINGS AND CONCLUSIONS IN THIS REPORT ARE IN PRESS AND SUBJECT TO CHANGE PRIOR TO FORMAL DISSEMINATION BY THE AGENCIES AND SHOULD NOT BE CONSTRUED TO REPRESENT AGENCY DETERMINATION OR POLICY.

USGS HUC	Province	Administrative unit	Watershed name	Subwatershed name	Year surveyed	Percent Federal
180102120505	Klamath / Siskiyou	Six Rivers NF	Grouse / Madden	Lower South Fork Trinity River		88.43
180101010204	Klamath / Siskiyou	Six Rivers NF	Middle Fork Smith River	Shelley Creek	2002	87.71
180102090801	Klamath / Siskiyou	Six Rivers NF	Bluff Creek	Cedar Creek	2008	94.91
180102091005	Klamath / Siskiyou	Six Rivers NF	Blue Creek	Lower Blue Creek		35.56
180102111203	Klamath / Siskiyou	Six Rivers NF	Trinity / South Fork To Tish Tang	Horse Linto Creek	2005	96.80
180201160503	Franciscan	Ukiah BLM	Lakeport	Lower Scotts Creek		34.01
171003010402	High Cascades	Umpqua NF	Clearwater	Bear Creek	2004	100.00
171003010103	High Cascades	Umpqua NF	Diamond Lake	Diamond Lake		100.00
171003010301	High Cascades	Umpqua NF	Upper North Umpqua	Warm Springs Creek	2005	100.00
171003010708	Western Cascades	Umpqua NF	Middle North Umpqua	Blitzen Facial	2009	99.52
171003020203	Western Cascades	Umpqua NF	Jackson Creek	Squaw Creek	2006	100.00
171003020302	Western Cascades	Umpqua NF	Middle South Umpqua River	Dumont Creek	2002	100.00
171003010801	Western Cascades	Umpqua NF	Steamboat Creek	Steamboat Headwaters	2004	96.86
171003011104	Western Cascades	Umpqua NF	Little River	Emile Creek	2004	82.88
171003011106	Western Cascades	Umpqua NF	Little River	Upper Cavitt Creek	2007	100.00
171003020403	Western Cascades	Umpqua NF	Elk Creek / South Umpqua	Drew Creek	2005	92.57
170900020101	Western Cascades	Umpqua NF	Row River	Layng Creek	2004	88.76
171003011101	Western Cascades	Umpqua NF	Little River	Little River Headwaters	2003	98.59
171003010501	Western Cascades	Umpqua NF	Fish Creek	Fish Creek Headwaters	2008	100.00
170200090203	North Cascades	Wenatchee NF	Upper Chelan	Fish Creek	2004	98.95
170300010301	North Cascades	Wenatchee NF	Middle Upper Yakima River	Swauk Creek	2002	74.68
170200110303	North Cascades	Wenatchee NF	Nason / Tumwater	Chiwaukum Creek	2008	89.97

THE FINDINGS AND CONCLUSIONS IN THIS REPORT ARE IN PRESS AND SUBJECT TO CHANGE PRIOR TO FORMAL DISSEMINATION BY THE AGENCIES AND SHOULD NOT BE CONSTRUED TO REPRESENT AGENCY DETERMINATION OR POLICY.


USGS HUC	Province	Administrative unit	Watershed name	Subwatershed name	Year surveyed	Percent Federal
170200110302	North Cascades	Wenatchee NF	Nason/tumwater	Upper Nason Creek	2009	90.14
170200110404	North Cascades	Wenatchee NF	Icicle / Chumstick	Chumstick Creek		45.30
170200110202	North Cascades	Wenatchee NF	Chiawa River	Middle Chiawa River		100.00
170200110203	North Cascades	Wenatchee NF	Chiawa River	Lower Chiawa River	2008	87.36
170300020303	North Cascades	Wenatchee NF	Naches River / Tieton River	North Fork Tieton River	2009	100.00
170900040501	Western Cascades	Willamette NF	Mckenzie River / Quartz Creek	Quartz Creek	2003	31.71
170900040201	Western Cascades	Willamette NF	Horse Creek	Upper Separation Creek	2004	100.00
170900060401	Western Cascades	Willamette NF	Quartzville Creek	Upper Quartzville Creek	2002	100.00
170900050301	Western Cascades	Willamette NF	Detroit Reservoir / Blow Out Divide Creek	Upper Blowout Creek	2008	99.81
170900050304	Western Cascades	Willamette NF	Detroit Reservoir / Blow Out Divide Creek	Detroit Reservoir / Kinney Creek	2009	39.79
170900010603	Western Cascades	Willamette NF	North Fork Of Middle Fork Willamette River	North Fork Of Middle Fork Willamette River / Fisher Creek	2003	100.00
170900050202	Western Cascades	Willamette NF	North Fork Breitenbush River	North Fork Breitenbush River	2006	94.39
170900050203	Western Cascades	Willamette NF	North Fork Breitenbush River	Humbug Creek	2006	100.00
170900010303	Western Cascades	Willamette NF	Salt Creek / Willamette River	Lower Salt Creek	2005	100.00
170900010902	Western Cascades	Willamette NF	Fall Creek	Fall Creek / Hehe Creek	2004	97.09
170900040302	Western Cascades	Willamette NF	South Fork Mckenzie River	South Fork Mckenzie River / Elk Creek	2009	100.00
170900010504	Western Cascades	Willamette NF	Hills Creek Reservoir	Middle Fork Willamette River / Larison Creek	2003	95.90

THE FINDINGS AND CONCLUSIONS IN THIS REPORT ARE IN PRESS AND SUBJECT TO CHANGE PRIOR TO FORMAL DISSEMINATION BY THE AGENCIES AND SHOULD NOT BE CONSTRUED TO REPRESENT AGENCY DETERMINATION OR POLICY.



USGS HUC	Province	Administrative unit	Watershed name	Subwatershed name	Year surveyed	Percent Federal
170900040102	Western Cascades	Willamette NF	Upper Mckenzie River	Hackleman Creek	2004	94.34
170900040307	Western Cascades	Willamette NF	South Fork Mckenzie River	South Fork Mckenzie River / Cougar Reservoir	2008	90.40
170900060503	Western Cascades	Willamette NF	Middle Santiam River	Donaca Creek	2002	86.19
170900050503	Western Cascades	Willamette NF	Little North Santiam River	Gold Creek	2005	100.00
170900050107	Western Cascades	Willamette NF	Upper North Santiam River	Boulder Creek / Marys Creek	2005	64.15
170900010702	Western Cascades	Willamette NF	Middle Fork Willamette / Lookout Point	Lookout Point Reservoir	2007	71.64
170900010106	Western Cascades	Willamette NF	Upper Middle Fork Willamette River	Lower Middle Fork Willamette River	2003	65.59
170900060604	Western Cascades	Willamette NF	South Santiam River	Trout Creek	2006	92.03
170900040107	Western Cascades	Willamette NF	Upper Mckenzie River	White Branch	2006	100.00
170900010202	Western Cascades	Willamette NF	Hills Creek	Upper Hills Creek	2009	100.00
180102030201	High Cascades	Winema NF	Klamath Lake	Threemile	2003	90.59

THE FINDINGS AND CONCLUSIONS IN THIS REPORT ARE IN PRESS AND SUBJECT TO CHANGE PRIOR TO FORMAL DISSEMINATION BY THE AGENCIES AND SHOULD NOT BE CONSTRUED TO REPRESENT AGENCY DETERMINATION OR POLICY.

Table 9—Evaluation criteria used in watershed-level decision-support model for each aquatic province.

Attribute and measure	Data value	Evaluated score	Curve shape	Source
High Cascades Province				
Roads—watershed road density Road mi/watershed area mi ²	0.5	+1		AREMP workshop 4/15-17/2008
	2	0		
	4	-1		
Roads—riparian—all streams 120-ft buffer Road mi/stream mi	0.01	+1		AREMP workshop 9/16/2009
	0.1	-1		
Roads—riparian—perennial streams 180-ft buffer Road mi/perennial stream mi	0.01	+1		AREMP workshop 4/15-17/2008
	0.1	-1		
Roads—crossings Number/stream mi	0.01	+1		AREMP workshop 4/15-17/2008
	1.25	-1		
Mass wasting—Landslide risk Change in average landslide density (per km ²) from an optimum forested, unroaded state	0.1	+1		AREMP workshop 4/15-17/2008
	0.3	-1		
Mass wasting—road crossings in high-risk areas Number/stream mi	0.01	+1		AREMP workshop 9/16/2009
	0.3	-1		
Vegetation—riparian QMD ¹ ≥ 8/14-inch (lodgepole/other) and canopy cover ≥ 40 percent, all species, 160-ft buffer mi ² /riparian forest-capable area	0.3	-1		AREMP workshop 4/15-17/2008
	0.5	0		
	0.7	+1		
Vegetation—watershed— rain on snow band QMD ≥ 8-inch and cover ≥ 40/70 % (dry/wet zone) mi ² /watershed forest-capable mi ²	0.35	-1		AREMP workshop 4/15-17/2008 Grant et al. (2008)
	0.85	+1		
Vegetation—watershed— rain only and snow only bands, QMD ≥ 8-inch and cover ≥ 40/70 % (dry/wet zone) mi ² /watershed forest-capable mi ²	0.1	-1		AREMP workshop 4/15-17/2008 Grant et al. (2008)
	0.45	+1		

THE FINDINGS AND CONCLUSIONS IN THIS REPORT ARE IN PRESS AND SUBJECT TO CHANGE PRIOR TO FORMAL DISSEMINATION BY THE AGENCIES AND SHOULD NOT BE CONSTRUED TO REPRESENT AGENCY DETERMINATION OR POLICY.

Attribute and measure	Data value	Evaluated score	Curve shape	Source
Klamath/Siskiyou and Franciscan Provinces				
Roads–crossings Number/stream mi	0.5	+1		AREMP workshop 2003
	2	-1		
Roads–riparian, 160-ft buffer Road mi/riparian area mi ²	0.4	+1		AREMP workshop 10/14/2009
	1	-1		
Roads– lower 1/3 of slope Road mi/lower 1/3 slope mi ²	1	+1		AREMP workshop 2003
	2	-1		
Roads–watershed, designated motorized off highway vehicle trail Trail mi/watershed area mi ²	0.5	+1		AREMP workshop 3/18/2008
	1.25	0		
	2	-1		
Roads–watershed Road mi/watershed area mi ²	1	+1		AREMP workshop 2003
	2.5	0		
	4	-1		
Vegetation–riparian cover– mixed species, 160-ft buffer Average canopy cover (%)	40	-1		AREMP workshop 10/14/2009
	60	+1		
Vegetation–riparian cover–oak woodlands 160-ft buffer Average canopy cover (%)	20	-1		AREMP analysis
	50	+1		
Vegetation–watershed–early seral (QMD < 5-inch) mi ² /watershed forest-capable mi ²	0.05	+1		AREMP workshop 3/18/2008
	0.25	-1		
Vegetation–watershed – < 7,000 ft elevation– mixed species Average canopy cover (%)	40	-1		AREMP workshop 10/14/2009
	70	+1		
Vegetation–watershed–< 7,000-ft elevation –oak woodlands Average canopy cover (%)	10	-1		AREMP workshop 10/14/2009
	40	+1		
Vegetation–watershed–≥ 7,000 ft elevation–mixed species Average canopy cover (%)	20	-1		AREMP workshop 10/14/2009
	40	+1		
Vegetation–watershed–≥ 7,000 ft elevation–oak woodlands Average canopy cover (%)	40	-1		AREMP workshop 10/14/2009
	70	+1		

THE FINDINGS AND CONCLUSIONS IN THIS REPORT ARE IN PRESS AND SUBJECT TO CHANGE PRIOR TO FORMAL DISSEMINATION BY THE AGENCIES AND SHOULD NOT BE CONSTRUED TO REPRESENT AGENCY DETERMINATION OR POLICY.

North Cascades Province

Landslide risk—roads and vegetation Change in average landslide density (per km ²) from an optimum forested, unroaded state	0.1	+1		AREMP analysis
	0.3	-1		
Roads—crossings # crossings/stream mi	0	+1		AREMP workshop 9/19/2008
	1.25	-1		
Roads—riparian—< 3 % gradient 30-m buffer Proportion of stream mi w/road in buffer	0	+1		AREMP workshop 9/19/2008
	0.05	-1		
Roads—riparian—≥ 3 % gradient 100-m buffer Proportion of stream mi w/road in buffer	0	+1		AREMP workshop 9/19/2008
	0.05	-1		
Road density— high-hazard areas Road mi/watershed area mi ²	0	+1		AREMP workshop 9/19/2008
	1.2	-1		
Road density— low-hazard areas Road mi/watershed area mi ²	0.7	+1		AREMP workshop 9/19/2008
	2.4	-1		
Vegetation—riparian east, QMD ≥12-inch, all species, 100-ft buffer mi ² /riparian forest-capable mi ²	0.3	-1		AREMP workshop 9/19/2008
	0.5	+1		
	0.8	+1		
	1	0		
Vegetation—riparian west, QMD ≥16-inch, all species, 160-ft buffer mi ² /riparian forest-capable mi ²	0.3	-1		AREMP workshop 9/19/2008
	0.6	+1		
Vegetation—watershed east, wet zone, QMD < 5-inch, all species mi ² /watershed forest capable mi ²	0.2	+1		AREMP workshop 2003
	0.3	-1		
Vegetation—watershed, east, dry zone, QMD ≥ 20-inch, conifers mi ² /watershed forest- capable mi ²	0.3	-1		AREMP workshop 9/19/2008
	0.6	+1		
Vegetation—watershed west ≥ 70 percent cover, conifers mi ² /watershed forest- capable mi ²	0.65	-1		AREMP workshop 2003
	0.88	+1		



THE FINDINGS AND CONCLUSIONS IN THIS REPORT ARE IN PRESS AND SUBJECT TO CHANGE PRIOR TO FORMAL DISSEMINATION BY THE AGENCIES AND SHOULD NOT BE CONSTRUED TO REPRESENT AGENCY DETERMINATION OR POLICY.

Olympic Peninsula Province

Landslide risk—roads and vegetation Change in average landslide density (per km ²) from an optimum forested, unroaded state	0.1	+1		AREMP analysis
	0.3	-1		
Roads—crossings Number/stream mi	0.25	+1		AREMP workshop follow-up 2003
	1	-1		
Roads—riparian, 160-ft buffer Road mi/stream mi	0	+1		AREMP workshop 4/10/08
	0.1	-1		
Roads—watershed Road mi/watershed area mi ²	1	+1		AREMP workshop 4/10/08
	3	-1		
Vegetation—riparian, QMD ≥20-inch (wet zone) QMD ≥12-inch (dry zone) 160-ft buffer mi ² /riparian forest-capable mi ²	0.25	-1		AREMP workshop 4/10/08
	0.65	+1		
Vegetation—urban/agriculture mi ² /watershed area mi ²	0.2	+1		AREMP workshop 4/10/08
	0.4	-1		
Vegetation—watershed, QMD < 5-inch mi ² /watershed forest-capable mi ²	0.05	+1		AREMP workshop 6/16/2009
	0.25	-1		

THE FINDINGS AND CONCLUSIONS IN THIS REPORT ARE IN PRESS AND SUBJECT TO CHANGE PRIOR TO FORMAL DISSEMINATION BY THE AGENCIES AND SHOULD NOT BE CONSTRUED TO REPRESENT AGENCY DETERMINATION OR POLICY.

Oregon Coast Province

Upslope–road density Road mi/watershed area mi ²	2.2	+1		Kaufman and Hughes 2006
	4.2	-1		
Upslope–urban/agriculture mi ² /watershed area mi ²	0.2	+1		AREMP workshop 2003
	0.4	-1		
Upslope–Landslide risk Difference in average landslide density (per km ²) from an optimum forested, unroaded state	0.1	+1		AREMP analysis
	0.3	-1		
Riparian–all streams–canopy cover 160-ft buffer, all species Average cover %	0.55	-1		AREMP analysis
	0.7	+1		
Riparian–all streams–conifers ≥20-inch QMD, 160-ft buffer mi ² /riparian forest-capable mi ²	0.25	-1		AREMP workshop 2003
	0.5	+1		
Riparian–perennial streams– QMD conifers ≥20-inch 300-ft buffer mi ² /riparian forest capable mi ²	0.25	-1		AREMP workshop 3/4/2008
	0.5	+1		
Riparian–perennial streams – road density 300-ft buffer Road mi/stream mi	0.075	+1		AREMP workshop 3/4/2008
	0.4	-1		

THE FINDINGS AND CONCLUSIONS IN THIS REPORT ARE IN PRESS AND SUBJECT TO CHANGE PRIOR TO FORMAL DISSEMINATION BY THE AGENCIES AND SHOULD NOT BE CONSTRUED TO REPRESENT AGENCY DETERMINATION OR POLICY.

Western Cascades Province

Roads–watershed road density Road mi/watershed area mi ²	0.5	+1		AREMP workshop 4/15-17/2008
	2	0		
	4	-1		
Roads–riparian–all streams 120-ft buffer Road mi/stream mi	0.01	+1		AREMP workshop 9/16/2009
	0.1	-1		
Roads–riparian–perennial streams 180-ft buffer Road mi/perennial stream mi	0.01	+1		AREMP workshop 4/15-17/2008
	0.1	-1		
Roads–crossings Number/stream mi	0.01	+1		AREMP workshop 4/15-17/2008
	1.25	-1		
Mass wasting–Landslide risk Change in average landslide density (per km ²) from an optimum forested, unroaded state	0.1	+1		AREMP workshop 4/15-17/2008
	0.3	-1		
Mass wasting–road crossings in high-risk areas Number/stream mi	0.01	+1		AREMP workshop 9/16/2009
	0.3	-1		
Vegetation–riparian QMD ≥ 8/14-inch (lodgepole/other) and canopy closure ≥ 40 %, all species, 160-ft buffer mi ² /riparian forest capable mi ²	0.5	-1		AREMP workshop 4/15-17/2008
	0.7	+1		
Vegetation–watershed– rain on snow band QMD ≥ 8-inch and cover ≥ 70% mi ² /watershed forest-capable mi ²	0.35	-1		AREMP workshop 4/15-17/2008 Grant et al. (2008)
	0.85	+1		
Vegetation–watershed– rain only and snow only bands, QMD ≥ 8-inch and cover ≥ 70% mi ² /watershed forest-capable mi ²	0.1	-1		AREMP workshop 4/15-17/2008 Grant et al. (2008)
	0.45	+1		

¹ QMD = quadratic mean diameter of all dominant and codominant trees.

Table 10—Evaluation criteria used in inchannel-level decision-support models by aquatic province.

Attribute and measure	Data value	Evaluated score	Curve shape	Source
High Cascades Province				
Amphibians Aquatic and terrestrial index	33	0		AREMP analysis
	100	+1		
Fines, < 3 percent gradient percent fines	0	-1		AREMP workshop 2003
	10	+1		
	20	+1		
	30	-1		
Fines 3 to 6 % gradient percent fines	20	+1		AREMP workshop 2003
	35	-1		
Flood-plain connectivity, < 3 % gradient No side channels No side channels visible Indeterminate Probably side channels visible Side channels obviously connected		-1		AREMP workshop 2008
		-0.5		
		0		
		0.5		
		+1		
Macroinvertebrates–EPT Percentage of taxa in <i>ephemeroptera</i> , <i>plecoptera</i> , and <i>triptoptera</i> (EPT)	61	-1		Herlihy (2009) analysis of reference sites
	72	+1		
Macroinvertebrates–intolerant Percentage of taxa	35	-1		Herlihy (2009) analysis of reference sites
	50	+1		
Pool frequency, < 3 percent gradient Bankfull widths per pool	5	+1		AREMP workshop 2003
	7	-1		
Pool frequency, 3 to 6 % gradient Bankfull widths per pool	2	+1		AREMP workshop 2003
	6	-1		
Water temperature, bull trout are not present Deg C	16	+1		AREMP workshop 2008
	18	0		
	23	-1		
Water temperature, bull trout are present Deg C	9	+1		AREMP workshop 2008
	12	0		
	13	-1		
Wood frequency, North,	1	-1		AREMP workshop 2008

THE FINDINGS AND CONCLUSIONS IN THIS REPORT ARE IN PRESS AND SUBJECT TO CHANGE PRIOR TO FORMAL DISSEMINATION BY THE AGENCIES AND SHOULD NOT BE CONSTRUED TO REPRESENT AGENCY DETERMINATION OR POLICY.

Min 12-inch diameter x 25-ft length Pieces/100 m	4	+1		
Wood frequency, South Min 18-inch diameter x 25-ft length Pieces/100 m	1	-1		AREMP workshop 2008
	4	+1		

DRAFT

THE FINDINGS AND CONCLUSIONS IN THIS REPORT ARE IN PRESS AND SUBJECT TO CHANGE PRIOR TO FORMAL DISSEMINATION BY THE AGENCIES AND SHOULD NOT BE CONSTRUED TO REPRESENT AGENCY DETERMINATION OR POLICY.

Klamath/Siskiyou and Franciscan Provinces


Amphibians Aquatic and terrestrial index	33	0		AREMP analysis
	100	+1		
Bankfull width:depth, < 4 % gradient Ratio	15	+1		AREMP workshop 3/2008
	35	-1		
D50 (median particle size) < 4 % gradient mm	2	-1		AREMP workshop 3/2008
	45	+1		
	200	+1		
	350	-1		
Fines < 4 % gradient % fines	30	-1		AREMP workshop 3/2008
	10	+1		
Macroinvertebrates–EPT Percentage of taxa in <i>ephemeroptera, plecoptera, and</i> <i>triptoptera</i> (EPT)	61	-1		Herlihy (2009) analysis of reference sites
	72	+1		
Macroinvertebrates,–intolerant Percentage of taxa	35	-1		Herlihy (2009) analysis of reference sites
	50	+1		
Pool frequency, < 4 % gradient and within unconfined valley type Bankfull widths per pool	5	+1		AREMP workshop 2003
	12	-1		
Water temperature Deg C	18	1		AREMP workshop 3/2008
	20	0.8		
	21	0		
	24	-1		
Wood frequency, Min 12-inch diameter x 25-ft length Pieces/100 m	1	-1		AREMP workshop 3/2008
	3	+1		
Wood frequency, Min 24-inch diameter (small end) x 50-ft length Pieces/100 m	0	-1		AREMP workshop 3/2008
	0.1	+1		

THE FINDINGS AND CONCLUSIONS IN THIS REPORT ARE IN PRESS AND SUBJECT TO CHANGE PRIOR TO FORMAL DISSEMINATION BY THE AGENCIES AND SHOULD NOT BE CONSTRUED TO REPRESENT AGENCY DETERMINATION OR POLICY.

North Cascades Province

Amphibians Aquatic and terrestrial index	33	0		AREMP analysis
	100	+1		
Bankfull width:depth, ≤ 2 % gradient, nonglacial Ratio	39.9	+1		AREMP Workshop 4/28/03
	40	-1		
Mean particle size (D50) ≤ 2 % gradient, nonglacial mm	20	-1		AREMP Workshop 4/28/03
	30	+1		
	100	+1		
	500	-1		
Dissolved oxygen mg/L	5	-1		AREMP Workshop 4/28/03
	10	+1		
Entrenchment, east ≤ 2 % gradient, nonglacial Ratio	1.4	-1		AREMP Workshop 4/28/03
	1.5	+1		
Entrenchment, west ≤ 2 % gradient, nonglacial Ratio	1.2	-1		AREMP Workshop 4/28/03
	1.3	+1		
Fines ≤ 2 % gradient, nonglacial	11	+1		AREMP Workshop 4/28/03
	17	-1		
Macroinvertebrates–EPT Percentage of taxa in <i>ephemeroptera, plecoptera, and</i> <i>triptoptera</i> (EPT)	61	-1		Herlihy (2009) analysis of reference sites
	72	+1		
Macroinvertebrates–intolerant Percentage of taxa	35	-1		Herlihy (2009) analysis of reference sites
	50	+1		
Pool frequency, east > 2 % gradient Bankfull widths per pool	0.9	+1		AREMP Workshop 2004
	1	0		
	4	0		
	4.1	-1		
Pool frequency, west ≤ 2 % gradient Bankfull widths per pool	4.9	+1		AREMP Workshop 2004
	5	0		
	7	0		
	7.1	-1		
Pool frequency, west Bankfull widths per pool	4	+1		AREMP Workshop 2004
	5	0		
	14	0		
	18	-1		
Water temperature, bull trout are present Deg C	3	-1		AREMP Workshop 2004
	6	+1		
	9	+1		
	13	-1		

THE FINDINGS AND CONCLUSIONS IN THIS REPORT ARE IN PRESS AND SUBJECT TO CHANGE PRIOR TO FORMAL DISSEMINATION BY THE AGENCIES AND SHOULD NOT BE CONSTRUED TO REPRESENT AGENCY DETERMINATION OR POLICY.

Water temperature Bull trout are not present Deg C	4	-1		AREMP Workshop 2004
	6	+1		
	15	+1		
	18	-1		
Wood frequency, east, Min 12-inch diameter x 25-ft length, wetter subsections Pieces/100 m	1.6	-1		AREMP Workshop 2004
	3.1	0		
	4.5	+1		
Wood frequency, east, min 12-inch diameter x 25-ft length, drier subsections Pieces/100 m	0.9	-1		AREMP Workshop 2004
	1.9	0		
	2.8	+1		
Wood frequency, west, min 12-inch diameter (small end) x 25-ft length, > 2 percent gradient Pieces/100 m	0.5	-1		AREMP Workshop 2004
	1	0		
	5	+1		
Wood frequency, west, min 12-inch diameter (small end) x 25-ft length, ≤ 2 percent gradient Pieces/100 m	0.5	-1		AREMP Workshop 2004
	2.5	0		
	7.5	+1		


THE FINDINGS AND CONCLUSIONS IN THIS REPORT ARE IN PRESS AND SUBJECT TO CHANGE PRIOR TO FORMAL DISSEMINATION BY THE AGENCIES AND SHOULD NOT BE CONSTRUED TO REPRESENT AGENCY DETERMINATION OR POLICY.

Olympic Peninsula Province

Amphibians	33	0		AREMP analysis 2009
Aquatic & terrestrial	100	+1		
Mean particle size (D50) ≤ 2 percent gradient, unconstrained mm	45	-1		AREMP workshop 2003
	65	+1		
	95	+1		
	128	-1		
Fines, ≤ 2 % gradient, unconstrained Percent	11	+1		AREMP workshop 2003
	17	-1		
Macroinvertebrates–EPT Percentage of taxa in <i>ephemeroptera, plecoptera, and</i> <i>triptoptera</i> (EPT)	61	-1		Herlihy (2009) analysis of reference sites
	72	+1		
Macroinvertebrates–intolerant Percentage of taxa	35	-1		Herlihy (2009) analysis of reference sites
	50	+1		
Pool frequency Bankfull widths/pool	8	+1		ODFW benchmarks
	20	-1		
Water temperature, bull trout are not present Deg C	16	+1		AREMP workshop 2008
	18	+1		
	23	-1		
Water temperature, bull trout are present Deg C	9	+1		AREMP workshop 2008
	12	0		
	13	-1		
Wood frequency, min 12-inch diameter (small end) x 25-ft length, > 2 % gradient Pieces/100 m	0.5	-1		AREMP workshop 2008
	1	0		
	5	+1		
Wood frequency, min 12-inch diameter (small end) x 25' length, ≤ 2 % gradient Pieces / 100 m	0.5	-1		AREMP workshop 2008
	2.5	0		
	7.5	+1		


THE FINDINGS AND CONCLUSIONS IN THIS REPORT ARE IN PRESS AND SUBJECT TO CHANGE PRIOR TO FORMAL DISSEMINATION BY THE AGENCIES AND SHOULD NOT BE CONSTRUED TO REPRESENT AGENCY DETERMINATION OR POLICY.

Oregon Coast Province

Amphibians Aquatic & terrestrial	33	0		AREMP analysis 2009
	100	+1		
Fines percent	11	+1		AREMP workshop 2003
	17	-1		
Flood-plain connectivity, < 3 % gradient No side channels		-1		AREMP workshop 2008
	No side channels visible	-0.5		
	Indeterminate	0		
	Probably side channels visible	0.5		
	Side channels obviously connected	+1		
Macroinvertebrates–EPT Percentage of taxa in <i>ephemeroptera</i> , <i>plecoptera</i> , and <i>triptoptera</i> (EPT)	61	-1		Herlihy (2009) analysis of reference sites
	72	+1		
Macroinvertebrates–intolerant Percentage of taxa	35	-1		Herlihy (2009) analysis of reference sites
	50	+1		
Pool depth meters	0.35	-1		AREMP workshop 2003
	0.75	+1		
Pool frequency Bankfull widths per pool	8	+1		AREMP workshop 2003
	20	-1		
Water temperature, Maximum 7-day average Deg C	16	1		AREMP workshop 2003
	23	-1		
Wood frequency, min 24-inch diameter (small end) x 33-ft length Pieces/100 m	1	-1		OR DFW standard
	3	+1		

THE FINDINGS AND CONCLUSIONS IN THIS REPORT ARE IN PRESS AND SUBJECT TO CHANGE PRIOR TO FORMAL DISSEMINATION BY THE AGENCIES AND SHOULD NOT BE CONSTRUED TO REPRESENT AGENCY DETERMINATION OR POLICY.

**Western Cascades
Province**

Amphibians Aquatic & terrestrial	33	0		AREMP analysis
	100	+1		
Mean particle size (D50) < 3 % gradient mm	40	-1		AREMP workshop 2003
	60	+1		
	140	+1		
	200	-1		
Fines, < 3 % gradient % fines	0	-1		AREMP workshop 2003
	10	+1		
	20	+1		
	30	-1		
Fines, 3-6 % gradient % fines	15	+1		AREMP workshop 2003
	30	-1		
Flood-plain connectivity, < 3 % gradient No side channels		-1		AREMP workshop 2008
	No side channels visible	-0.5		
	Indeterminate	0		
	Probably side channels visible	0.5		
	Side channels obviously connected	+1		
Macroinvertebrates–EPT Percentage of taxa in <i>ephemeroptera, plecoptera, and</i> <i>triptoptera</i> (EPT)	61	-1		Herlihy (2009) analysis of reference sites
	72	+1		
Macroinvertebrates-intolerant Percentage of taxa	35	-1		Herlihy (2009) analysis of reference sites
	50	+1		
Pool frequency, < 3 % gradient Bankfull widths per pool	5	+1		AREMP workshop 2003
	7	-1		
Pool frequency 3-6 % gradient Bankfull widths per pool	2	+1		AREMP workshop 2003
	6	-1		
Water temperature, bull trout are not present Deg C	16	+1		AREMP workshop 2008
	18	+1		
	23	-1		
Water temperature, bull trout are present Deg C	9	+1		AREMP workshop 2008
	12	0		
	13	-1		
Wood frequency, 24-inch dbh x 50-ft Pieces/100 m	1	-1		AREMP workshop 2008

THE FINDINGS AND CONCLUSIONS IN THIS REPORT ARE IN PRESS AND SUBJECT TO CHANGE PRIOR TO FORMAL DISSEMINATION BY THE AGENCIES AND SHOULD NOT BE CONSTRUED TO REPRESENT AGENCY DETERMINATION OR POLICY.

Table 11—Accuracy of land use category assignments to individual subwatersheds.

Land use category	Mean	Standard Deviation	Minimum	Maximum
Congressionally reserved	82%	18%	38%	100%
Late-successional reserves	76%	18%	35%	100%
Matrix	77%	18%	35%	100%

Table 12--Percent difference in aerial photography-based road attributes compared to corporate GIS road attributes.

Attribute	Mean	Standard Deviation	Range
Upslope Road Density (road mi / upslope mi ²)	14%	13.5%	0 to 57%
Riparian Road Density (road mi / riparian mi ²)	18%	20%	-0 to 80%
Road Crossings (crossings / stream mi)	28%	38%	-0 to 150%

Table 13—Summary of vegetation attribute accuracy assessment metric ranges.

Attribute	R ¹	NRMSE ²
Quadratic mean diameter of all dominant and codominant trees (QMDA_DOM)	0.44 - 0.78	0.41 - 0.55
Quadratic mean diameter of all dominant and codominant conifers (QMDC_DOM)	0.42 - 0.72	0.45 - 0.7
Canopy cover of all live trees (CANCOV)	0.7 - 0.8	0.2 - 0.31
Canopy cover of all conifers (CANCOV_CON)	0.65 - 0.82	0.22 - 0.49

¹ Coefficient of correlation

² Normalized root mean squared error

Table 14—Field attributes evaluated in the Quality Assurance Program report. The environment column represents the percent of total variation in the data due to the differences between watersheds plus the differences between sites within watersheds. The Observer column represents the percent of the total variation in the data due to measurement by crews and unexplained error. The two columns do not total 100% because other terms related to time (differences through time) account for the remained of the total variation.

Attribute Type	Attribute	Environment	Observer
Channel morphology	Site length	75	17
	Average bankfull width	95	4
	Average bankfull depth	36	54
	Average Bankfull width:Depth	84	1
	Bedload gradient	96	2
Physical habitat	Stream channel sinuosity	5	87
	D50 (median particle size)	20	80
	D50 without Bedrock	97	0
	Pool Tail Crest fines	42	2
	Wood frequency	85	8
	Pool frequency	57	19
	Pool depth	45	50
Water chemistry	Dissolved oxygen	29	12
	Specific conductance	74	25
Biological	EPT ¹ richness	26	0
	Intolerant taxa richness	15	37
	Terrestrial amphibian simpsons index	55	16

¹ Ephemeroptera, Plecoptera, Trichoptera taxa

Figure numbering

Figures are numbered using Word's Insert Caption command. For compound figures (a, b, c), the first figure receives the caption and the sub-identifier (a) is put on a separate line and not in Caption style. Then References/Cross-reference is used to insert the labels for subsequent sub-figures, so they are tied to the numbering of the first sub-figure (and update automatically).




Figure 1—Timeline of the monitoring program development and implementation.

Aquatic Province Boundaries



Figure 2—Aquatic provinces used to assess watershed condition in the Northwest Forest Plan (NWFP) area. The NWFP area extends from the U.S.-Canada border to Point Reyes, California, and includes the eastern flank of the Cascade Mountain range and encompasses the range of

THE FINDINGS AND CONCLUSIONS IN THIS REPORT ARE IN PRESS AND SUBJECT TO CHANGE PRIOR TO FORMAL DISSEMINATION BY THE AGENCIES AND SHOULD NOT BE CONSTRUED TO REPRESENT AGENCY DETERMINATION OR POLICY.

the northern spotted owl. The NWFP area being evaluated includes USDA Forest Service, USDI Bureau of Land Management, and USDI National Park Service lands.




Figure 3—Land use allocations in the Northwest Forest Plan (NWFP) area.

THE FINDINGS AND CONCLUSIONS IN THIS REPORT ARE IN PRESS AND SUBJECT TO CHANGE PRIOR TO FORMAL DISSEMINATION BY THE AGENCIES AND SHOULD NOT BE CONSTRUED TO REPRESENT AGENCY DETERMINATION OR POLICY.



Figure 4—Program personnel from FS, BLM, NPS, Pacific Northwest Research Station, Washington Department of Fish and Wildlife, Washington Department of Ecology, Washington Forest Practices Board Cooperative Monitoring, Evaluation, and Research Committee, Environmental Protection Agency, National Oceanic and Atmospheric Administration Fisheries, Oregon Fish and Wildlife, and California Fish and Game provided technical expertise and local knowledge for decision-support model refinement during a series of aquatic province workshops. Specialists also provided feedback on how well the

THE FINDINGS AND CONCLUSIONS IN THIS REPORT ARE IN PRESS AND SUBJECT TO CHANGE PRIOR TO FORMAL DISSEMINATION BY THE AGENCIES AND SHOULD NOT BE CONSTRUED TO REPRESENT AGENCY DETERMINATION OR POLICY.

output from refined models matched up with their perspective of 'on-the-ground' conditions, so the models could be further fine-tuned.

DRAFT




Figure 5—Illustrative model diagrams showing how indicators are combined into overall condition scores (actual structures for each province detailed in appendix 4).




Figure 6—Example of randomly selected sample sites in a sixth-field watershed. The sampled stream reaches (red dots) were selected from 1:100,000 stream layers by using a generalized random stratified tessellation survey, a process that guarantees a spatially balanced sample (Stevens and Olsen 2004).

THE FINDINGS AND CONCLUSIONS IN THIS REPORT ARE IN PRESS AND SUBJECT TO CHANGE PRIOR TO FORMAL DISSEMINATION BY THE AGENCIES AND SHOULD NOT BE CONSTRUED TO REPRESENT AGENCY DETERMINATION OR POLICY.

DRAFT

THE FINDINGS AND CONCLUSIONS IN THIS REPORT ARE IN PRESS AND SUBJECT TO CHANGE PRIOR TO FORMAL DISSEMINATION BY THE AGENCIES AND SHOULD NOT BE CONSTRUED TO REPRESENT AGENCY DETERMINATION OR POLICY.



Figure 7—Bankfull width-to-depth ratios were calculated from cross-sectional profiles.



Figure 8—Large pieces of wood in the stream channel were counted in each sampled stream reach.



Figure 9—Fine sediment was measured at pool tail crests.

THE FINDINGS AND CONCLUSIONS IN THIS REPORT ARE IN PRESS AND SUBJECT TO CHANGE PRIOR TO FORMAL DISSEMINATION BY THE AGENCIES AND SHOULD NOT BE CONSTRUED TO REPRESENT AGENCY DETERMINATION OR POLICY.



THE FINDINGS AND CONCLUSIONS IN THIS REPORT ARE IN PRESS AND SUBJECT TO CHANGE PRIOR TO FORMAL DISSEMINATION BY THE AGENCIES AND SHOULD NOT BE CONSTRUED TO REPRESENT AGENCY DETERMINATION OR POLICY.

Figure 10—Naturally occurring landslides provide large wood and spawning gravels to rivers. However, excessive landslides caused by management activities can overwhelm river systems and negatively impact aquatic organisms.




Figure 11—Road density (miles per square mile of watershed) was used as an indicator of watershed condition. In this example of a sixth-field watershed (purple line), the black lines are roads, blue lines are streams, and red dots are road-stream crossings.



Figure 12—Watershed condition evaluations included metrics of average tree diameter and percent canopy cover in riparian and upslope areas.

THE FINDINGS AND CONCLUSIONS IN THIS REPORT ARE IN PRESS AND SUBJECT TO CHANGE PRIOR TO FORMAL DISSEMINATION BY THE AGENCIES AND SHOULD NOT BE CONSTRUED TO REPRESENT AGENCY DETERMINATION OR POLICY.



Figure 13—Disturbance from fire and harvest was used an alternative metric for assessing watershed condition.




Figure 14—Inchannel condition scores by status category for the 193 randomly selected watersheds in the Northwest Forest Plan area that have been sampled for inchannel attributes as of 2009.




Figure 15—Distribution of attribute scores for aquatic pools, substrate, wood, temperature, and aquatic invertebrates and for the 193 randomly selected watersheds in the Northwest Forest Plan area that were sampled for inchannel attributes as of 2009.

DRAFT

THE FINDINGS AND CONCLUSIONS IN THIS REPORT ARE IN PRESS AND SUBJECT TO CHANGE PRIOR TO FORMAL DISSEMINATION BY THE AGENCIES AND SHOULD NOT BE CONSTRUED TO REPRESENT AGENCY DETERMINATION OR POLICY.




Figure 16—Distribution of inchannel condition scores for the 193 randomly selected

THE FINDINGS AND CONCLUSIONS IN THIS REPORT ARE IN PRESS AND SUBJECT TO CHANGE PRIOR TO FORMAL DISSEMINATION BY THE AGENCIES AND SHOULD NOT BE CONSTRUED TO REPRESENT AGENCY DETERMINATION OR POLICY.

watersheds in the Northwest Forest Plan (NWFP) area that have been sampled for inchannel attributes as of 2009.

DRAFT




Figure 17—Distribution of inchannel scores by land use category for the 193 randomly selected watersheds in the Northwest Forest Plan area that were sampled for inchannel attributes as of 2009 (reserved = congressional reserves, LSR = late-successional reserves).





Figure 18—Distribution of inchannel attribute scores by land use category (reserved = congressional reserves, LSR = late-successional reserves).

a



DRK





Figure 18

b





DR





Figure 18

c



DRK





Figure 18

d





DR




Figure 18

d






Figure 18

e.






Figure 19—Watershed condition scores in the Northwest Forest Plan area by status category (2008).




Figure 20—Watershed condition status scores (2008) in the Northwest Forest Plan (NWFP) area.

THE FINDINGS AND CONCLUSIONS IN THIS REPORT ARE IN PRESS AND SUBJECT TO CHANGE PRIOR TO FORMAL DISSEMINATION BY THE AGENCIES AND SHOULD NOT BE CONSTRUED TO REPRESENT AGENCY DETERMINATION OR POLICY.

DRAFT

THE FINDINGS AND CONCLUSIONS IN THIS REPORT ARE IN PRESS AND SUBJECT TO CHANGE PRIOR TO FORMAL DISSEMINATION BY THE AGENCIES AND SHOULD NOT BE CONSTRUED TO REPRESENT AGENCY DETERMINATION OR POLICY.




Figure 21—Distribution of watershed condition status scores (2008) by land use allocation (reserved = congressional reserves, LSR = late-successional reserves).




Figure 22—Frequency of watershed status scores (2008) by land use allocation (reserved = congressional reserves, LSR = late-successional reserves). These curves show the data in a continuous manner, rather than by data bins (i.e., histograms).




Figure 23—Frequency of watershed status scores (2008) for key and nonkey watersheds. These curves show the data in a continuous manner, rather than by data bins (i.e., histograms).




Figure 24—Watershed condition status (2008) by land use allocation in the Northwest Forest Plan (NWFP) area.

THE FINDINGS AND CONCLUSIONS IN THIS REPORT ARE IN PRESS AND SUBJECT TO CHANGE PRIOR TO FORMAL DISSEMINATION BY THE AGENCIES AND SHOULD NOT BE CONSTRUED TO REPRESENT AGENCY DETERMINATION OR POLICY.

a

DRAFT

THE FINDINGS AND CONCLUSIONS IN THIS REPORT ARE IN PRESS AND SUBJECT TO CHANGE PRIOR TO FORMAL DISSEMINATION BY THE AGENCIES AND SHOULD NOT BE CONSTRUED TO REPRESENT AGENCY DETERMINATION OR POLICY.




Figure 24

b

THE FINDINGS AND CONCLUSIONS IN THIS REPORT ARE IN PRESS AND SUBJECT TO CHANGE PRIOR TO FORMAL DISSEMINATION BY THE AGENCIES AND SHOULD NOT BE CONSTRUED TO REPRESENT AGENCY DETERMINATION OR POLICY.

DRAFT

THE FINDINGS AND CONCLUSIONS IN THIS REPORT ARE IN PRESS AND SUBJECT TO CHANGE PRIOR TO FORMAL DISSEMINATION BY THE AGENCIES AND SHOULD NOT BE CONSTRUED TO REPRESENT AGENCY DETERMINATION OR POLICY.




Figure 24

c

THE FINDINGS AND CONCLUSIONS IN THIS REPORT ARE IN PRESS AND SUBJECT TO CHANGE PRIOR TO FORMAL DISSEMINATION BY THE AGENCIES AND SHOULD NOT BE CONSTRUED TO REPRESENT AGENCY DETERMINATION OR POLICY.

DRAFT

THE FINDINGS AND CONCLUSIONS IN THIS REPORT ARE IN PRESS AND SUBJECT TO CHANGE PRIOR TO FORMAL DISSEMINATION BY THE AGENCIES AND SHOULD NOT BE CONSTRUED TO REPRESENT AGENCY DETERMINATION OR POLICY.




Figure 25—Watershed condition status (2008) for key and nonkey watersheds in the Northwest Forest Plan (NWFP) area.

THE FINDINGS AND CONCLUSIONS IN THIS REPORT ARE IN PRESS AND SUBJECT TO CHANGE PRIOR TO FORMAL DISSEMINATION BY THE AGENCIES AND SHOULD NOT BE CONSTRUED TO REPRESENT AGENCY DETERMINATION OR POLICY.

a

DRAFT

THE FINDINGS AND CONCLUSIONS IN THIS REPORT ARE IN PRESS AND SUBJECT TO CHANGE PRIOR TO FORMAL DISSEMINATION BY THE AGENCIES AND SHOULD NOT BE CONSTRUED TO REPRESENT AGENCY DETERMINATION OR POLICY.




Figure 25

b

THE FINDINGS AND CONCLUSIONS IN THIS REPORT ARE IN PRESS AND SUBJECT TO CHANGE PRIOR TO FORMAL DISSEMINATION BY THE AGENCIES AND SHOULD NOT BE CONSTRUED TO REPRESENT AGENCY DETERMINATION OR POLICY.

DRAFT

THE FINDINGS AND CONCLUSIONS IN THIS REPORT ARE IN PRESS AND SUBJECT TO CHANGE PRIOR TO FORMAL DISSEMINATION BY THE AGENCIES AND SHOULD NOT BE CONSTRUED TO REPRESENT AGENCY DETERMINATION OR POLICY.




Figure 26—Frequency of watershed condition scores in 1994 and 2008. These curves show the data in a continuous manner, rather than by data bins (i.e., histograms).




Figure 27—Distribution of changes in watershed condition scores between 1994 and 2008.




Figure 28—Watershed condition scores by status category in 2008.




Figure 29—Number of watersheds with category scores from 1994 and 2008. The bubble sizes range from few watersheds (smallest bubbles) to many watersheds (largest watersheds).




Figure 30—Change in watershed condition score 1994 to 2008.






Figure 31—Distribution of changes in watershed scores from 1994 to 2008 by land use allocation.




Figure 32—Frequency of 1994 and 2008 watershed scores by land use allocation. These curves show the data in a continuous manner, rather than by data bins (i.e., histograms) (reserved = congressional reserves, LSR = late-successional reserves).





Figure 33—Distribution of changes in watershed condition scores between 1994 and 2008 for key and nonkey watersheds.




Figure 34—Frequency of 1994 and 2008 watershed condition scores for key and nonkey watersheds. These curves show the data in a continuous manner, rather than by data bins (i.e., histograms).




Figure 35—Watershed road density (mi/mi²) in 1994 and 2008.




Figure 36—Changes in watershed road density (mi/mi²) from 1994 to 2008.




Figure 37—Road density by land category, 2008 (reserved = congressional reserves, LSR = late-successional reserves).




Figure 38—Change in road density (mi/mi²) by land use category, 1994 to 2008.

A

DRAFT




Figure 38

b

DRAFT




Figure 38

c

DRAFT




Figure 38

d

DRAFT




Figure 38

e

DRAFT




Figure 39—Percentage of watershed area in late-successional old-growth vegetation (LSOG; conifer quadratic mean diameter \geq 20 inches and canopy cover \geq 10 percent) in 1994 and 2008.




Figure 40—Change in percentage of watershed area in late-successional old-growth vegetation (LSOG; conifer quadratic mean diameter > 20 inches) from 1994 to 2008.

DRAFT




Figure 41—Percent of watershed area in late-successional (LSOG; conifer quadratic mean diameter at breast height ≥ 20 inches and canopy cover ≥ 10 percent) by land use allocation, 2008 (Reserved = Congressional reserves, LSR = late-successional reserves).




Figure 42—Change in percentage of watershed in late-successional old-growth (LSOG; diameter at breast height \geq 20 inches and canopy cover \geq 10 percent) from 1994 to 2008 by land category.

a




Figure 42

b

DRAFT




Figure 42

c

DRAFT




Figure 42

d

DRAFT




Figure 42

e

DRAFT




Figure 43—Average percentage of watershed area affected by harvest and fire disturbances 1985 to 2008.




Figure 44—Average yearly pre- and post-Northwest Forest Plan (NWFP) fire and harvest disturbance levels by land use allocation, and for key and nonkey watersheds (Reserved = congressionally reserved, LSR = late-successional reserve).

DRAFT




Figure 45—Watershed versus inchannel current status scores. These curves show the data in a continuous manner, rather than by data bins (i.e., histograms).




Figure 46—Upslope/riparian versus inchannel current status scores for 193 watersheds with both data types available.




Figure 47—Federal lands in the headwaters of the Clearwater River watershed (Mount Baker-Snoqualmie National Forest) are within a wilderness. Nonfederal lands in the upper half of the same watershed are managed for timber production, and that area is heavily roaded. Including the nonfederal lands would misrepresent the watershed score if the watershed is classified as a federal watershed.




Figure 48—High Cascades province evaluation model structures for watershed and inchannel conditions (EPT = Percentage of taxa in *ephemeroptera*, *plecoptera*, and *triptera*).




Figure 49—Klamath-Siskiyou and Franciscan province evaluation model structures for watershed and inchannel conditions (EPT = Percentage of taxa in *ephemeroptera*, *plecoptera*, and *tricoptera*; D50 = mean particle size).




Figure 50—North Cascades province evaluation model structures for watershed and inchannel conditions (EPT = Percentage of taxa in *ephemeroptera*, *plecoptera*, and *tricoptera*; D50 = mean particle size).






Figure 51—Olympic Peninsula province evaluation model structures for watershed and inchannel conditions (EPT = Percentage of taxa in *ephemeroptera*, *plecoptera*, and *tricoptera*; D50 = mean particle size).

Oregon Coast Province

Watershed condition
(Upslope and riparian)



Inchannel condition



KEY
(Relative weights in parentheses)
[Context switches in brackets]

Figure 52—Oregon Coast province evaluation model structures for watershed and inchannel conditions (EPT = Percentage of taxa in *ephemeroptera*, *plecoptera*, and *tricoptera*).




Figure 53—West Cascades province evaluation model structures for watershed and inchannel conditions (EPT = Percentage of taxa in *ephemeroptera*, *plecoptera*, and *tricoptera*; D50 = mean particle size).

High/Western Cascades Stream Scores 2009






Figure 54-Distribution of inchannel condition scores for the randomly selected watersheds in the High Cascade and West Cascade Provinces in the Northwest Forest Plan (NWFP) area that have been sampled for inchannel attributes as of 2009.

DRAFT




Figure 55-Watershed condition status scores (2008) for the High Cascade and West Cascade Provinces in the Northwest Forest Plan (NWFP) area.




Figure 56- Watershed condition trend scores (1994 to 2008) for the High Cascade and West Cascade Provinces in the Northwest Forest Plan (NWFP) area.

THE FINDINGS AND CONCLUSIONS IN THIS REPORT ARE IN PRESS AND SUBJECT TO CHANGE PRIOR TO FORMAL DISSEMINATION BY THE AGENCIES AND SHOULD NOT BE CONSTRUED TO REPRESENT AGENCY DETERMINATION OR POLICY.




Figure 57-Distribution of inchannel condition scores for the randomly selected watersheds in the Klamath/Siskiyou and Franciscan Provinces in the Northwest Forest Plan (NWFP) area that have been sampled for inchannel attributes as of 2009.

THE FINDINGS AND CONCLUSIONS IN THIS REPORT ARE IN PRESS AND SUBJECT TO CHANGE PRIOR TO FORMAL DISSEMINATION BY THE AGENCIES AND SHOULD NOT BE CONSTRUED TO REPRESENT AGENCY DETERMINATION OR POLICY.




Figure 58-Watershed condition status scores (2008) for the Klamath/Siskiyou and Franciscan Provinces in the Northwest Forest Plan (NWFP) area.

THE FINDINGS AND CONCLUSIONS IN THIS REPORT ARE IN PRESS AND SUBJECT TO CHANGE PRIOR TO FORMAL DISSEMINATION BY THE AGENCIES AND SHOULD NOT BE CONSTRUED TO REPRESENT AGENCY DETERMINATION OR POLICY.




Figure 59-Watershed condition trend scores (1994 to 2008) for the Klamath/Siskiyou and Franciscan Provinces in the Northwest Forest Plan (NWFP) area.

THE FINDINGS AND CONCLUSIONS IN THIS REPORT ARE IN PRESS AND SUBJECT TO CHANGE PRIOR TO FORMAL DISSEMINATION BY THE AGENCIES AND SHOULD NOT BE CONSTRUED TO REPRESENT AGENCY DETERMINATION OR POLICY.




Figure 60-Distribution of inchannel condition scores for the randomly selected watersheds in

THE FINDINGS AND CONCLUSIONS IN THIS REPORT ARE IN PRESS AND SUBJECT TO CHANGE PRIOR TO FORMAL DISSEMINATION BY THE AGENCIES AND SHOULD NOT BE CONSTRUED TO REPRESENT AGENCY DETERMINATION OR POLICY.

the North Cascades Province in the Northwest Forest Plan (NWFP) area that have been sampled for inchannel attributes as of 2009.

DRAFT

North Cascades Watershed Condition 2008





Figure 61-Watershed condition status scores (2008) for the North Cascades Province in the Northwest Forest Plan (NWFP) area.

THE FINDINGS AND CONCLUSIONS IN THIS REPORT ARE IN PRESS AND SUBJECT TO CHANGE PRIOR TO FORMAL DISSEMINATION BY THE AGENCIES AND SHOULD NOT BE CONSTRUED TO REPRESENT AGENCY DETERMINATION OR POLICY.

North Cascades Watershed Condition 1994 to 2008






Figure 62-Watershed condition trend scores (1994 to 2008) for the North Cascades Province in the Northwest Forest Plan (NWFP) area.

DRAFT




Figure 63-Distribution of inchannel condition scores for the randomly selected watersheds in the Olympic Peninsula Province in the Northwest Forest Plan (NWFP) area that have been sampled for inchannel attributes as of 2009.

THE FINDINGS AND CONCLUSIONS IN THIS REPORT ARE IN PRESS AND SUBJECT TO CHANGE PRIOR TO FORMAL DISSEMINATION BY THE AGENCIES AND SHOULD NOT BE CONSTRUED TO REPRESENT AGENCY DETERMINATION OR POLICY.




Figure 64-Watershed condition status scores (2008) for the Olympic Peninsula Province in the Northwest Forest Plan (NWFP) area.

THE FINDINGS AND CONCLUSIONS IN THIS REPORT ARE IN PRESS AND SUBJECT TO CHANGE PRIOR TO FORMAL DISSEMINATION BY THE AGENCIES AND SHOULD NOT BE CONSTRUED TO REPRESENT AGENCY DETERMINATION OR POLICY.




Figure 65-Watershed condition trend scores (1994 to 2008) for the Olympic Peninsula Province in the Northwest Forest Plan (NWFP) area.

THE FINDINGS AND CONCLUSIONS IN THIS REPORT ARE IN PRESS AND SUBJECT TO CHANGE PRIOR TO FORMAL DISSEMINATION BY THE AGENCIES AND SHOULD NOT BE CONSTRUED TO REPRESENT AGENCY DETERMINATION OR POLICY.




Figure 66-Distribution of inchannel condition scores for the randomly selected watersheds in

THE FINDINGS AND CONCLUSIONS IN THIS REPORT ARE IN PRESS AND SUBJECT TO CHANGE PRIOR TO FORMAL DISSEMINATION BY THE AGENCIES AND SHOULD NOT BE CONSTRUED TO REPRESENT AGENCY DETERMINATION OR POLICY.

the Oregon Coast Province in the Northwest Forest Plan (NWFP) area that have been sampled for inchannel attributes as of 2009.

DRAFT

THE FINDINGS AND CONCLUSIONS IN THIS REPORT ARE IN PRESS AND SUBJECT TO CHANGE PRIOR TO FORMAL DISSEMINATION BY THE AGENCIES AND SHOULD NOT BE CONSTRUED TO REPRESENT AGENCY DETERMINATION OR POLICY.




Figure 67-Watershed condition status scores (2008) for the Oregon Coast Province in the Northwest Forest Plan (NWFP) area.

THE FINDINGS AND CONCLUSIONS IN THIS REPORT ARE IN PRESS AND SUBJECT TO CHANGE PRIOR TO FORMAL DISSEMINATION BY THE AGENCIES AND SHOULD NOT BE CONSTRUED TO REPRESENT AGENCY DETERMINATION OR POLICY.




Figure 68-Watershed condition trend scores (1994 to 2008) for the Oregon Coast Province in the Northwest Forest Plan (NWFP) area.

THE FINDINGS AND CONCLUSIONS IN THIS REPORT ARE IN PRESS AND SUBJECT TO CHANGE PRIOR TO FORMAL DISSEMINATION BY THE AGENCIES AND SHOULD NOT BE CONSTRUED TO REPRESENT AGENCY DETERMINATION OR POLICY.

Overview of the Watershed Evaluation Modeling Process

Our watershed assessment process uses the Ecosystem Management Decision Support system (EMDS) to conduct consistent evaluations that integrate diverse kinds of data, such as vegetation and roads. EMDS is not a cause-and-effect statistical model, rather it is a framework for integrating diverse sources of information (from statistical relationships to expert opinion) consistently across time and space.

Advantages of using such a decision-support model include:

- All aspects of the analysis process are shown; therefore, it is easy to explain to customers.
- Models can be developed to assess ecological condition at any spatial or temporal scale.
- As we learn more about how watersheds function, models can be refined and rerun on data from earlier periods to correct deficiencies.


The simplified model structure below illustrates how each watershed was scored using the EMDS process. The actual model structures we used are presented in appendix 4.

Step 1: Each attribute (i.e., indicator) is scored to a common scale between +1.0 (for “good”) and -1.0 (for “poor”) by comparing it to a set of evaluation criteria.

Criteria are based on published literature, field data, and professional judgment. Usually one evaluation criterion value is selected for the +1 threshold, another for the -1 threshold, and intermediate values are scored based on a linear interpolation between these thresholds. Evaluation function examples are shown for the road components below.

Step 2: The evaluation scores for each of the attributes are aggregated together for each general model component (e.g., Roads) by using user-defined rules. Selection of the rules is based on experts’ knowledge of the system and ecological processes. Rules can produce an aggregated score weighted toward the resource with either the highest or lowest attribute score, or use the average of scores (as shown in this example by “AVE”). An aggregated score can also be based on the weighted or unweighted average of the indicator evaluation scores, e.g., as shown in step 3.

Step 3: The component scores are further aggregated based on the model structure. In this model structure, the watershed condition is determined by using the weighted average of roads (60 percent) and vegetation (40 percent) scores. The watershed condition score will always range from -1.0 to +1.0.



Example of a simplified decision-support model. In step 1, individual attributes are evaluated by using evaluation criteria. In steps 2 and 3, the evaluation scores of the attributes are aggregated to determine the overall watershed condition score. AVE = average of scores. Revised from Gallo et al. 2005.

Macroinvertebrate Metric

An evaluation of aquatic macroinvertebrates was added into the decision-support model (DSM) for this report. The diversity and environmental sensitivity of macroinvertebrates make them useful indicators of stream condition, and they respond predictably to anthropogenic disturbances on the landscape or in the stream (Karr and Chu 1999). To determine which macroinvertebrate metrics to include in the DSM, we engaged the assistance of Alan Herlihy, an Oregon State University scientist who had used data from the Environmental Protection Agency's Environmental Monitoring and Assessment Program (EMAP) to develop indices of biotic integrity for macroinvertebrates. We analyzed Aquatic and Riparian Effectiveness Monitoring Program data and EMAP data collected within the Northwest Forest Plan area independently to select macroinvertebrate metrics that are sensitive to management in the Plan area and eliminate redundant metrics. The EMAP data were included to increase sample size and because the data set includes forested lands that are not federally owned, which increased the range of environmental conditions.

Three metrics were selected for inclusion in the DSM. The first attribute, proportion of the macroinvertebrate taxa in the orders of Ephemeroptera (mayflies), Tricoptera (caddisflies), and Plecoptera (stoneflies), is a community composition index that focuses on species known to exist in cold, high-quality waters (Merritt and Cummins 1996). The second metric, the proportion of the macroinvertebrate taxa that are intolerant, can be used to detect nutrient enrichment, high sediment load, high water temperature, and organic pollution (Hilsenhoff 1988). The final measure, proportion of the macroinvertebrate taxa that are climbers, is an indicator of habitat. These taxa tend to live in structurally complex habitats, such as those with high-quality spawning gravel or cover for juvenile fish (Karr and Chu 1999).

Determination of evaluation criteria for each metric was conducted by distinguishing sample locations that were in minimally disturbed or "reference" condition



Steve Lamigan

Three metrics were used in the decision-support model for macroinvertebrate data: (1) a community composition index that focuses on species known to exist in cold, high-quality waters; (2) the proportion of the macroinvertebrate taxa that are intolerant, to detect nutrient enrichment, high sediment load, high water temperature, and organic pollution; and (3) the proportion of the macroinvertebrate taxa that are climbers.

from those in poor condition, then looking at the range of metric values to distinguish the two types of locations. The median data value of the "reference" sites was designated as the +1 evaluation criterion and the median value of the poor sites was designated as the -1 criterion.

Amphibian Metric

An evaluation of terrestrial and aquatic amphibians was added into the decision support model (DSM) for this report. Amphibians have characteristics that make them useful for monitoring watershed health including their association with riparian areas (Olson et al. 2007), their sensitivity to environmental stressors (Blaustein et al. 1994), and they are inexpensive to sample (relative to many other field attributes). Linkages between inchannel impacts of management practices and amphibians in aquatic life stages (*Dicamptodon* spp and larval *Ascaphus truei*) have been demonstrated in the literature (Olson et al. 2007).

The amphibian data came from two survey protocols: time-constrained searches for terrestrial amphibians done in a splash zone (within 2 m of the wetted edge), and single-pass electrofishing, without block nets, done throughout the survey reach.¹ Because the two different protocols were not optimal in their ability to capture animals for a variety of reasons (including time of




Steve Lamigan

Aquatic and terrestrial amphibians were included in our assessment of stream reach condition.

year and moisture conditions), the lower range of the DSM score was truncated to 0. This resulted in giving positive credit to the watershed score if animals are captured, while not penalizing the score when we failed to find any amphibians.

¹ Aquatic amphibian data was available from 2002 – 2007. Electrofishing was discontinued after 2007.



A Simpson's Index of species richness was used to convert the captures of each of the aquatic amphibians and the terrestrial amphibians to a 0 to +1 scale. There are three possible outcomes for generating a decision-support model (DSM) score for each branch of the amphibians model: (1) If no aquatic amphibians are found at a site, then the aquatic amphibian score is zero. (2) If only one species of aquatic amphibian is captured (approximately 58 percent of the sites with aquatic amphibian captures and 41 percent of the sites with terrestrial amphibian captures had only one species present) or the Simpson's Index value is <0.33, then the DSM score is arbitrarily changed to a value of 0.33. (3) If more than one species of aquatic amphibian is present and the Simpson's Index value is ≥ 0.33 , then the Simpson's Index value is used as the DSM score². The two DSM scores—aquatic and terrestrial—were averaged with a weight of 0.75 and 0.25³, respectively for the overall site amphibian score.

² Olson, D.H. 2010. Personal communication. Research Ecologist. Pacific Northwest Research Station, 3200 SW Research Way, Corvallis, OR 97331.
³ Ibid

Distribution Functions

This report makes use of three types of distribution graphs: box plots (A), histograms (B), and frequency of occurrence graphs (C). Box plots break the data into four groups each representing a quarter of the range of data values or observations. The box represents the middle 50 percent of the data called the second and third quartiles, and the lines outside the box represent the first and fourth quartiles. The line inside the box represents the median value. In the plots showing inchannel data, which are samples of all watersheds, the shaded area represents possible variation in the median value. The median does not vary for watershed condition scores because the scores are based on upslope and riparian data for all watersheds, i.e., a census.


Histograms and frequency of occurrence graphs represent the distribution of a data set in a similar way. The x-axis represents the range of values within the data set, and the y-axis represents the number of observations that occur at different places within the range. In a histogram, the total range of values is segmented into smaller groups called bins. The x-axis represents the bins and the y-axis represents how many observations fall into each bin. Because the data are grouped together, the size of the bin affects the shape of the histogram. Smaller bins provide a closer approximation of the shape of the distribution for continuous data such as watershed and attribute scores.

Frequency of occurrence graphs also display the range of the data on the x-axis. However, the y-axis represents the relative number of watersheds that are in the vicinity of a vertical line to the x-axis. The relative amount is determined using bins as in the histograms. Instead of counting the occurrences of observations in each bin, the count is compared to the rest of the observations in a mathematical interpolation function. Interpolation infers observations in places where no data are present, making it possible to display a continuous representation of the data. In the case of density graphs, this creates a line describing the shape of the data where higher counts of observations result in higher relative amount. Because the line is continuous, it provides insight into shape of the distribution that can not be derived from a histogram.


To interpret a frequency of occurrence graph, consider the height, slope, and features of the line. The higher

on the y-axis the more observations around the x-axis value, steeper line slopes identify larger increases or decreases in the number of observations around the x-axis value, and peaks represent high concentrations of observations. With overlaid graphs, at any point along the x-axis the top line has more observations than the bottom line. When frequency of occurrence graphs are used to represent the same attribute at different times, shifts along the x-axis of features such as peaks implies a concentration of observations has increased or decreased its highest frequency x-axis value.


A - box plot



B - histogram



C - frequency of occurrence graph



Modeling Inchannel Variables from Landscape Conditions

Aquatic and Riparian Effectiveness Monitoring Program (AREMP) staff contracted with Kelly Burnett (Pacific Northwest Research Station [PNW]), Ken Vance-Borland (Oregon State University [OSU]), Rebecca Flitcroft (PNW), and Lisa Ganio (OSU) to develop statistical models to explore relationships between inchannel variables and landscape conditions (geographic information system and satellite imagery data).

Numerous landscape conditions were explored as independent variables. One of these, contributing area (km²), was determined for the entire catchment upstream of each sampled reach. All other landscape conditions were summarized in 100-m buffers and 2-km catchments for each sampled reach. For modeling each in-channel variable, a core set of six landscape conditions (table 4) was considered. Contributing area (km²), reach gradient, and mean annual precipitation (cm) were selected as influencing stream power and thus numerous biophysical characteristics of fluvial systems (e.g., Knighton 1999, Stanford 1998). Road density (km/km²), percentage of area harvested 1972–2002, and % area with >65 percent hardwood cover were selected as indicators of riparian and stream disturbance

(e.g., Lee et al. 1997, Pabst and Spies 1999). Inchannel data used included pool frequency (number of pools per meter), frequency of key (0.6 m x 15 m) large wood pieces (number per m), frequency of all large (0.3 m x 3 m) wood pieces (number per m), and aquatic macroinvertebrate species richness (number of Ephemeroptera species in the reach).

Statistical models indicated that inchannel variables were generally more strongly related to landscape conditions describing topography, lithology, or climate than those considered to be sensitive to human activities. The relatively weak relationship observed between inchannel variables and landscape conditions sensitive to human activities may stem from several sources, including use of probability sampled data, among-reach variability that was relatively low for land use but high for characteristics unrelated to management, and a legacy of splash-damming effects in streams.

Although the current AREMP probability sample design was considered useful for collecting inchannel data to assess status and monitor trends in streams, a modified sampling design may be more conducive for developing models to explain these trends or to predict inchannel data in nonsampled watersheds from GIS and remote-sensing data.

Table 4—Landscape conditions that were considered in developing final models for each inchannel variable.

	Landscape Condition	Pool frequency	Key wood frequency	All large wood frequency	Macroinvert. family richness	Ephemeroptera species richness
Core set	Contributing area (km ²)	X	X	X	X	X
	Reach gradient (%)	X	X	X	X	X
	Mean annual precipitation (cm)	X	X	X	X	X
	Road density (km/km ²)	X	X	X	X	X
	Percentage of (%) area harvested 1972 - 2002	X	X	X	X	X
	% area with >65% hardwood cover	X	X	X	X	X
Disurbance	% area with urban cover	X				X
	Average quadratic mean diameter (QMD) of coniferous trees (cm)	X				X
	% area with conifer QMD 10.2-25.4 cm				X	
	% area with conifer QMD > 50.8 cm		X	X		
Topography	% area with hillslope 3-6%					X
	% area with hillslope 20-44%	X	X	X	X	
	Mean elevation (m)	X			X	
	% area with elevation <198.1 m					X
	% area with elevation <609.6 m		X	X		
Lithology	% area with unconsolidated deposits				X	X
	% area with weak rock		X	X	X	

Fish Passage at Culverts


When culvert fish-passage data become available, decision-support models can reflect improved conditions as fish passage barriers are removed.

Forest Service—An initial culvert assessment was done by Forests in 2002-2003 that covered almost all streams with the range of anadromy and 80 percent of the streams with resident fishes.¹ However, culvert location data are known to have spatial errors that need to be corrected. An effort is underway by the Forest Service (FS) to compile and update all their culvert location data for where fish are present. Documentation of actions to correct fish-passage problems are also incomplete; the FS corporate database only tracks work funded out of the regional office (about 75 percent of the culvert-related projects). Additional culvert improvement/removal projects funded at the forest level, e.g., Knutson-Vandenberg Act funds,

¹D.Heller, 2010. Personal communication, Regional Fish Biologist, retired, USDA Forest Service Pacific Northwest Region, 333 SW First Ave., Portland, OR 97204.

Burned Area Emergency Response, partnership funding, Emergency Relief for Federally Owned Roads, and Highway Trust Funds used for county and state roads are not being tracked.

Bureau of Land Management— Bureau of Land Management (BLM) culvert data varies in its quality, consistency and completeness. Working in concert with BLM, Oregon Department of Fish and Wildlife has converted a substantial quantity of BLM culvert/barrier data into the Oregon Fish Passage Barrier Data Standard. As of April 2009, a total of 3,981 barrier records from five western Oregon BLM districts were “standardized,” with several thousand still to be evaluated. Culverts constitute the majority of the barriers that will be tracked by the data set, but dams, diversions, tidegates, weirs, falls, and cascades/gradient barriers are also included. Data collected from BLM units starting in 1999 for aquatic restoration project data, including culvert removal, replacement, and upgrade, are stored in a corporate geographic information system data warehouse.



All other things being equal, these two watersheds would have received the same watershed condition score, despite the fact that the watershed on the left is almost entirely blocked to fish passage. When culvert locations and culvert fish passage assessment data become available we intend to use the amount of fish habitat available to fish as part of our watershed condition assessment.

Road Data Challenges

Roads are a major part of our evaluation models, however there are a number of issues with available road data.



Not all roads on federal lands are digitized in existing geographic information system (GIS) layers—Figure A is a digital orthoquad with an overlay of the existing Willamette National Forest roads layer. Road locations are not exact and some roads seen in the orthoquad are not included in the Forest Service (FS) roads layer. This is because the FS does not track “nonsystem” roads, even though they may affect hydrological processes. These are usually smaller spur roads built for timber sales. Gallo et al. 2005 found that the Bureau of Land Management (BLM) and FS underrepresented roads by 10 and 15 percent, respectively. Roads on nonfederal lands were underrepresented by 37 percent.




Figure B shows the same area with a surface derived from LiDAR (Light Detection and Ranging). LIDAR is a remote sensing data collection system that collects topographic information using aircraft-mounted lasers. A road layer derived from LIDAR imagery has more accurate road locations and includes roads that are not part of existing GIS road layers. Specifically, roads not included in the FS roads layer, but visible on the orthoquad and roads that were “hidden” under the canopy show up in LIDAR imagery. Complete LIDAR coverage of the Northwest Forest Plan area is not available now, but is expected to become available as more areas are flown and agencies share their data. However, LIDAR can provide only road locations and gradient. The BLM and FS road databases will still be needed to provide information about the roads, such as maintenance level and surface type.



BLM and FS have separate road layers—We derived our current road layer by putting together the BLM ground transportation layer with the FS Infra travel route transportation data. However, because the two agencies manage their road information in different data structures and have different road attributes, the data did not go together easily.

We clipped each agency’s data to their ownership to avoid overlap when combining road layers (Fig C).

However, this potentially caused roads or their attributes to be clipped off if they were not contained in the databases of both agencies. Evaluating road attributes that could affect watershed condition, such as surface type and maintenance levels, will continue to be very challenging unless the FS and BLM manage their road data cooperatively.

Better Tracking of Road Improvements Is Needed—We currently evaluate the effect of roads on watershed condition by using a rather simplistic approach because of the lack of corporate road data. The only road information available is whether a road is present or it has been decommissioned (and is therefore assumed to be benign with respect to sediment delivery to streams). Therefore, decommissioned roads are the only “road improvements” tracked in federal agency databases. However, the term “decommissioned” can be applied to a road that has been closed by a gate or a tank trap, or to a road that has been obliterated. Because other road condition improvements (e.g., outsloping, water bars, drivable fords, hardening surfaces) can also reduce sediment delivery, we recommend that additional road attributes should be tracked in federal agency databases. Availability of these data will allow future decision-support models to better reflect the effects of roads on watersheds and allow managers to take credit for those improvements.

Support to Local Units



Steve Lanigan

Protocol comparison test—Aquatic and Riparian Effectiveness Monitoring Program (AREMP) staff helped initiate and coordinate an interagency side-by-side protocol test in the John Day Basin (eastern-central Oregon). The goal of this effort was to assess the performance and compatibility of measurements obtained from seven monitoring groups that all use different monitoring protocols to assess stream habitat throughout the Pacific Northwest. The following were examined: (1) consistency of measurements within a monitoring group, (2) the ability of each monitoring protocol to detect environmental heterogeneity, (3) the compatibility of measurements between monitoring groups, and (4) the relationship of measurements to more intensive stream measurements that may better describe the true character of stream habitat. Understanding how the results of different monitoring programs are related to each other may foster improvement in the quality of stream habitat data, increase the sharing of data across monitoring groups, and increase statistical power to detect environmental trends (Roper et al. 2010).




Steve Lanigan

Aquatic invasive species surveys—Aquatic and Riparian Effectiveness Monitoring Program (AREMP) staff participated in a pilot regional survey effort to locate aquatic invasive species on federal lands. Training and field protocols we developed with Oregon State University Sea Grant College Program personnel were used to survey for 23 aquatic plants and animals identified as threats to northwest watersheds. We are continuing to work with regional invasive species coordinators to develop a process for alerting agency managers when aquatic invasives occur on lands they manage, so appropriate control and eradication efforts can be made (Andersen and Lanigan 2009).

Support to Local Units

Forest plan revisions—Program personnel worked with specialists on the Okanogan-Wenatchee and Colville National Forests, the Forest Service (FS) Pacific Northwest Regional Office, and forests in the Blue Mountains (Umatilla, Malheur, and Wallowa-Whitman) to apply decision-support models in their forest plan revisions. The Aquatic and Riparian Effectiveness Monitoring Program (AREMP) watershed condition model is being used by these forests as part of the key watershed designation process and to evaluate the contribution of the FS.

Assessment of temperature problems—We initiated and provided a summary of hydrograph temperature data we collected to BLM and FS regional water quality coordinators. These data will be shared with the Oregon Department of Environmental Quality and Washington Department of Ecology for assessment of temperature total maximum daily load (TMDL).



Restoration project survey support—We assisted the Fisheries and Hydrology staff of the Roseburg Bureau of Land Management district on a project to map existing channel configuration at the beginning of a restoration project so that changes could be measured through time. Six sites (in two watersheds) totaling approximately 6,000 feet were intensively mapped to document the existing channel and habitat features. Mapped habitat features included different types of substrate bar classifications (distinguished from bed load material), wood (both natural and placed), exposed bedrock sheets, and information about the existing pools. We resurveyed the same sites after the project was in place to detect differences in substrate as the result of a winter flood.



Bull trout reintroduction—The Mount Hood National Forest is working with the U.S. Fish and Wildlife Service, Oregon Department of Fish and Wildlife, and researchers from the U.S. Geological Survey to reintroduce bull trout into the Clackamas River. Aquatic and Riparian Effectiveness Monitoring Program personnel consulted on the project and built a decision-support model to document the decision process used to determine the suitability of an area for reintroduction and the likelihood of success. The model can be used for reintroduction of any species in any ecosystem. It evaluates data related to historical occupation of the species in the ecosystem, likelihood that the species is still present, natural recolonization potential, potential of habitat to support the reintroduction, threats, and impacts to donor stocks (Dunham and Gallo 2008).

Cummins and Tenmile Watershed, Siuslaw National Forest, Oregon

Land acquisition and conservation easement tools are being used for restoring watersheds and recovering species


—Land acquisitions and conservation easements are a major component of the restoration strategy in the Cummins/Tenmile watershed. Although most of the watershed is managed by the Forest Service, a large number of the major fish-bearing streams were located on private land prior to implementation of the land acquisition program.

Working with the Trust for Public Lands and local landowners, the Forest Service, Audubon Society of Portland, and The Nature Conservancy have acquired 1,900 acres and 10 miles of critical habitat for threatened coho salmon in the Cummins/Tenmile

watershed. In addition, the Tenmile Conservation Program worked with the McKenzie River Trust and local landowners to complete a 500-acre conservation easement package to protect habitat for multiple species dependent on older forest habitat, including an additional 2 miles of coho salmon streams. Funding for the land acquisitions and easements came from the Federal Land and Water Conservation Fund, Landowner Incentive Program, Oregon Watershed Enhancement Board, and Oregon State Parks.

The restoration strategy was recently extended into the Pacific Ocean adjacent to the Cummins/Tenmile watershed with a proposal to make a marine reserve in Oregon's state waters. These ongoing efforts combined with the Aquatic Conservation Strategy in the Northwest Forest Plan have protected ecologically significant habitat, and they have enabled implementation of a variety of restoration activities that work with natural processes to recover depleted and Federally Endangered Species Act listed species.

Contact Jack Sleeper (jsleeper@fs.fed.us) for more information




Elkhorn Creek Watershed, Salem BLM District, Oregon


Natural disturbance and past management affected coho habitat—The Elkhorn Creek watershed is the most productive salmon habitat in the Trask River drainage. However, these streams were lacking large wood due in part to a series of large-scale fires collectively called the Tillamook Burn and the extensive salvage logging that followed in the late 1940s through the early 1960s. Many roads and skid trails were built for post-fire logging that are still affecting hydrologic processes. Historic removal of large wood from streams also occurred on public and privately managed lands. The lack of large wood negatively affected many inchannel processes including: creation of complex pools, sediment and nutrient storage, spawning gravel sorting and retention, and providing cover for fish during high winter stream flows. Pre-project Rapid Bio-Assessment (RBA) surveys had shown Elkhorn and Cruiser Creeks to have high summer rearing capacity but minimal capability to support juvenile salmonids during high winter flow regimes.

A diverse partnership is restoring the watershed—Tillamook Bureau of Land Management, Oregon Dept of Fish and Wildlife, Oregon Dept of Forestry, U.S. Fish and Wildlife Service, Tillamook Estuaries Partnership, Tillamook Bay Watershed Council, Weyerhaeuser LLC, Tillamook Futures Council, Nestucca Valley High School, and Bio-Surveys LLC have all worked together to enhance summer rearing and over-winter habitat on a watershed scale by placing large wood and boulder structures in Cruiser Creek and Elkhorn Creek to increase stream complexity, improve pool/riffle ratios, and retain more of the quality spawning gravels that were being washed out of these streams. Other restoration actions included riparian plantings, replacing culverts, and obliterating and decommissioning riparian roads.

Fish numbers are up!—The treated reaches have shown dramatic increases in complex pools, and retention of spawning gravels and increased stream complexity. RBA surveys show that coho over-winter survival rates have gone from ~ 5 percent pre-project for both Cruiser and Elkhorn Creeks, to a post-project high of 17.4 percent in Cruiser and 25 percent in Elkhorn! This represents significant production increases for Oregon Coast coho from these watersheds.



Russ Chapman photos



B - Felled alders and newly placed wood worked together to catch coarse woody debris coming downstream and improved channel functions and complexity almost immediately.

A - An old remnant channel (top photo) became active again with multiple braided flood-plain channels after wood was added (bottom photo).

Contact Russ Chapman (rchapman@blm.gov) for more information.

Legacy Roads and Trails Program

Upgrading and maintaining needed forest roads and decommissioning unnecessary ones can maximize the many benefits they provide, while minimizing their risks to water quality and aquatic ecosystems

Roads provide many benefits to society, including access for recreation, forest management, fire management, and monitoring and research. Some road systems, however, can adversely impact water quality and the health of aquatic ecosystems in a variety of ways. These include increasing peak flows, erosion, and stream temperatures; constricting streams and decreasing their interactions with floodplains; and fragmenting habitats. Because the National Forest transportation system is so vast, was largely built using older standards, and contains many aging components, road restoration has been a major part of implementing the NWFP-ACS and similar aquatic strategies on federal forests throughout the Pacific Northwest. Restoration treatments generally include relocating roads away from streams, riparian areas, and unstable terrains; upgrading and improving road drainage systems; replacing road-stream crossings to provide passage for fish and other biota; “storing” roads that are not currently being used, but will be needed in the future; and decommissioning unneeded roads.

In recent years, Congress has placed greater emphasis on these activities and provided additional sources of funding. For example, between fiscal year (FY) 2008 and FY 2010, the U.S. Forest Service (FS) Pacific Northwest Region was allocated almost \$40 million in Legacy Roads and Trails funding to correct or reduce road and trail impacts and risks to watershed and aquatic



Figure 1. Field crews evaluate a decommissioned road.

resources. Given the importance of these activities in protecting and restoring watersheds, the FS is implementing two regional-scale studies to evaluate their effectiveness, which are described below. These finer-scale, more intensive effectiveness monitoring studies are intended to complement the coarser, broader-scale status and trend monitoring being conducted by AREMP.

Hydrogeomorphic Effectiveness Monitoring—To evaluate the effectiveness of road treatments in reducing hydrologic and geomorphic impacts, the FS Rocky Mountain Research Station and Pacific Northwest Region are using detailed, field-based inventories and a suite of robust environmental models to develop and compare multiple road impact-risk metrics, before and after road treatments at treated and control sites (fig. 1). Each site also includes a final validation evaluation following a large storm event. To date, evaluations have been initiated or completed at 25 locales where road decommissioning, road storage, or road drainage improvements have been or will be implemented. Results from the one site where monitoring results have been fully analyzed indicate that some road treatments can significantly reduce road impacts and risks to aquatic ecosystems.

Fish Passage Effectiveness Monitoring—Since 2002, the Region has treated more than 200 crossings using the “stream simulation” technique. This is a design process intended to ensure that conditions within a culvert or bridge mimic the natural upstream and downstream conditions. Fish passage, sediment transport and debris conveyance within the crossing are designed to function as they would in a natural channel. Given the considerable past and planned investments in removing fish barriers, the Region initiated a pilot program in 2008 to develop and apply a cost-effective method for evaluating whether new crossings are providing fish passage and simulating natural stream channels.


Preliminary results suggest that passage was achieved at 100% of the 25 monitored crossings and that 75% of them “simulated” the natural stream channel conditions. The Region is now working in partnership with the FS San Dimas Technology Center, the Pacific Northwest Research Station and the U.S. Geological Survey as part of a national effort to further refine and apply the monitoring protocol.

Contact Brian Staab brianstaab@fs.fed.us for more information


Methow Valley Subbasin, Okanogan and Wenatchee National Forest, Washington

Improving stream connectivity for fluvial bull trout in the Methow subbasin—The Methow Valley subbasin is a core bull trout (*Salvelinus confluentus*) area in the Upper Columbia River Basin. Bull trout are listed in the Endangered Species Act as a threatened species. A key criterion in the bull trout recovery plan is removing artificial barriers to allow unimpeded access to these important life stage areas.

Historically, irrigation dams and road culverts throughout the Methow Valley contributed to the decline in bull trout in the subbasin by blocking migratory corridors and restricting connectivity to upstream spawning areas and downstream overwintering areas.




Bull trout distribution showing restored or improved habitat connectivity.



Recent radio telemetry studies documented bull trout use in Goat Creek, Cold Creek, and Libby Creek, where access was previously blocked by artificial barriers.

Restoring migratory bull trout passage throughout the Methow was a large cooperative effort that included the Bureau of Reclamation, Washington (WA) Department Fish and Wildlife, Okanogan County, WA Department of Transportation, Methow Salmon Recovery Foundation, Chewuch Basin Council, private landowners and the Okanogan and Wenatchee National Forest. These efforts have greatly improved connectivity and migration corridors for bull trout in the Methow subbasin by restoring or improving migratory bull trout access to approximately 112 miles of habitat within the Methow subbasin.



Fluvial bull trout redd numbers from six streams monitored since 1999. The data do not suggest a clear trend. From 1999 to 2003, the redd counts fluctuated downward then had a somewhat consistent increase until 2007. In 2008 and 2009, bull trout redd counts were down across the Methow. Although the data does not demonstrate any clear trends, we expect the improved access has improved production.

Contact Gene Shull (gshull@fs.fed.us) for more information.

Paradise Creek, Coos Bay BLM District, Oregon

This creek needs help—Paradise Creek, a tributary to the Umpqua River, was designated as a tier one key watershed under the Northwest Forest Plan. However, Oregon Department of Fish and Wildlife (ODFW) habitat inventories showed the majority of the fish-bearing habitat throughout the watershed was in poor to fair condition because instream habitat was dominated by bedrock with minimal structure or stream complexity.

A partnership to the rescue—The Coos Bay District Bureau of Land Management (BLM) and partners Roseburg Resources Inc. (a private timber company), ODFW, the Partnership for Umpqua Rivers (watershed council) and private landowners (Paradise Creek Ranch) implemented a watershed-scale Paradise Creek restoration project during 2006-2007 affecting about 11 stream miles. It was apparent that achieving properly functioning conditions in the Paradise Creek watershed was not likely to occur within the foreseeable future without active intervention. Like the majority of forested lands managed by the Coos Bay District BLM, the Paradise Creek watershed has a legacy of logging and other land management activities that have degraded instream and riparian habitats on both public and private



Dan Van Slyke

lands. Roads now occupy a significant amount of historical flood plains, the conifers adjacent to stream channels have been reduced substantially, and logs and boulders were removed from virtually all stream reaches accessible by roads or logging systems.

Immediate benefits—

Adding large trees to the stream channel immediately benefited fish, amphibians,

and crayfish because the logs and boulders provided a tremendous amount of cover habitat that did not exist previously. The structures also provided immediate low-velocity refuge during high flows, which is critical for over-wintering juvenile salmonids. Post-project monitoring has shown that fine stream substrate is already being stored behind placed structures, and through time, stable gravel riffles will develop and provide quality spawning habitat. Spawning Chinook and coho salmon were observed in the boulder placement project reach in the first winter following placement.

Contact Dan VanSlyke (dan_vanslyke@blm.gov) for more information.



Jen Feola - ODFW

A helicopter, excavator, and cable yarding system were used to place over 900 conifer logs and hundreds of boulders throughout the watershed in 2006-2007 to provide quality spawning and rearing habitat for coho salmon (listed as threatened), Chinook salmon, cutthroat trout, steelhead trout, and both resident and Pacific lamprey. An average of 82 logs were placed per stream mile!

Scott River, Klamath National Forest, California

The 1.9-million-acre Klamath National Forest administers public lands that contain portions of the Klamath River and three of its significant tributaries: Scott, Salmon, and Shasta Rivers. A key beneficial use for these rivers is the coldwater salmonid fisheries for Chinook salmon, coho salmon, and steelhead.

Tributaries are considered impaired under the Clean Water Act

—All of these rivers are listed under the Clean Water Act as impaired for either nonpoint sources of temperature or sediment (303(d) listings). The advent of 303(d) listings and the subsequent development of Total Daily Maximum Loads (TMDLs) has placed a new urgency and focus on watershed restoration. Many TMDL requirements are consistent with past restoration efforts to disconnect the road network from the tributaries. For example:

- Unneeded roads were decommissioned. Old sediment-producing roads were converted from having inboard ditches to being out-sloped and drained by rolling dips.
- The amount of fill on many stream crossings was greatly reduced, which eliminated the potential for the streams to flow down the road.

However, TMDLs also caused a shift in watershed focus from key watersheds identified by the NW Forest Plan to nonkey impaired watersheds. Also, the use of anecdotal information for some 303(d) listings indicated a need for



This retaining wall was installed on Bucher Road to stop sediment from a 2006 slide from entering the Scott River.

scientifically credible data to verify some listings and focus scarce funds on the truly impaired watersheds.

The forest leveraged the use of appropriated funds by partnering with state and federal agencies and nonprofit organizations to achieve much of this road work. For example, by “piggybacking” on earlier efforts the forest just received over \$550,000 of federal stimulus funds to stormproof 140 miles of road.

Scott River TMDL—The first TMDL was developed for the Scott River watershed (520,000 acres). The challenge for the Forest is that it only administers 37 percent of this watershed. It also contains significant agricultural lands, industrial forest lands, and grazing lands. Portions of the Scott River tributaries were also turned literally upside-down by gold dredging in the early 20th century. The forest and the North Coast Water Quality Control Board are in the process of creating a bilateral memorandum of understanding (MOU) that will guide TMDL implementation compliance. One important requirement of the MOU is the already ongoing process of sediment source inventory, prioritization, scheduling and implementation of corrective actions. The model used for the Scott River may be applied in the future as an integrated and comprehensive forestwide approach to all TMDLs.

Integrated monitoring approach—Another new approach coming out of the Scott River TMDL is the development of an integrated monitoring approach, to be applied forestwide. The monitoring plan includes protocols for field collection of instream sediment data and stream shading. The plan also stratifies the forest’s seventh-field field watersheds into geologic (sandy, nonsandy, volcanic) and management categories (reference, over cumulative watershed effect threshold, sediment control, etc.) to facilitate analysis and interpretation of results. The monitoring plan objectives include:

- Track compliance with the TMDL.
- Conduct instream “best management practices” effectiveness monitoring.
- Ensure compliance of projects enrolled under categorical waiver(s).
- Create lines of evidence for the forest to use in proposing the delisting of unimpaired streams.
- Create data that can be used to calibrate and refine cumulative watershed effect (CWE) models.

Contact John Schuyler (jschuyler@fs.fed.us) for more information.

Six Rivers National Forest, California

Water quality problems—Most watersheds on the Six Rivers National Forest (NF) are listed as water quality impaired under section 303(d) of the Clean Water Act for sediment, temperature, nutrient, or dissolved oxygen. With the exception of sediment, these water quality concerns are primarily attributable to management activities outside of the forest boundary. On the Six Rivers NF, roads are the leading source of management-related sediment inputs. Where forest roads are located on steep terrain, mass soil movement is a common mechanism of erosion and sediment delivery. Most road-related erosion and sediment delivery is associated with large storm events that trigger culvert failures, stream diversions, and mass wasting. With the decline of road maintenance funding over the last 15 years, the risk of road failures and elevated sediment delivery is increasing, especially during large storm events.

Watershed restoration efforts for the purpose of protecting important anadromous streams, began about 1990 and were focused on minimizing surface erosion from inner gorge landslides by using tree planting and other landslide stabilization techniques. However,

these efforts proved costly and not very effective. When the Northwest Forest Plan was signed and subsequently incorporated into our forest plan in 1995, decommissioning of abandoned forest roads became the focus of restoration efforts on the Six Rivers NF.

Since 1994, the Six Rivers NF has secured over \$6.8 million dollars for road decommissioning and stormproofing (primarily correcting stream crossing diversion potential). To date, 384 miles have been decommissioned and 301 stream crossings have had diversion potential corrected. The forest received over \$1 million in 2009 for legacy road funding that will be spent on road-associated water quality improvements and road decommissioning.

Tribal partnerships are a key for success—

Forming partnerships between the Six Rivers NF and the Karuk and Yurok Tribes has proven to be an effective method of garnering competitive grant funding to restore fisheries habitat through road decommissioning effects. As part of the cost-share partnership, each Tribe contributes 30 to 35 percent of project costs. In addition to contributing funding, the tribes provide skilled restoration specialists to implement the work, thereby keeping jobs within local communities.

Contact Corrine Black (cblack@fs.fed.us) for more information.



Left Photo: Karuk Tribes restoration crews marked the limit of excavation and reviewed site plans prior to decommissioning this perennial stream crossing in the Bluff Creek watershed. Middle photo: View of completed work. The arrow marks the spot where the left photo was taken. Right photo: Closeup view of newly excavated stream channel.