

US Department of Agriculture Forest Service  
Pacific Northwest Research Station  
Olympic Peninsula Spotted Owl Demography Study  
2014 Annual Report  
January 2015

**Title**

Demographic characteristics of northern spotted owls (*Strix occidentalis caurina*) in the Olympic National Forest, Washington, 1987–2014.

**Principal investigator**

Eric Forsman, USDA Forest Service, Pacific Northwest Research Station, Corvallis, Oregon.

**Biologists**

Debaran Kelso, Senior Faculty Research Assistant; Jim Swingle, Faculty Research Assistant; and Kari Williamson, Biological Science Research Technician; Department of Fisheries and Wildlife, Oregon State University.

**Study objectives**

The primary study objectives are to estimate population trends, population age structure, reproductive rates, and survival rates of spotted owls in the Olympic Peninsula Demography Study Area. A secondary study objective is to estimate changes in occupancy of barred owls (*Strix varia*) within territories that were historically occupied by spotted owls.

**Potential benefits and utility of the study**

The Olympic Peninsula spotted owl demography study consists of parallel efforts conducted by the USDA Forest Service Pacific Northwest Research Station and by the Olympic National Park. This document reports the results of surveys conducted by the Pacific Northwest Research Station. Gremel (2014) reported the results from the Olympic National Park surveys. The Olympic Peninsula demography study is 1 of 8 long-term demographic studies that constitute the federal monitoring program for the northern spotted owl under the Effectiveness Monitoring Program for the Northwest Forest Plan (Lint et al. 1999). These demographic studies are designed to monitor vital rates and populations trends of spotted owls on federal lands in the Pacific Northwest. During regional meta-analyses that occur every 5 years, data collected from these demographic studies are used to make inferences regarding the effects of biological covariates on vital rates and population trends of spotted owls (Forsman et al. 1996, Franklin et al. 1999, Anthony et al. 2006, Forsman et al. 2011).

## **Study area**

The 94,800 ha Pacific Northwest Research Station study area is located on the northern Pacific and Hood Canal Districts of the Olympic National Forest and includes 40 sites that have been surveyed each year for spotted owls since 1991 (Fig. 1). The adjacent area surveyed by the National Park Service includes 128,000 ha within Olympic National Park (Gremel 2014; Fig. 2).

## **Methods**

Field and data collection methods used in this study have been described in a variety of sources (Franklin et al. 1996, Reid et al. 1999, Lint et al. 1999, Anthony et al. 2006). Study areas are surveyed  $\geq 3$  times each year using standardized protocols to estimate nesting status, and reproductive success of all territorial spotted owls detected within the area (Franklin et al. 1996, Lint et al. 1999). All owls detected are marked with color bands and US Fish and Wildlife Service numeric bands so that they can be identified visually, without trapping them every year. Capture histories of banded owls are used to estimate apparent annual survival, and apparent survival and recruitment are used to estimate the rate of population growth (Pradel 1996, Anthony et al. 2006, Forsman et al. 2011).

All spotted owl surveys and captures are conducted under US Fish and Wildlife Service recovery permit TE-026280-11, Washington State Scientific collection permit 10-139, master banding permit 21249, and Oregon State University's Animal Care and Use Permit 3628.

## **Results**

During the 2014 field season, we conducted 231 visits to 40 historical survey areas (mean survey visits per site =  $6.0 \pm 0.38 SE$ ). Spotted owl pairs were detected at 5 sites (Tables 1–2). We detected spotted owls at 3 other sites during night surveys. Follow-up visits at the latter sites indicated that the spotted owl detections were from paired individuals that had been previously documented at adjacent sites. Of the 10 adult spotted owls detected in 2014, 6 were previously banded and 4 (2 male, 2 females) were banded for the first time (Table 3). The total number of territorial owls detected in 2014 decreased slightly from 2013 and was still far below historical levels (Table 1). In contrast, the number of survey areas in which we detected barred owls remained high (Fig. 3).

All 5 females located in 2014 attempted to nest (Table 4) and 4 of the 5 females that nested produced young (Table 5). We documented 7 juveniles that successfully left the nest and 1 juvenile that was found dead at the base of a nest tree.

## **Discussion**

As the total number of territorial adults detected on the study area in 2014 decreased slightly from 2013, the number of owls detected on the study area remained far below historical levels. This was the first year in which significant numbers of spotted owls nested on the study area

since 2010. Although this is a positive sign, the generally low level of reproduction in most years since 2010 is resulting in an aging population.

### **Problems encountered**

There were no new problems encountered in 2014. Access to sites continued to be an issue because of road closures, and reduced road and trail maintenance.

### **Publications, presentations, and technology transfer completed in 2014**

Data collected in our study were used in a range-wide meta-analysis of spotted owl populations that was conducted in January 2014, in Corvallis, Oregon. The results of the meta-analysis will be published in 2015.

### **Duration of study**

Study was initiated in 1987 and will continue in 2015.

### **Acknowledgements**

This study was funded by the USDI Bureau of Land Management, USDA Forest Service Region 6, and the USDA Forest Service Pacific Northwest Research Station. We would like to thank Brian Biswell for his many years of service on the Olympic Peninsula study. Before he retired in January 2014, Brian was the crew leader on the study and managed and analyzed the data from 1996–2013. He will be missed!

### **Literature cited**

- Anthony, R.G., E.D. Forsman, A.B. Franklin, et al. 2006. Status and trends in demography of northern spotted owls, 1985–2003. Wildlife Monographs No. 163.
- Forsman, E.D., S. DeStefano, M.G. Raphael, and R.J. Gutiérrez, eds. 1996. Demography of the northern spotted owl. Studies in Avian Biology No. 17.
- Franklin, A.B., D.R. Anderson, E.D. Forsman, K.P. Burnham, and F.W. Wagner. 1996. Methods for collecting and analyzing demographic data on the northern spotted owl. Pp. 12–20 in E.D. Forsman, S. DeStefano, M.G. Raphael, and R.J. Gutiérrez, eds. Demography of the northern spotted owl. Studies in Avian Biology No 17.
- Forsman, E.D., R.G. Anthony, K.M. Dugger, et al. 2011. Population demography of northern spotted owls. Studies in Avian Biology No. 40.
- Franklin, A.B., K.P. Burnham, G.C. White, et al. 1999. Range-wide status and trends in northern spotted owl populations. Unpublished report. Colorado Cooperative Fish and Wildlife Research Unit.

- Gremel, S. 2014. Spotted owl monitoring in Olympic National Park: 2014 annual report. Unpublished report. USDI National Park Service, Olympic National Park, Port Angeles, WA.
- Lint, J., B. Noon, R. Anthony, E. Forsman, M. Raphael, M. Collopy, and E. Starkey. 1999. Northern spotted owl effectiveness monitoring plan for the Northwest Forest Plan. USDA Forest Service, Pacific Northwest Research Station, General Technical Report PNW-GTR-440.
- Pradel, R. 1996. Utilization of capture-mark-recapture for the study of recruitment and population growth rate. *Biometrics* 52:703–709.
- Reid, J.A., R.B. Horn, and E.D. Forsman. 1999. Detection rates of spotted owls based on acoustic-lure and live-lure surveys. *Wildlife Society Bulletin* 27:986–990.

Table 1. Number of spotted owls detected per year on the USDA Forest Service Pacific Northwest Research Station Olympic Peninsula Demography Study, 1987–2014. Age codes indicate adult, subadult, or age unknown spotted owls.

Year	Sites	Males			Females			Total
		Adults	Subadults	Age unknown	Adults	Subadults	Age unknown	
1987	14	10	0	1	9	0	2	22
1988	20	11	2	2	12	0	1	28
1989	28	22	0	1	16	0	3	42
1990	36	20	2	2	26	0	2	52
1991	40	27	1	2	25	1	3	59
1992	40	29	3	2	30	2	1	67
1993	40	27	3	2	27	0	5	64
1994	40	27	0	6	30	1	1	65
1995	40	28	0	3	24	0	1	56
1996	40	26	1	2	26	0	0	55
1997	40	25	0	1	20	1	4	51
1998	40	26	1	3	22	1	4	57
1999	40	10	0	2	10	0	1	23
2000	40	21	1	0	13	0	3	38
2001	40	13	0	5	17	0	1	36
2002	40	14	0	5	11	0	2	32
2003	41	13	0	3	8	0	3	27
2004	40	13	0	1	12	1	2	29
2005	40	12	0	0	11	2	0	25
2006	40	8	0	1	6	0	1	16
2007	40	9	0	2	4	0	4	19
2008	40	11	0	0	10	0	3	24
2009	40	6	0	0	3	1	1	11
2010	40	5	1	2	6	0	2	16
2011	40	3	0	3	4	0	2	12
2012	40	3	0	3	4	0	3	13
2013	40	4	1	0	5	0	2	12
2014	40	5	0	0	5	0	0	10

Table 2. Number of spotted owl territories on the USDA Forest Service Pacific Northwest Research Station Olympic Peninsula Demography Study in which we located pairs, singles, floaters, or no spotted owls, 1987–2014.

Year	Sites	Pairs	Singles	Floaters	Status uncertain	No owls
1987	14	9	3	0	0	2
1988	20	12	3	0	0	5
1989	29	19	4	0	0	6
1990	36	23	5	0	0	8
1991	40	24	5	2	2	7
1992	40	32	2	0	0	6
1993	40	28	6	0	0	6
1994	40	30	2	1	1	6
1995	40	22	9	0	1	8
1996	40	26	3	0	0	11
1997	40	20	6	1	1	12
1998	40	23	6	1	0	10
1999	40	6	9	0	1	24
2000	40	14	8	0	0	18
2001	40	15	4	1	2	18
2002	40	13	3	0	3	21
2003	40	8	6	0	4	22
2004	40	13	0	0	1	26
2005	40	11	2	0	1	26
2006	40	7	2	0	0	31
2007	40	4	7	0	3	26
2008	40	9	3	0	2	26
2009	40	3	2	0	3	32
2010	40	6	3	0	1	30
2011	40	2	4	0	3	31
2012	40	5	2	2	0	31
2013	40	4	3	0	0	33
2014	40	5	0	0	0	35

Table 3. Number of spotted owls banded on the USDA Forest Service Pacific Northwest Research Station Olympic Peninsula Spotted Owl Demography Study, 1987–2014. Non-fledglings are classified as adults ( $\geq 3$  years) and subadults (S1 = 1 year and S2 = 2 years).

Year	Fledglings	Males			Females			Total
		Adults	S1	S2	Adults	S1	S2	
1987	0	15	2	1	15	0	0	33
1988	13	11	1	3	13	0	0	41
1989	46	22	1	0	25	0	1	95
1990	62	19	6	3	22	1	7	120
1991	31	17	5	3	15	2	2	75
1992	78	23	1	2	21	0	1	126
1993	0	15	1	1	12	1	1	31
1994	32	8	1	1	11	1	1	55
1995	0	13	3	1	2	0	0	19
1996	58	5	0	2	9	0	3	77
1997	25	2	0	1	6	1	0	35
1998	26	2	1	1	4	2	0	36
1999	0	0	0	0	1	0	0	1
2000	1	6	0	0	5	0	0	12
2001	26	2	1	0	7	1	0	37
2002	28	1	1	0	4	0	0	34
2003	0	5	1	0	1	1	0	8
2004	36	6	0	0	5	1	0	48
2005	1	1	2	0	3	3	3	13
2006	6	0	0	0	0	0	0	6
2007	0	1	0	0	1	0	0	2
2008	11	2	0	0	3	0	0	16
2009	0	0	0	0	0	0	1	1
2010	0	0	0	1	0	0	0	1
2011	1	0	0	0	1	0	0	2
2012	1	0	0	0	1	0	0	2
2013	0	0	0	0	0	0	0	0
2014	5	2	0	0	2	0	0	9

Table 4. Annual reproductive statistics for female spotted owls from the USDA Forest Service Pacific Northwest Research Station Olympic Peninsula Demography Study, 1987–2014. Sample size only includes females where protocols for nesting status or the number of young produced where met.

Year	Proportion of females that nested			Proportion of females that produced young			Proportion of nesting females that produced young		
	n	$\pi$	95% C.I.	n	$\pi$	95% C.I.	n	$\pi$	95% C.I.
1987	16	0.19	0.00–0.38	19	0.11	-0.03–0.24	3	0.67	0.13–1.20
1988	19	0.26	0.07–0.46	27	0.33	0.16–0.51	5	1.00	—
1989	20	0.40	0.19–0.61	39	0.67	0.52–0.81	8	1.00	—
1990	35	0.71	0.56–0.86	52	0.56	0.42–0.69	24	0.63	0.43–0.81
1991	46	0.41	0.27–0.56	53	0.34	0.21–0.47	19	0.79	0.61–0.97
1992	48	0.90	0.81–0.98	63	0.78	0.68–0.88	43	0.86	0.76–0.96
1993	51	—	—	54	—	—	0	—	—
1994	49	0.84	0.73–0.94	56	0.54	0.41–0.67	41	0.66	0.51–0.80
1995	35	—	—	36	—	—	0	—	—
1996	37	0.89	0.79–0.99	50	0.68	0.55–0.81	33	0.67	0.51–0.83
1997	34	0.50	0.33–0.67	45	0.36	0.22–0.50	17	0.76	0.56–0.97
1998	43	0.56	0.41–0.71	45	0.42	0.28–0.57	24	0.71	0.53–0.89
1999	10	—	—	12	—	—	0	—	—
2000	25	0.12	-0.01–0.25	30	0.03	-0.03–0.10	3	0.33	-0.20–0.87
2001	31	0.55	0.37–0.72	34	0.44	0.27–0.61	17	0.88	0.73–1.04
2002	29	0.76	0.60–0.91	30	0.50	0.23–0.54	22	0.68	0.49–0.88
2003	26	—	—	26	—	—	18	—	—
2004	32	0.78	0.64–0.93	32	0.75	0.60–0.90	25	0.84	0.70–0.98
2005	29	0.03	-0.03–0.10	29	0.03	-0.03–0.10	29	0.03	-0.03–0.10
2006	8	0.88	0.65–1.10	9	0.67	0.36–0.98	8	0.75	0.45–1.05
2007	7	—	—	0	—	—	0	—	—
2008	4	0.50	0.01–0.99	9	0.78	0.51–1.05	4	0.50	0.01–0.94
2009	6	—	—	6	—	—	0	—	—
2010	5	0.80	0.45–1.15	5	—	—	5	—	—
2011	4	—	—	4	—	—	0	—	—
2012	6	—	—	6	0.33	-0.04–0.71	5	0.2	-0.15–0.55
2013	2	—	—	0	—	—	0	—	—
2014	5	1.00	—	5	0.80	0.45–1.15	5	0.80	0.45–1.15

Table 5. Estimated fecundity ( $\beta \pm SE$ ) of female spotted owls on the USDA Forest Service Pacific Northwest Research Station Olympic Peninsula Demography Study, 1987–2014. Fecundity is defined as the number of female young produced per female, assuming a 50:50 sex ratio of offspring.

Year	Adults	Subadults	Adults	Subadults	Combined <sup>1</sup>
			$\beta \pm SE$	$\beta \pm SE$	$\beta \pm SE$
1987	18	1	0.08 ± 0.06	—	0.08 ± 0.06
1988	25	2	0.24 ± 0.08	—	0.24 ± 0.07
1989	39	0	0.54 ± 0.07	—	0.54 ± 0.07
1990	46	5	0.47 ± 0.07	0.10 ± 0.10	0.42 ± 0.06
1991	50	3	0.31 ± 0.06	0.17 ± 0.17	0.30 ± 0.06
1992	57	6	0.66 ± 0.05	0.50 ± 0.18	0.64 ± 0.05
1993	49	0	—	—	—
1994	53	1	0.42 ± 0.06	—	0.39 ± 0.06
1995	36	0	—	—	—
1996	43	3	0.56 ± 0.07	0.33 ± 0.17	0.54 ± 0.06
1997	43	0	0.31 ± 0.07	—	0.30 ± 0.06
1998	39	3	0.31 ± 0.07	0.50 ± 0.29	0.31 ± 0.06
1999	11	0	—	—	—
2000	29	0	0.02 ± 0.02	—	0.02 ± 0.02
2001	33	0	0.36 ± 0.08	—	0.38 ± 0.08
2002	28	0	0.45 ± 0.09	—	0.45 ± 0.08
2003	22	1	—	—	—
2004	23	4	0.74 ± 0.08	0.38 ± 0.24	0.60 ± 0.08
2005	22	5	0.02 ± 0.02	—	0.02 ± 0.02
2006	8	0	0.50 ± 0.16	—	0.50 ± 0.14
2007	7	0	—	—	—
2008	9	0	0.63 ± 0.16	—	0.61 ± 0.14
2009	5	1	—	—	—
2010	5	0	—	—	—
2011	3	0	—	—	—
2012	6	0	0.17 ± 0.11	—	—
2013	6	0	—	—	—
2014	5	0	0.70 ± 0.20	—	0.70 ± 0.20

<sup>1</sup>Includes 14 females of unknown age that were not reported in this table.




Figure 1. USDA Forest Service Pacific Northwest Research Station spotted owl demography study area, Washington. The dark box in the lower left insert indicates the study area. Black dots indicate spotted owl sites that were surveyed from 1987–2014.




Figure 2. Olympic National Park spotted owl demography study area, Washington. The dark box in the lower left insert indicates the study area. Black dots indicate spotted owl sites that were surveyed from 1992–2014.




Figure 3. Proportion of monitored owl territories with detection of  $\geq 1$  spotted owls (gray dots) and  $\geq 1$  barred owl (black dots) on the USDA Forest Service Pacific Northwest Research Station Olympic Peninsula spotted owl demography study, 1987–2014.